

**SNAPSHOTS OF SYSTEMS - METABOLIC CONTROL ANALYSIS AND BIOTECHNOLOGY IN
THE POST-GENOMIC ERA**

by

^{1,*}Douglas B. Kell & ^{1,2}Pedro Mendes

¹Institute of Biological Sciences, Cledwyn Building, University of Wales, Aberystwyth SY23 3DD, UK

²Present address: National Center for Genome Resources, 1800 A Old Pecos Trail, Santa Fe, New Mexico
87505, USA

Corresponding author: phone +44 1970 622334 fax +44 1970 622354

email dbk@aber.ac.uk

<http://gepasi.dbs.aber.ac.uk/home.htm>

Institute of Biological Sciences, Cledwyn Building, University of
Wales, Aberystwyth SY23 3DD, UK

SUMMARY

We provide a critical survey of some of the major limitations of the principles and applications of metabolic control analysis, with special reference to the enhancement of fluxes of biotechnological interest. Experimental methods of single-cell analysis such as flow cytometry show that the implicit assumption that we study and model ensembles of identical cells is often completely untenable, and that cellular heterogeneity is much greater than we normally assume. A feature of the post-genomic era is the recognition that many more genes exist, and are expressed, than we had ever recognised, and that methods are being developed for the quantitative assessment of this. Even if the individual flux-control coefficients of these gene products were each very small their enormous number means that over-simplified analyses that ignore them are very likely to lead to erroneous analyses of the true structure and organisation of a metabolic system or subsystem of interest. Even the assumption that the components of moiety-conserved cycles do not change their total concentrations during an experiment ignores what may be a large impact of this on the control structure. In the absence of compartmentation or channelling, such cycles also serve to connect segments of metabolism usually considered rather distant from each other. Simplified ('top-down') methods in which the system structure is assumed *a priori* often will not work to give unequivocal answers for complex systems where the combinatorial explosion of possible interactions requires much more sophisticated methods for system identification. Dual-inhibitor titrations can reveal unsuspected direct kinetic interactions between individual catalytic activities in appropriate cases, but these are cleanly apparent only in the regime of large changes (such that experimental studies in which only small perturbations are revealed will cause (or allow) them to be missed). No example exists in which one can extrapolate the conventional control coefficients to provide reliable and quantitative predictions *a priori* about the behaviour of metabolic systems subjected to 'large' changes in parameters, although these may well be a valuable guide. A "Universal Method" proposed for solving this suffers from technical problems such as being unable to manipulate fluxes at constant metabolite concentrations. It is thus unclear if there are situations in which it could be applied in practice (especially if there are substantial interactions *between* pathways, usually involving conserved moieties such as adenine and pyridine nucleotide couples).

Metabolic control analysis and functional genomics share the same agenda in that they seek to relate the presence and activities of individual genes and gene products to higher level processes of cellular biochemistry and physiology. They can be considered to differ, however, in a philosophical sense since the former is essentially deductive in character (and as practised, although some of us theorems were initially captured by induction) while the latter is of necessity inductive, at least initially, because so many ORFs are of unknown function. Inductive methods of machine learning applied to large-scale data from analyses of the transcriptome, the proteome and the metabolome should prove of value in unravelling their properties.

SNAPSHOTS OF SYSTEMS - METABOLIC CONTROL ANALYSIS AND BIOTECHNOLOGY IN THE POST-GENOMIC ERA

“So the first requirement will be for a theoretical framework in which to embed all the detailed knowledge we have accumulated, to allow us to compute outcomes of the complex interactions and to start to understand the dynamics of the system. The second will be to make parallel measurements of the behaviour of many components during the execution by the cell of an integrated action in order to test whether the theory is right. Is there some other approach? If I knew it I would be doing it, and not writing about the problem.”

Sydney Brenner, 1997, in *Loose Ends* publ. Current Biology, London, p.73

“But one thing is certain: to understand the whole you must look at the whole”

Kacser, H. (1986). On parts and wholes in metabolism. In *The organization of cell metabolism* (ed. G. R. Welch and J. S. Clegg), pp. 327-337. Plenum Press, New York.

Introduction - Holism and reductionism, and MCA's view of the operations of complex biochemical systems

Following its original formulation in 1973 (Heinrich & Rapoport, 1973; Heinrich & Rapoport, 1974; Kacser & Burns, 1973) as a means of understanding the contribution of the individual steps of a biochemical pathway to the values of flux and metabolite concentrations observed, some 13 years were to pass before we first surveyed (Kell & Westerhoff, 1986a; Kell & Westerhoff, 1986b) how the formalism, tools and terms of metabolic control analysis might usefully be applied to such systems in a biotechnological context. Since another such **period** has now elapsed, it is timely to take stock of progress, to recognise that the take-up of these methods among biotechnologists has been less than widespread, and (as requested by the Editor) to give a personal and critical review of successes, failures, problems and prospects for the use of MCA in biotechnology. In what follows, it is taken that the reader has a good working knowledge of the essential principles of MCA, as summarised for instance in (Cornish-Bowden & Cárdenas, 1990; Fell, 1992; Fell, 1996; Heinrich & Schuster, 1996; Kell, van Dam & Westerhoff, 1989; Kell & Westerhoff, 1986a; Ovádi, 1995) and on the Internet at http://gepasi.dbs.aber.ac.uk/metab/mca_home.htm and in links therefrom. In addition, we shall concentrate on unicellular systems, implicitly those most commonly exploited to make products of biotechnological interest.

Perhaps the chief intellectual benefits of MCA have been the recognition (i) that in the steady state of a (linear) pathway where all steps are proceeding at the same rate it is nevertheless appropriate to recognise that each contributes quantitatively to the control of flux, in a manner which (for small or infinitesimal changes) can be summed to unity, (ii) that the flux-control coefficients so determined are consequently necessarily small, and (iii) that the activities of many steps must be changed simultaneously if fluxes are to be enhanced substantially. MCA thus constituted a bridge between the rather reductionistic view then prevalent (that we can understand a systems by looking at its component parts in isolation, without considering the *interactions* between them - see (Kell & Welch, 1991; Mendes, Kell & Welch, 1995)) and the holistic one (which *in extremum* - and in practice for many real, nonlinear, coherent, self-organising systems (Kell & Hitchens, 1983) - would hold that the whole is so much more than the sum of its parts that it is essentially pointless to consider the individual parts in isolation at all (Ho, 1998)).

Coupled to these aperçus has been the recognition that computer simulation can be a powerful tool in solving the forward problem of metabolism: given the parameters of the system (usually the external metabolite and effector concentrations and the kinetic properties of the enzymes) one can solve the relevant differential equations and predict the time course and - if such exist - the steady-state values of flux and metabolite concentrations. Software such as the program *Gepasi* produced in Aberystwyth by one of us (Mendes, 1993; Mendes, 1997; Mendes & Kell, 1998b) has been designed for (and indeed by) biologists (and successfully hides the mathematical details from the typical user), and given a simulation of a pathway it is easy to extract the ‘MCA properties’ such as flux- and concentration-control coefficients by numerical simulation of the effects of small changes in parameter values or analytically by differentiating the rate equations to acquire the elasticities and inverting the elasticity matrix so obtained (Fell, 1996; Fell & Sauro,

1985; Mendes & Kell, 1998a; Reder, 1988; Westerhoff & Kell, 1987). The metabolic control analysis of a system is thus normally 'merely' a snapshot of a typically rather restricted subset of the cellular biochemistry actually taking place in time and space.

Assumptions in MCA, implicit and explicit.

With its concentration on small or infinitesimal changes, a domain where for (spatially) homogeneous systems its analysis is both exact and complete, MCA necessarily represents an *approximation* to a more complex reality, and this begs the question of how adequate this approximation is. In view of the recognition that it is but a subset of a full simulation of whatever system it is desired to simulate, it is probably unsupportable in the general case. Some other common assumptions of MCA (and its usual implementations) which will be explored later are summarised in Table 1. They include the implicit view that all cells in a suspension are the same, that it is possible to lump together large segments of metabolism without losing important knowledge of the behaviour of the overall system (or at least the ability to discriminate the model from other 'competing' models), and that there are 'universal' methods which can permit the rational and practical optimisation of metabolic fluxes in systems of arbitrarily complex organisation.

All cells in an axenic culture are not the same; microbial differentiation

Whilst it is rather obvious that the phenotypes of all cells in a *differentiated* organism are not the same (so much so that there is no such thing as a biochemically "normal" individual (Williams, 1956)), it is implicit in a standard MCA analysis that they are; in other words we tend to treat the system under study as an ensemble in the thermodynamic sense (Welch & Kell, 1986; Westerhoff & van Dam, 1987). In fact, the essence of the problem (Kell *et al.*, 1991) is that one is trying, typically, to correlate a rate of change (v) of a certain variable with respect to the value of a certain parameter or property (p), but a correlation may be

expected between the mean values \bar{v} and \bar{p} *only* if v is kinetically of first order with respect to p . This is completely unrealistic even for the axenic microbial cultures that are the focus of this review, and with the availability of techniques such as flow cytometry (Davey & Kell, 1996; Kell *et al.*, 1991; Shapiro, 1995) it becomes possible to determine the heterogeneity of cellular properties *directly*. In one example of our own (Kaprelyants & Kell, 1992), the extent to which *chemostat-grown* (and thus as near as one can get to genuinely steady-state cultures of) *Micrococcus luteus* cells could take up the membrane energisation probe rhodamine 123 varied by more than 1000-fold under conditions in which uptake was fully uncoupler-sensitive and neither efflux pumps nor lack of membrane permeability were an issue. The unwanted consequence of the failure to take culture heterogeneity into account (in terms of being led to erroneous conclusions about causality and mechanism) reaches its apotheosis in the study of microbial viability/culturability (Kell *et al.*, 1998). Such an analysis ignoring heterogeneity will also tend to mask intercellular interactions (Fuqua, S.C. & Greenberg, 1996; Kell, Kaprelyants & Grafen, 1995) such as those in which culturable cells secrete a factor necessary for the resuscitation and growth of non-growing cells of the same organism (Kaprelyants & Kell, 1993; Kaprelyants, Mukamolova & Kell, 1994; Kaprelyants *et al.*, 1999; Mukamolova *et al.*, 1998). Note of course that in transitions to states such as dormancy and non-culturability we are here talking about what MCA would regard as 'large' changes.

Not all organisms are the same....

So far as the typical textbook of biochemistry is concerned, cells are inevitably taken to be essentially similar, with a great majority of their broad activities (and the 'housekeeping' genes which code for them) being common throughout biology (at least at the level of prokaryote, eukaryote and archaean). Specific features such as photosynthesis or nitrogen fixation are seen merely as occasional adjuncts. Our attempts to simulate metabolism seem to rely implicitly on this, and it is perhaps assumed that models have a validity beyond the system for which they are constructed. However, as we enter the post-genomic era, two major facts have become evident; (i) many or most ORFs code for products of unknown function (Blattner *et al.*, 1997; Bork *et al.*, 1998; Cole *et al.*, 1998; Goffeau *et al.*, 1996; Hinton, 1997; Oliver, 1996) with many being conserved but most comparatively unique (Koonin & Galperin, 1997), and (ii) large-scale, genome-wide comparisons of orthologous genes point up the prevalence of horizontal gene transfer (Forterre, 1997a; Forterre, 1997b; Koonin *et al.*, 1997; Rivera *et al.*, 1998) and the consequent inadequacy of gradualist views of evolution. (Note however that these analyses are to date restricted to the very small fraction (Amann,

Ludwig & Schleifer, 1995) of cultured microbes, and that many close relative of existing taxa remain to be cultured (Kaprelyants *et al.*, 1999; Kell *et al.*, 1998; McVeigh, Munro & Embley, 1996.) Consequently, we now recognise that many more genes contribute to fitness than had previously been considered.

The major approaches to functional genomics currently being undertaken involve the systematic knocking out of individual genes *seriatim*; where this is being done, e.g. in *S. cerevisiae* (Dujon, 1998; Oliver & Baganz, 1998; Oliver *et al.*, 1998; Teusink *et al.*, 1998), it is found that (“only”) some 15% are ‘essential’, and the question arises as to the role of the others - do they have a very high contribution to fitness under a restricted set of conditions met only occasionally and never in the laboratory, or do they all provide a marginal contribution to fitness? At least as judged by the fact that they are both transcribed under laboratory conditions in rich media and can be shown to contribute to fitness in sensitive growth rate tests (Eisen *et al.*, 1998; Smith, Botstein & Brown, 1995; Smith *et al.*, 1996; Thatcher, Shaw & Dickinson, 1998), one is led to attach significance to the latter view. The mental picture which emerges then is that whilst there may be core or major blocks of primary metabolism which are important, the contribution of the rest of the cellular activities which are normally neglected is likely to be just as great or greater. (Note here the point - shown up in some of the pioneering control analysis of mitochondrial respiration (Wanders *et al.*, 1984) - that the control structure, even for a given respiratory flux, depended enormously on the reaction (hexokinase vs creatine kinase) that used the ATP.) When we come to intact microbial cells, we must recognise that although these other interactions may be individually small they are collectively numerous (the totals of genes in *E. coli*, baker’s yeast and *Streptomyces coelicolor* A3(2) are some 4000, 6000 and 8000 respectively), and while fewer will contribute to a flux than to the overall fitness as correctly judged (Kell, 1987; Westerhoff, Hellingwerf & van Dam, 1983) by growth *rate*, the emerging paradigm is of a much greater complexity and sophistication of unicellular controls than we had heretofore recognised. One example of an important (and probably excessive) simplification in common usage is that the total values of pyridine nucleotides in cellular compartments are not of particular significance.

Moiety conservation and flux enhancement

One aspect of metabolism that has received comparatively little attention for the purposes of flux maximisation is the existence of moiety-conserved cycles (Reich & Sel’kov, 1981). These are ubiquitous in metabolism and a few of them (e.g. NAD/NADH and ATP/ADP) act as major links between various parts of metabolism and impose constraints on the behaviour of the system. Whilst affecting the poise of these cofactor couples has been shown to be useful in metabolic engineering (Lopez de Felipe *et al.*, 1998) the flux through a pathway, as seen within the MCA formalism (Hofmeyr, Kacser & van der Merwe, 1986), is also controlled by the *total amount* of conserved moieties. We have observed by computer simulation that the flux of several model pathways responds to the total amount of conserved moiety according to a bell-shaped curve. This suggests that for such systems there is an optimal amount of cofactor for a given pathway flux (and that compartmentation of pathways would be necessary to optimise them separately), otherwise the flux will be somewhat limited by the availability of the conserved moiety. To manipulate the total amount of the moiety we will thus need to target the pathways of their biosynthesis and degradation. Alternatively we could as well manipulate the number and/or affinity of moiety binding sites which modulate the amount of *available* (i.e. free) total moiety. Fig. 1 depicts such a bell-shaped relation between the total amount of the moiety and the flux for a simple branched pathway in which other parameters are held constant. As nicely shown by Rohwer, Olivier & Hofmeyr {?this volume?}, the addition of extra reactions to this scheme can change the shape of these curves significantly, reinforcing the importance of recognising that the control structure of simple systems may be changed dramatically when we embed them in more complex ones (such as a cell). Indeed one must be very careful not to discard too much detail about the system when constructing models. Others (Bakker, ?this volume?, Cornish-Bowden, ?this volume?) have presented model pathways in which the constraints imposed by mass conservation (in the parasite *Trypanosoma brucei*) can indeed be used to advantage in the design of drug therapies.




Figure 1. Dependence of entry steady-state flux on the total amount of conserved moiety in a model branched pathway. The pathway simulated is shown in the inset, the arrows representing the positive direction of flux. All reactions are fully reversible, all kinetic and equilibrium constants are unity, $[A] = 10$, $[C] = [D] = 1$ such that the overall disequilibrium ratio is 0.1 on both branches.

On modelling at the right scale

We have traditionally treated our systems relatively simply, due in part to the difficulty of measuring everything. With the emergence of measurements of the transcriptome (Bowtell, 1999; Brown & Botstein, 1999; Chu *et al.*, 1998; de Saizieu *et al.*, 1998; Debouck & Goodfellow, 1999; DeRisi, Iyer & Brown, 1997; Duggan *et al.*, 1999; Iyer *et al.*, 1999; Lipshutz *et al.*, 1999; Marton *et al.*, 1998; Schena *et al.*, 1996; Spellman *et al.*, 1998; Velculescu *et al.*, 1997; Wodicka *et al.*, 1997), the proteome (Anderson & Anderson, 1998; Blackstock & Weir, 1999; Boucherie *et al.*, 1996; Cash, 1998; Garrels, 1996; Humphery-Smith, Cordwell & Blackstock, 1997; Wang & Hewick, 1999; Wilkins *et al.*, 1996; Wilkins *et al.*, 1997) and the metabolome (Oliver & Baganz, 1998; Oliver *et al.*, 1998) we now have the ability to carry out hundreds of measurements on macromolecular and metabolic variables *simultaneously*. The outcomes of the pioneering studies are in many cases given in the form of lists of expression ratios for the hundreds of genes of interest, which are hard to interpret - the appropriate scale for easy understanding is not a life-sized model (Eisen *et al.*, 1998). Treating related segments of metabolism as 'blocks' is one solution (Kacser, 1983; Kell *et al.*, 1989), sometimes referred to as top-down analysis (Brand, 1996; Brand, 1998; Brown, Hafner & Brand, 1990), and is being exploited in functional genomics as the FANCY method (Oliver & Baganz, 1998; Teusink *et al.*, 1998), but this approach fails to give a true account of the system of interest under a number of circumstances, and in some cases may lead only to the system being underdetermined.

A critique of 'top-down' methods in which segments of metabolism are treated as one

In the early literature of MCA (Kacser & Burns, 1973) it was already proposed that one could group sequential enzymes and treat them as one unit for the purposes of control analysis. This is possible due to the summation theorem (Kacser & Burns, 1973) and the fact that the elasticity concept can be applied to groups of reactions (Kacser, 1983). Brown *et al.* (Brown *et al.*, 1990) took this one step further and formally proposed the 'top-down' method for determining control coefficients. In this method one builds two groups of metabolic steps around (upstream and downstream of) one single intermediate metabolite. Provided that this central intermediate metabolite is the only kinetic link between the two groups of steps then one can determine the elasticities of the two groups towards that metabolite with just two single-modulation experiments. Group control coefficients can then be calculated using the connectivity and summation

theorems (the assumption therefore being that no other interactions - such as feedback loops or metabolic channelling (Agius & Sherratt, 1997; Mendes *et al.*, 1995; Ovádi, 1995) - exist between the two groups). The control coefficients thus obtained for the two groups of steps could reveal which of them has more control, e.g. as in (Simpson, Shimizu & Stephanopoulos, 1998; Stephanopoulos & Simpson, 1997), where knowledge of this can allow one to vary the control structure of a pathway to improve fluxes of biotechnological interest (Simpson, Colon & Stephanopoulos, 1995).

An attractive application of this method would be to apply it repeatedly by subdividing each group into smaller groups. This would result in the determination of the complete control structure of the pathway, each time with more resolution (hence the 'top-down' name). Although elegant in conception, this method may be impossible to apply with impunity in practice, due to the requirement of grouping steps such that there is only one kinetic link (via the intermediate metabolite) between them. Ainscow and Brand have recently extended the method (Ainscow & Brand, 1995) such that it can be applied to the case when the groups of steps are connected by more than one metabolite. Nevertheless, for the method to work as intended one must be absolutely sure that all the kinetic links between the groups of steps are known and included in the analysis explicitly (or are unaffected by the modulations performed (Ainscow & Brand, 1998a)). Thus, one can never be sure that the coefficients determined by this method are correct as there could be extra kinetic interactions between the two groups of steps other than the ones taken into consideration. The classical method of direct determination of control coefficients by perturbation of enzyme activities is immune from this problem and so could (and should) be used to confirm the results with the top-down method - but (while this may be hard if the number of reactions in individual blocks is large) this rather defeats at least some of the purpose of using it in the first place! (Recently, Kholodenko and colleagues have presented a combination of top-down MCA and the perturbation method which they refer to as 'Metabolic Design Analysis' (Kholodenko *et al.*, 1998).) But there are extra problems when one wants to use this method in general: (i) there are several known (and certainly many more unknown) feedback loops in metabolism and (ii) many reactions include cosubstrates such as NAD/NADH or ATP/ADP which form kinetic links between steps normally considered distant. Both these reduce the number of metabolites that can be effectively used in the top-down approach to separate groups of steps, and as such the determination of whether the particular blocks chosen for the simplification are the most appropriate ones should be seen as a system identification or inverse (Mendes & Kell, 1996) problem. As such the method is not amenable, in general, to a true 'top-down' approach of measuring *all* control coefficients. The problems are greatly compounded by the propagation of errors and bias contingent on the measurement of elasticities (Schlosser, Holcomb & Bailey, 1993; Thomas & Fell, 1995) and control coefficients (Ehlde & Zacchi, 1996; Small, 1993), and such errors may not be normally distributed (Ainscow & Brand, 1998b). We therefore find that the (correct) application of the top-down method to large, complex biosystems may be rather limited in practice, and above all dangerous if results are not validated by an independent method. Indeed, the effect of error propagation can be such that it is almost impossible to falsify a chosen model (in the sense of discriminating it from a better model of metabolic flux and its control) within the limits of the experimental precision attainable, especially when the perturbations are held small (within the range in which elasticities are unchanged).

Note that this is not a critique of simplification *per se*, since in many cases the intrinsic dimensionality of the major blocks of a complex system of interest may well be comparatively small and the level of understanding that we require, and indeed good precision in our models (Broadhurst *et al.*, 1997; Kell & Sonnleitner, 1995; Shaw *et al.*, 1997), is more easily attained with small models than with large ones. But this is something that we find out afterwards, when measurements of many variables have been made and evaluated (Eisen *et al.*, 1998; Tamayo *et al.*, 1999), not something to build in beforehand! Hence our stress that these types of problem should first be seen as problems of system identification.

Proton-coupled electron transport-linked phosphorylation - an example of a channelled system, assessed using dual-inhibitor titrations

A consequence of lumping reactions together in the macroscopic way typified by the top-down approach is that it is assumed that their intermediates are delocalised. One of the major areas of interest of this laboratory has been in the problem of channelling, most recently in terms of intermediary metabolites (Mendes *et al.*, 1995; Mendes, Kell & Westerhoff, 1992; Mendes, Kell & Westerhoff, 1996) but more classically in terms of the problem of whether the energetic intermediates of electron transport-linked phosphorylation are delocalised or not (Kell, 1979; Kell, 1988; Kell & Westerhoff, 1990). The basic idea is

as follows. In the classical chemiosmotic coupling model (Mitchell, 1966; Nicholls & Ferguson, 1992), electron transport generates a transmembrane proton gradient which, due to the rapid diffusion rates of protons in aqueous media, leads to a delocalised protonmotive force Δp consisting of a membrane potential $\Delta\psi$ and a pH gradient $z\Delta\text{pH}$ which is consequently available to *all* ATP synthase enzymes in the organelle in whose membrane the pmf-generators are embedded. Uncouplers act by dissipating the pmf as heat. (In addition, the pmf can in principle feed back to inhibit electron transport via ‘slip’ or be dissipated ‘naturally’ to heat via a pmf-dependent ‘leak slip’ which does not differ formally from the imperfect coupling naturally present.) This is depicted in Fig 2.




Figure 2. The classical chemiosmotic coupling paradigm for electron transport-linked phosphorylation

The consequence is that the rate of phosphorylation J_p should depend only and monotonically on Δp , typically (when compared with the data that may be obtained experimentally in acid-bath-type experiments) according to a sigmoidal function of the pmf and with a threshold of ca 150 mV in which no phosphorylation occurs, i.e.

$$J_p = [\text{ATP-synthase}] \cdot f(\Delta p) \quad (\text{Eq. 1})$$

In a typical dual-inhibitor titration using an uncoupler and an ATP synthase inhibitor, we first study the effect of uncoupler on J_p , with results similar to those in Fig 3 (open circles). We then inhibit half of the ATP synthases using a tight-binding (or better covalent) inhibitor such that J_p falls to one half of its original value (Fig 3). The effect of the uncoupler titration that must be predicted from the delocalised chemiosmotic type of uncoupling model is similar to that given by the closed triangles in Fig 3, since the pmf cannot be made smaller by this treatment and it is probably slightly larger due to the smaller drain on it: the shape of the curve is the same but the rate at any level of uncoupler (and putatively pmf) is just one half of the control. Similarly, the amount of uncoupler needed to achieve full uncoupling is the same. Unfortunately for this view, the experimental curve is quite different: lowering the initial J_p to one-half of its original value with the ATP synthase inhibitor decreases by one half the amount of uncoupler necessary to achieve full uncoupling. *No delocalised coupling model can account for this type of behaviour in uncoupler/energy transfer inhibitor titrations (Herweijer, Berden & Slater, 1986; Hitchens & Kell, 1983a; Hitchens & Kell, 1983b; Kell, 1988; Kell, 1992; Westerhoff & Kell, 1988), and indeed none has made a serious attempt to do so.* Note, however (from Fig 3), that if we had carried out the experiment in the limit of small changes -not allowing the flux to change by more than say 5-10% - and included error bars as well, we might easily have allowed ourselves the conclusion that the titration curve (normalised to the flux in the absence of uncoupler) had been unchanged by the presence of the energy transfer inhibitor (and thus consistent with the delocalised model). *Only the extension to large changes gave a clear an unequivocal discrimination between the competing models, and the conclusion is that if we wish to test our models to destruction, rather than simply seeking to parametrise those we believe to be true, we must carry out large changes in the external parameters.*




Fig 3. Theoretical and experimental traces from dual uncoupler/energy transfer inhibitor titrations.

Between the Bud and the Rose: Large Changes and Metabolic Control Analysis

As is well-known, and is mentioned above, the theorems of MCA work only for parameter changes which are small (and in principle infinitesimal). This said, we should be reminded of the classical paper (Kacser & Burns, 1981) illustrating that we should expect genetic dominance to be rare (as is observed) due to the fact that only rarely (e.g. as in (Agius & Peak, 1997; Agius *et al.*, 1996)) is an individual flux-control coefficient sufficiently high to give a major change in phenotype when the amount of active enzyme decreases to 50% of the wild type. The necessary corollary of this is exactly that enzymes with low flux-control coefficients at wild-type levels should also be expected to have low flux-control coefficients at the lower enzymes concentrations, i.e. following large changes! Notwithstanding, it is to be assumed (and see below) that substantial increases in flux towards metabolites of biotechnological interest do require large changes in at least some of the parameters such as enzyme concentrations (and the same is true for the phenotypic manifestation of disease states when a threshold loss in enzymatic function is induced (Durrieu *et al.*, 1997; Letellier *et al.*, 1998; Mazat *et al.*, 1997; Mazat *et al.*, 1998)). Small and Kacser tackled the problem of exactly how great an inaccuracy in estimating control coefficients via large changes in parameters might be involved. They first introduced the idea of a *deviation index* as the relative change in a metabolic variable such as a flux to a *large* change in a parameter (Small & Kacser, 1993a), and showed that for unbranched chains of enzymes with linear kinetics there was a direct relationship between deviation indices and flux-control coefficients. They also pointed out that combined changes of the activity of individual enzymes will produce a more-than-additive response (and see below). The behaviour of branched and non-linear pathways was more complex (Small & Kacser, 1993b), and though it was stated that many metabolic systems behave in practice as quasi-linear systems, the differences between the actual and predicted amplification factors were often quite great. Indeed, a detailed study by Schuster & Holzhütter (Schuster & Holzhütter, 1995) of erythrocyte properties resulting from large-scale alterations in enzymatic activities concluded that no existing extrapolation method using the conventional control coefficients was able to provide reliable predictions.

The not-very Universal Method

It is worth noting that, as proven in the summation theorem of Metabolic Control Analysis (e.g. (Cornish-Bowden, Hofmeyr & Cárdenas, 1995; Fell, 1996; Heinrich & Rapoport, 1974; Heinrich & Schuster, 1996; Kacser & Burns, 1973; Kell & Westerhoff, 1986a)), changes in the concentrations of individual

enzymes tend to have little effect on particular metabolic fluxes (nor, indeed, on the gross phenotype under most laboratory conditions (Thatcher *et al.*, 1998)). However, in part because of the so-called connectivities of MCA, changes in individual enzyme concentrations can and do have substantial effects on metabolite *concentrations*, even when the changes in flux are negligible (Mendes *et al.*, 1995; Mendes *et al.*, 1996). It is therefore very reasonable that attempts to increase metabolic fluxes by increasing the concentrations of metabolic enzymes may lead to substantial increases in metabolite levels, and that these may either prove cytotoxic or at least necessarily lead to the diversion of flux to pathways other than that desired. It would therefore be desirable (if it were indeed possible) to seek to modulate fluxes by changing enzyme activities in a manner that managed to *preserve* the steady-state levels of metabolites. Thus Kacser & Acerenza (Kacser & Acerenza, 1993) introduced the so-called Universal Method that purported to have this effect and to be ‘entirely general’. In the Universal Method, it is recognised that in any pathway leading to the output of interest, the activities of whose enzymes one would wish to increase, there will be branch points leading to other parts of metabolism which should not be perturbed. Because of the conservation of mass, the fluxes down each branch point following a change in flux are related both to the changes in flux before and after the branchpoint in the pathway of interest and to the ratio r of enzyme activities before and after the change in flux. For unimolecular reactions, there is a unique value of r for each such reaction at which the fluxes down the branches remain unchanged.

However, the Universal Method seems to be inapplicable in practice in real, large systems, due to its requirement for maintaining constant metabolite concentrations. This is basically impossible to achieve: while for some cases (branch points) a small change in the metabolite concentration may not affect the overall outcome significantly, in general one can never be sure this will not be the case (and the number of interactions will greatly exceed the number of fluxes), and to date we do not know of any experimental attempt to exploit the method in practice.

Multi-site modulation

Whilst the Universal Method probably cannot be made to work as advertised, it does draw attention to the need - whatever the effects on the rest of metabolism - for multisite modulations to be performed if there is to be a substantial increase in flux, and this is now widely recognised (Cornish-Bowden, 1995; Cornish-Bowden *et al.*, 1995; Fell, 1998; Fell & Thomas, 1995; Niederberger *et al.*, 1992; Thomas & Fell, 1998). This does not contradict any of the insights of MCA, and in fact it can be shown both by analysis and simulation (Small & Kacser, 1993a; Small & Kacser, 1993b) that this result is to be expected: as soon as the step with higher control becomes faster (as happens with overexpression) the control shifts to other steps in the pathway. It is now evident that for any strategy to be successful in increasing the flux of a pathway substantially there is a requirement for the manipulation of *several* steps. This was clearly demonstrated experimentally by Niederberger and colleagues in their classical study (Niederberger *et al.*, 1992) and has been discussed at some length by Fell and Thomas (Fell, 1998; Fell & Thomas, 1995; Thomas & Fell, 1998). In a recent conference it was evident that the metabolic engineering community (both research and industry) is converging to this conclusion, and it is now largely accepted that to increase flux one should manipulate at least two metabolic steps (Mendes & Kell, 1997). Removing the fluxes to unproductive pathways is likely to be much more significant for mature fermentations than seeking solely to stimulate the flux through the desired one *per se* (Holms, 1996; Holms, Hamilton & Mousdale, 1991).

Active learning and a post-Baconian approach to science in the post-genomic era

The commonest conventional method of experimental science, generally referred to as ‘the scientific method’, involves the preparation of an experimental system in a specified state and the manipulation of, preferably, a single parameter, whereupon one observes the time-evolution of the values of one or more variables compared to that of a control in which the ‘triggering’ manipulation is not performed. The parameter may then be moved to different set points. Each of those variables might also be controlled at a fixed level, i.e. as a parameter, and comparable experiments performed. If the system is comparatively simple and well behaved (e.g. asymptotically stable, and not chaotic as in (Davey *et al.*, 1996)) and the problem well-posed it is usually possible to determine the form and parameterisation of the system equations by mathematical fitting procedures (Mendes & Kell, 1998a), leading to what is usually considered an understanding of the system. However, this is true only for simple systems, and one may put forward the views that (i) this kind of deductive reasoning is that usually practised in the MCA community (though there

is evidence that some of the theorems of MCA were originally induced from the results of simulations on an analogue computer), and (ii) the functional genomics agenda (Kell, 1998) is likely to be much better attacked under current conditions via an inductive type of approach.

Indeed, we consider that complex systems cannot be treated to best advantage (Westerhoff & Kell, 1996) in this more classical, deductive way. First of all, there are far too many variables and potential parameters for an exhaustive set of experiments to be performed, and those parameter sets producing ‘desirable’ outcomes may be few and far between. (For n parameters which might adopt m values the number of combinations is obviously m^n , and even if m is only a miserable 2 for $n=100$, $2^n \sim 10^{31}$, and the lifetime of the Universe is ‘only’ some 10^{17} s (Barrow & Silk, 1995).) The inevitable conclusion for the study of complex systems is that we must vary *many* (or at least several) parameters at a time (over a large range - see above) and use the methods of multivariate statistics and machine learning to deconvolute the data so obtained to extract those features most relevant to the operation of the system. Then, because of the high dimensionality of the system and problem, we must iterate this process further (somewhat in the way in which we traditionally need to provide *rounds* of mutation and selection in fermentation development programmes (Crueger & Crueger, 1989)). Indeed, our own approach in recent years to the understanding of complex cellular systems has been to exploit spectroscopic methods such as pyrolysis mass spectrometry (Broadhurst *et al.*, 1997; Gilbert *et al.*, 1997; Goodacre & Kell, 1996; Goodacre, Kell & Bianchi, 1993; Goodacre, Neal & Kell, 1994a; Goodacre, Neal & Kell, 1996a; Goodacre *et al.*, 1994b; Taylor *et al.*, 1998), FT-IR (Goodacre, Rooney & Kell, 1998a; Goodacre *et al.*, 1996b; Oliver *et al.*, 1998; Winson *et al.*, 1997; Winson *et al.*, 1998), and dispersive Raman (Goodacre *et al.*, 1998b) in which hundreds of variables are measured *simultaneously* and to couple these measurements with advanced chemometric and related analyses based on the methods of artificial intelligence and evolutionary computing (Bäck, Fogel & Michalewicz, 1997; Rich & Knight, 1991; Weiss & Kulikowski, 1991).

Although the above described the overall structure of a *single* experiment, scientific research of course proceeds by a *process* of experimental hypothesis testing (e.g. (Oldroyd, 1986)), and it is appropriate to end by outlining one way of computer-assisted knowledge acquisition with which we think important progress might be made. This process is an active approach, which in fact differs markedly from the passive nature of most ‘scientific discovery’ systems (Langley *et al.*, 1987), which either receive data all at once in a single batch, or have no choice over the next example (Raju & Cooney, 1998), and suffer from the problem that most of the observables have little bearing on the overall outcome (Blum & Langley, 1997) and for the purposes of this analysis amount to ‘noise’. The study of systems that can choose the next experiment is known as ‘active learning’. There are two computational tasks in active learning: formation of hypotheses that are consistent with known background knowledge and experimental results, and selection of the best experiment (or set of experiments) to discriminate between hypotheses. It should be noted that experiment selection in active learning is not to be confused with the traditional statistical study of experimental design, where the difference is between deciding which question to ask next (active learning) versus ensuring that a set of experiments can answer a question (traditional experimental design).

To conclude, it seems reasonable that active learning approaches can lead us efficiently to means for asking and answering the right kinds of question at the right kind of complexity in the post-genomic era.

Acknowledgments

We thank Ross King and Steve Oliver for useful discussions, and the BBSRC for financial support.

Table 1. Some explicit and implicit assumptions of modern MCA, and some inadequacies of its usual implementations in a biotechnological context.

ASSUMPTION/MISAPPLICATION	COMMENTS/CONSEQUENCES
All cells are the same	Heterogeneity is very much greater than normally assumed, and this can be determined experimentally using single-cell analyses
Simple models are adequate	Genome sequencing has uncovered the fact that we know the function of fewer than half of their genes, and there is evidence that almost all contribute to fitness even in laboratory conditions. Much more of metabolism is relevant to a flux than is normally recognised.
The Universal Method permits a rational approach to the optimisation of flux in any metabolic system	It doesn't work if (i) the end-product feeds back to inhibit its own synthesis, whether kinetically or by mass action, and/or (ii) there are interactions between pathway branches involving moiety-conserved cycles.
The coefficients of MCA determined using large changes are not too badly different from those obtained via very small parameter changes	Nonlinearities, unknown interactions and the overall complexity of biological systems mean that deviation indices are reasonably small only in simple systems.

References

- Agius, L. & Peak, M. (1997). Binding and translocation of glucokinase in hepatocytes. *Biochemical Society Transactions* **25**, 145-150.
- Agius, L., Peak, M., Newgard, C. B., GomezFoix, A. M. & Guinovart, J. J. (1996). Evidence for a role of glucose-induced translocation of glucokinase in the control of hepatic glycogen synthesis. *Journal of Biological Chemistry* **271**, 30479-30486.
- Agius, L. & Sherratt, H. S. A. (1997). Channelling in intermediary metabolism. Portland Press, London.
- Ainscow, E. K. & Brand, M. D. (1995). Top-down control analysis of systems with more than one common intermediate. *Eur. J. Biochem.* **231**, 579-586.
- Ainscow, E. K. & Brand, M. D. (1998a). Control analysis of systems with reaction blocks that 'cross-talk'. *Biochim. Biophys. Acta* **1366**, 284-290.
- Ainscow, E. K. & Brand, M. D. (1998b). Errors associated with metabolic control analysis. Application of Monte Carlosimulation of experimental data. *J. Theoret. Biol.* **194**, 223-233.
- Amann, R. I., Ludwig, W. & Schleifer, K. H. (1995). Phylogenetic identification and *in situ* detection of individual microbial cells without cultivation. *Microbiological Reviews* **59**, 143-169.
- Anderson, N. L. & Anderson, N. G. (1998). Proteome and proteomics: New technologies, new concepts, and new words. *Electrophoresis* **19**, 1853-1861.
- Bäck, T., Fogel, D. B. & Michalewicz, Z. (1997). Handbook of evolutionary computation. IOPPublishing/Oxford University Press, Oxford.
- Barrow, J. D. & Silk, J. (1995). *The left hand of creation: the origin and evolution of the expanding universe*. Penguin, London.
- Blackstock, W. P. & Weir, M. P. (1999). Proteomics: quantitative and physical mapping of cellular proteins. *Tibtech* **17**, 121-127.
- Blattner, F. R., Plunkett, G., Bloch, C. A., Perna, N. T., Burland, V., Riley, M., ColladoVides, J., Glasner, J. D., Rode, C. K., Mayhew, G. F., Gregor, J., Davis, N. W., Kirkpatrick, H. A., Goeden, M. A., Rose, D. J., Mau, B. & Shao, Y. (1997). The complete genome sequence of *Escherichia coli* K-12. *Science* **277**, 1453-1462.
- Blum, A. L. & Langley, P. (1997). Selection of relevant features and examples in machine learning. *Artificial Intelligence* **97**, 245-271.
- Bork, P., Dandekar, T., Diaz-Lazcoz, Y., Eisenhaber, F., Huynen, M. & Yuan, Y. P. (1998). Predicting function: From genes to genomes and back. *Journal of Molecular Biology* **283**, 707-725.
- Boucherie, H., Sagliocco, F., Joubert, R., Maillet, I., Labarre, J. & Perrot, M. (1996). Two-dimensional gel protein database of *Saccharomyces cerevisiae*. *Electrophoresis* **17**, 1683-1699.
- Bowtell, D. D. L. (1999). Options available - from start to finish - for obtaining expression data by microarray. *Nature Genetics* **21**, 25-32.
- Brand, M. D. (1996). Top down metabolic control analysis. *Journal of Theoretical Biology* **182**, 351-360.
- Brand, M. D. (1998). Top-down elasticity analysis and its application to energy metabolism in isolated mitochondria and intact cells. *Molecular and Cellular Biochemistry* **184**, 13-20.
- Broadhurst, D., Goodacre, R., Jones, A., Rowland, J. J. & Kell, D. B. (1997). Genetic algorithms as a method for variable selection in multiple linear regression and partial least squares regression, with applications to pyrolysis mass spectrometry. *Analytica Chimica Acta* **348**, 71-86.
- Brown, G. C., Hafner, R. P. & Brand, M. D. (1990). A top-down approach to the determination of control coefficients in metabolic control theory. In *European Journal of Biochemistry*, vol. 188, pp. 321-325.
- Brown, P. O. & Botstein, D. (1999). Exploring the new world of the genome with DNA microarrays. *Nature Genetics* **21**, 33-37.
- Cash, P. (1998). Characterisation of bacterial proteomes by two-dimensional electrophoresis. *Analytica Chimica Acta* **372**, 121-145.
- Chu, S., DeRisi, J., Eisen, M., Mulholland, J., Botstein, D., Brown, P. O. & Herskowitz, I. (1998). The transcriptional program of sporulation in budding yeast. *Science* **282**, 699-705.
- Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V., Eglmeier, K., Gas, S., Barry, C. E., Tekaiia, F., Badcock, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., Hamlin, N., Holroyd, S., Hornby, T., Jagels, K., Krogh, A., McLean, J., Moule, S., Murphy, L., Oliver, K., Osborne, J., Quail, M. A., Rajandream, M. A., Rogers, J., Rutter, S., Seeger, K., Skelton, J., Squares, R., Squares, S., Sulston, J. E., Taylor, K., Whitehead, S. &

- Barrell, B. G. (1998). Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence. *Nature* **393**, 537-544.
- Cornish-Bowden, A. (1995). Kinetics of multi-enzyme systems. In *Biotechnology*, vol. 9 (ed. H. J. Rehm, G. Reed, A. Pühler and P. Stadler), pp. 121-136. Verlag Chemie, Weinheim.
- Cornish-Bowden, A. & Cárdenas, M. L. (1990). Control of metabolic processes. Plenum Press, New York.
- Cornish-Bowden, A., Hofmeyr, J.-H. S. & Cárdenas, M. L. (1995). Strategies for manipulating metabolic fluxes in biotechnology. *Bioorg. Chem.* **23**, 439-449.
- Crueger, W. & Crueger, A. (1989). *Biotechnology: a textbook of industrial microbiology*. Sinauer Associates, Sunderland, MA.
- Davey, H. M., Davey, C. L., Woodward, A. M., Edmonds, A. N., Lee, A. W. & Kell, D. B. (1996). Oscillatory, stochastic and chaotic growth rate fluctuations in permissively-controlled yeast cultures. *Biosystems* **39**, 43-61.
- Davey, H. M. & Kell, D. B. (1996). Flow cytometry and cell sorting of heterogeneous microbial populations: the importance of single-cell analysis. *Microbiol. Rev.* **60**, 641-696.
- de Saizieu, A., Certa, U., Warrington, J., Gray, C., Keck, W. & Mous, J. (1998). Bacterial transcript imaging by hybridization of total RNA to oligonucleotide arrays. *Nature Biotechnol.* **16**, 45-48.
- Debouck, C. & Goodfellow, P. N. (1999). DNA microarrays in drug discovery and development. *Nature Genetics* **21**, 48-50.
- DeRisi, J. L., Iyer, V. R. & Brown, P. O. (1997). Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* **278**, 680-686.
- Duggan, D. J., Bittner, M., Chen, Y. D., Meltzer, P. & Trent, J. M. (1999). Expression profiling using cDNA microarrays. *Nature Genetics* **21**, 10-14.
- Dujon, B. (1998). European Functional Analysis Network (EUROFAN) and the functional analysis of the *Saccharomyces cerevisiae* genome. *Electrophoresis* **19**, 617-624.
- Durrieu, G., Letellier, T., Antoch, J., Deshouillers, J. M., Malgat, M. & Mazat, J. P. (1997). Identification of mitochondrial deficiency using principal component analysis. *Molecular and Cellular Biochemistry* **174**, 149-156.
- Ehlde, M. & Zacchi, G. (1996). Influence of experimental errors on the determination of flux control coefficients from transient metabolic concentrations. *Biochemical Journal* **313**, 721-727.
- Eisen, M. B., Spellman, P. T., Brown, P. O. & Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. *Proc. Natl. Acad. Sci.* **95**, 14863-14868.
- Fell, D. A. (1992). Metabolic Control Analysis - a Survey of its Theoretical and Experimental Development. *Biochemical Journal* **286**, 313-330.
- Fell, D. A. (1996). *Understanding the Control of Metabolism*. Portland Press, London.
- Fell, D. A. (1998). Increasing the flux in metabolic pathways: A metabolic control analysis perspective. *Biotechnology and Bioengineering* **58**, 121-124.
- Fell, D. A. & Sauro, H. M. (1985). Metabolic control and its analysis. Additional relationships between elasticities and control coefficients. *European Journal of Biochemistry* **148**, 555-561.
- Fell, D. A. & Thomas, S. (1995). Physiological control of metabolic flux: the requirement for multisite modulation. *Biochemical Journal* **311**, 35-39.
- Forterre, P. (1997a). Archaea: what can we learn from their sequences? *Current Opinion in Genetics & Development* **7**, 764-770.
- Forterre, P. (1997b). Protein versus rRNA: Problems in rooting the universal tree of life. *ASM News* **63**, 89-95.
- Fuqua, C., S.C., W. & Greenberg, E. P. (1996). Census and consensus in bacterial ecosystems: the *LuxR-LuxI* family of quorum-sensing transcriptional regulators. *Annual Review of Microbiology* **50**, 727-751.
- Garrels, J. I. (1996). YPD - a Database For the Proteins of *Saccharomyces cerevisiae*. *Nucleic Acids Research* **24**, 46-49.
- Gilbert, R. J., Goodacre, R., Woodward, A. M. & Kell, D. B. (1997). Genetic programming: A novel method for the quantitative analysis of pyrolysis mass spectral data. *Analytical Chemistry* **69**, 4381-4389.
- Goffeau, A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., Galibert, F., Hoheisel, J. D., Jacq, C., Johnston, M., Louis, E. J., Mewes, H. W., Murakami, Y., Philippsen, P., Tettelin, H. & Oliver, S. G. (1996). Life With 6000 Genes. *Science* **274**, 546-567.
- Goodacre, R. & Kell, D. B. (1996). Pyrolysis mass spectrometry and its applications in biotechnology. *Curr. Opinion Biotechnol.* **7**, 20-28.

- Goodacre, R., Kell, D. B. & Bianchi, G. (1993). Rapid assessment of the adulteration of virgin olive oils by other seed oils using pyrolysis mass spectrometry and artificial neural networks. *Journal of the Science of Food and Agriculture* **63**, 297-307.
- Goodacre, R., Neal, M. J. & Kell, D. B. (1994a). Rapid and quantitative analysis of the pyrolysis mass spectra of complex binary and tertiary mixtures using multivariate calibration and artificial neural networks. *Analytical Chemistry* **66**, 1070-1085.
- Goodacre, R., Neal, M. J. & Kell, D. B. (1996a). Quantitative analysis of multivariate data using artificial neural networks: a tutorial review and applications to the deconvolution of pyrolysis mass spectra. *Zentralblatt. für Bakteriologie* **284**, 516-539.
- Goodacre, R., Rooney, P. J. & Kell, D. B. (1998a). Rapid analysis of microbial systems using vibrational spectroscopy and supervised learning methods: application to the discrimination between methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*. In *Proc. SPIE (Infrared Spectroscopy: New Tool in Medicine)*, vol. 3257 (ed. M. Jackson and H. H. Mantsch), pp. 220-229. SPIE, San Jose, California, USA.
- Goodacre, R., Timmins, É. M., Burton, R., Kaderbhai, N., Woodward, A., Kell, D. B. & Rooney, P. J. (1998b). Rapid identification of urinary tract infection bacteria using hyperspectral, whole organism fingerprinting and artificial neural networks. *Microbiology* **144**, 1157-1170.
- Goodacre, R., Timmins, É. M., Rooney, P. J., Rowland, J. J. & Kell, D. B. (1996b). Rapid identification of *Streptococcus* species using diffuse reflectance-absorbance Fourier transform infrared spectroscopy and artificial neural networks. *FEMS Microbiol. Lett.* **140**, 233-239.
- Goodacre, R., Trew, S., Wrigley-Jones, C., Neal, M. J., Maddock, J., Ottley, T. W., Porter, N. & Kell, D. B. (1994b). Rapid screening for metabolite overproduction in fermentor broths, using pyrolysis mass spectrometry with multivariate calibration and artificial neural networks. *Biotechnol. Bioeng.* **44**, 1205-1216.
- Heinrich, R. & Rapoport, T. A. (1973). Linear theory of enzymatic chains: its application for the analysis of the crossover theorem and of the glycolysis of human erythrocytes. *Acta Biologica et Medica Germanica* **31**, 479-94.
- Heinrich, R. & Rapoport, T. A. (1974). A linear steady-state treatment of enzymatic chains. General properties, control and effector strength. *European Journal of Biochemistry* **42**, 89-95.
- Heinrich, R. & Schuster, S. (1996). *The Regulation of Cellular Systems*. Chapman & Hall, New York.
- Herweijer, M. A., Berden, J. A. & Slater, E. C. (1986). Uncoupler-Inhibitor Titrations of ATP-Driven Reverse Electron- Transfer in Bovine Submitochondrial Particles Provide Evidence For Direct Interaction Between ATPase and NADH-Q Oxidoreductase. *Biochimica Et Biophysica Acta* **849**, 276-287.
- Hinton, J. C. D. (1997). The *Escherichia coli* genome sequence: the end of an era or the start of the FUN? *Mol. Microbiol.* **26**, 417-422.
- Hitchens, G. D. & Kell, D. B. (1983a). On the functional unit of energy coupling in photophosphorylation by bacterial chromatophores. *Biochim. Biophys. Acta* **723**, 308-316.
- Hitchens, G. D. & Kell, D. B. (1983b). Uncouplers can shuttle rapidly between localised energy coupling sites during photophosphorylation by chromatophores of *Rhodospseudomonas capsulata* N22. *Biochem. J.* **212**, 25-30.
- Ho, M. W. (1998). *The Rainbow and the Worm: the Physics of Organisms*, Second edition. World Scientific, Singapore.
- Hofmeyr, J.-H. S., Kacser, H. & van der Merwe, K. J. (1986). Metabolic control analysis of moiety-conserved cycles. *Eur. J. Biochem.* **155**, 631-641.
- Holms, H. (1996). Flux analysis and control of the central metabolic pathways in *Escherichia coli*. *Fems Microbiology Reviews* **19**, 85-116.
- Holms, W. H., Hamilton, I. D. & Mousdale, D. (1991). Improvements to Microbial Productivity By Analysis of Metabolic Fluxes. *Journal of Chemical Technology and Biotechnology* **50**, 139-141.
- Humphery-Smith, I., Cordwell, S. J. & Blackstock, W. P. (1997). Proteome research: complementarity and limitations with respect to the RNA and DNA worlds. *Electrophoresis* **18**, 1217-1242.
- Iyer, V. R., Eisen, M. B., Ross, D. T., Schuler, G., Moore, T., Lee, J. C. F., Trent, J. M., Staudt, L. M., Hudson, J., Boguski, M. S., Lashkari, D., Shalon, D., Botstein, D. & Brown, P. O. (1999). The transcriptional program in the response of human fibroblasts to serum. *Science* **283**, 83-87.
- Kacser, H. (1983). The control of enzyme systems in vivo: elasticity analysis of the steady state. *Biochem. Soc. Trans.* **11**, 35-40.

- Kacser, H. & Akerenza, L. (1993). A universal method for achieving increases in metabolite production. *Eur. J. Biochem.* **216**, 361-367.
- Kacser, H. & Burns, J. A. (1973). The control of flux. In *Rate Control of Biological Processes. Symposium of the Society for Experimental Biology Vol 27* (ed. D. D. Davies), pp. 65-104. Cambridge University Press, Cambridge.
- Kacser, H. & Burns, J. A. (1981). The molecular basis of dominance. *Genetics* **97**, 639-636.
- Kaprelyants, A. S. & Kell, D. B. (1992). Rapid assessment of bacterial viability and vitality using rhodamine 123 and flow cytometry. *Journal of Applied Bacteriology* **72**, 410-422.
- Kaprelyants, A. S. & Kell, D. B. (1993). Dormancy in stationary-phase cultures of *Micrococcus luteus*: flow cytometric analysis of starvation and resuscitation. *Applied and Environmental Microbiology* **59**, 3187-3196.
- Kaprelyants, A. S., Mukamolova, G. V. & Kell, D. B. (1994). Estimation of dormant *Micrococcus luteus* cells by penicillin lysis and by resuscitation in cell-free spent medium at high dilution. *FEMS Microbiology Letters* **115**, 347-352.
- Kaprelyants, A. S., Mukamolova, G. V., Kormer, S. S., Weichart, D. H., Young, M. & Kell, D. B. (1999). Intercellular signalling and the multiplication of prokaryotes: bacterial cytokines. *Symp. Soc. Gen. Microbiol.* **57**, 33-69.
- Kell, D. B. (1979). On the functional proton current pathway of electron transport phosphorylation: an electrodic view. *Biochim./Biophys. Acta* **549**, 55-99.
- Kell, D. B. (1987). Forces, fluxes and the control of microbial growth and metabolism. The twelfth Fleming lecture. *J. Gen. Microbiol.* **133**, 1651-1665.
- Kell, D. B. (1988). Protonmotive energy-transducing systems: some physical principles and experimental approaches. In *Bacterial Energy Transduction* (ed. C. J. Anthony), pp. 429-490. Academic Press, London.
- Kell, D. B. (1992). The protonmotive force as an intermediate in electron transport-linked phosphorylation: problems and prospects. *Curr. Top. Cell. Reg.* **33**, 279-289.
- Kell, D. B. (1998). From code to mode for orphan genes. *Trends Biotechnol.* **16**, 491-493.
- Kell, D. B. & Hitchens, G. D. (1983). Coherent properties of the membranous systems of electron transport phosphorylation. In *Coherent Excitations in Biological Systems* (ed. H. Fröhlich and F. Kremer), pp. 178-198. Springer-Verlag, Berlin.
- Kell, D. B., Kaprelyants, A. S. & Grafen, A. (1995). On pheromones, social behaviour and the functions of secondary metabolism in bacteria. *Trends Ecol. Evolution* **10**, 126-129.
- Kell, D. B., Kaprelyants, A. S., Weichart, D. H., Harwood, C. L. & Barer, M. R. (1998). Viability and activity in readily culturable bacteria: a review and discussion of the practical issues. *Antonie van Leeuwenhoek* **73**, 169-187.
- Kell, D. B., Ryder, H. M., Kaprelyants, A. S. & Westerhoff, H. V. (1991). Quantifying heterogeneity: Flow cytometry of bacterial cultures. *Antonie van Leeuwenhoek* **60**, 145-158.
- Kell, D. B. & Sonnleitner, B. (1995). GMP - Good Modelling Practice: an essential component of good manufacturing practice. *Trends Biotechnol.* **13**, 481-492.
- Kell, D. B., van Dam, K. & Westerhoff, H. V. (1989). Control analysis of microbial growth and productivity. *Symp. Soc. Gen. Microbiol.* **44**, 61-93.
- Kell, D. B. & Welch, G. R. (1991). No turning back, Reductonism and Biological Complexity. *Times Higher Educational Supplement* **9th August**, 15.
- Kell, D. B. & Westerhoff, H. V. (1986a). Metabolic control theory: its role in microbiology and biotechnology. *FEMS Microbiol. Rev.* **39**, 305-320.
- Kell, D. B. & Westerhoff, H. V. (1986b). Towards a rational approach to the optimization of flux in microbial biotransformations. *Trends Biotechnol.* **4**, 137-142.
- Kell, D. B. & Westerhoff, H. V. (1990). Control analysis of organised multienzyme systems. In *Structural and organizational aspects of metabolic regulation* (ed. P. Srere, M. E. Jones and C. Mathews), pp. 273-289. Alan R. Liss, New York.
- Kholodenko, B. N., Cascante, M., Hoek, J. B., Westerhoff, H. V. & Schwaber, J. (1998). Metabolic design: How to engineer a living cell to desired metabolite concentrations and fluxes. *Biotechnology and Bioengineering* **59**, 239-247.
- Koonin, E. V. & Galperin, M. Y. (1997). Prokaryotic genomes: the emerging paradigm of genome-based microbiology. *Current Opinion in Genetics & Development* **7**, 757-763.

- Koonin, E. V., Mushegian, A. R., Galperin, M. Y. & Walker, D. R. (1997). Comparison of archaeal and bacterial genomes: computer analysis of protein sequences predicts novel functions and suggests a chimeric origin for the archaea. *Molecular Microbiology* **25**, 619-637.
- Langley, P., Simon, H. A., Bradshaw, G. L. & Zytlow, J. M. (1987). *Scientific Discovery: computational exploration of the creative processes*. MIT Press, Cambridge, MA.
- Letellier, T., Malgat, M., Rossignol, R. & Mazat, J. P. (1998). Metabolic control analysis and mitochondrial pathologies. *Molecular and Cellular Biochemistry* **184**, 409-417.
- Lipshutz, R. J., Fodor, S. P. A., Gingeras, T. R. & Lockhart, D. J. (1999). High density synthetic oligonucleotide arrays. *Nature Genetics* **21**, 20-24.
- Lopez de Felipe, F., Kleerebezem, M., de Vos, W. M. & Hugenholtz, J. (1998). Cofactor engineering: a novel approach to metabolic engineering in *Lactococcus lactis* by controlled expression of NADH oxidase. *J. Bacteriol.* **180**, 3804-3808.
- Marton, M. J., DeRisi, J. L., Bennett, H. A., Iyer, V. R., Meyer, M. R., Roberts, C. J., Stoughton, R., Burchard, J., Slade, D., Dai, H. Y., Bassett, D. E., Hartwell, L. H., Brown, P. O. & Friend, S. H. (1998). Drug target validation and identification of secondary drug target effects using DNA microarrays. *Nature Medicine* **4**, 1293-1301.
- Mazat, J. P., Letellier, T., Bedes, F., Malgat, M., Korzeniewski, B., Jouaville, L. S. & Morkuniene, R. (1997). Metabolic control analysis and threshold effect in oxidative phosphorylation: Implications for mitochondrial pathologies. *Molecular and Cellular Biochemistry* **174**, 143-148.
- Mazat, J. P., Letellier, T., Malgat, M., Rossignol, R., Korzeniewski, B., Demaugre, F. & Leroux, J. P. (1998). Inborn errors of metabolism in the light of metabolic control analysis. *Biochemical Society Transactions* **26**, 141-145.
- McVeigh, H. P., Munro, J. & Embley, T. M. (1996). Molecular evidence for the presence of novel actinomycete lineages in a temperate forest soil. *Journal of Industrial Microbiology* **17**, 197-204.
- Mendes, P. (1993). GEPASI: a software package for modelling the dynamics, steady states and control of biochemical and other systems. *Computer Appl. Biosci.* **9**, 563-571.
- Mendes, P. (1997). Biochemistry by numbers: simulation of biochemical pathways with Gepasi 3. *Trends Biochem. Sci.* **22**, 361-363.
- Mendes, P. & Kell, D. B. (1996). On the analysis of the inverse problem of metabolic pathways using artificial neural networks. *Biosystems* **38**, 15-28.
- Mendes, P. & Kell, D. B. (1997). Making cells work: metabolic engineering for everyone. *Trends Biotechnol.* **15**, 6-7.
- Mendes, P. & Kell, D. B. (1998a). Non-linear optimization of biochemical pathways: applications to metabolic engineering and parameter estimation. *Bioinformatics* **14**, 869-883.
- Mendes, P. & Kell, D. B. (1998b). Numerical optimisation and simulation for rational metabolic engineering. In *Biothermokinetics in the post genomic era*, vol. 4. *Biothermokinetics* (ed. C. Larsson, I.-L. Pålman and L. Gustaffson), pp. 345-349. Chalmers Repro Service, Göteborg.
- Mendes, P., Kell, D. B. & Welch, G. R. (1995). Metabolic channeling in organized enzyme systems: experiments and models. In *Enzymology in vivo* (ed. K. M. Brindle), pp. 1-19. JAI Press, London.
- Mendes, P., Kell, D. B. & Westerhoff, H. V. (1992). Channelling can decrease pool size. *European Journal of Biochemistry* **204**, 257-266.
- Mendes, P., Kell, D. B. & Westerhoff, H. V. (1996). Why and when channeling can decrease pool size at constant net flux in a simple dynamic channel. *Biochimica et Biophysica Acta-General Subjects* **1289**, 175-186.
- Mitchell, P. (1966). Chemiosmotic coupling in oxidative and photosynthetic phosphorylation. *Biol. Rev.* **41**, 445-502.
- Mukamolova, G. V., Kaprelyants, A. S., Young, D. I., Young, M. & Kell, D. B. (1998). A bacterial cytokine. *Proc. Natl. Acad. Sci.* **95**, 8916-8921.
- Nicholls, D. G. & Ferguson, S. J. (1992). *Bioenergetics 2*. Academic Press, London.
- Niederberger, P., Prasad, R., Miozzari, G. & Kacser, H. (1992). A Strategy for increasing an in vivo flux by genetic manipulations - the tryptophan system of yeast. *Biochemical Journal* **287**, 473-479.
- Oldroyd, D. (1986). *The arch of knowledge: an introduction to the history of the philosophy and methodology of science*. Methuen, New York.
- Oliver, S. G. (1996). From DNA sequence to biological function. *Nature* **379**, 597-600.

- Oliver, S. G. & Baganz, F. (1998). The yeast genome: systematic analysis of DNA sequence and biological function. In *Genomics: commercial opportunities from a scientific revolution* (ed. L. G. Copping, G. K. Dixon and D. J. Livingstone), pp. 37-51. Bios, Oxford.
- Oliver, S. G., Winson, M. K., Kell, D. B. & Baganz, F. (1998). Systematic functional analysis of the yeast genome. *Trends Biotechnol.* **16**, 373-378.
- Ovádi, J. (1995). *Cell architecture and metabolic channeling*. Springer-Verlag, New York.
- Raju, G. K. & Cooney, C. L. (1998). Active learning from process data. *AIChE Journal* **44**, 2199-2211.
- Reder, C. (1988). Metabolic control theory: a structural approach. *J. Theoret. Biol.* **135**, 175-201.
- Reich, J. G. & Sel'kov, E. E. (1981). *Energy metabolism of the cell a theoretical treatise*. Academic Press, London.
- Rich, E. & Knight, K. (1991). *Artificial Intelligence*. McGraw Hill, New York.
- Rivera, M. C., Jain, R., Moore, J. E. & Lake, J. A. (1998). Genomic evidence for two functionally distinct gene classes. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 6239-6244.
- Schena, M., Shalon, D., Heller, R., Chai, A., Brown, P. O. & Davis, R. W. (1996). Parallel human genome analysis - microarray-based expression monitoring of 1000 genes. *Proc. Natl. Acad. Sci.* **93**, 10614-10619.
- Schlosser, P. M., Holcomb, T. & Bailey, J. E. (1993). Determining metabolic sensitivity coefficients directly from experimental data. *Biotechnology and Bioengineering* **41**, 1027-1038.
- Schuster, R. & Holzhutter, H. G. (1995). Use of Mathematical Models For Predicting the Metabolic Effect of Large-Scale Enzyme-Activity Alterations: Application to Enzyme Deficiencies of Red Blood Cells. *European Journal of Biochemistry* **229**, 403-418.
- Shapiro, H. M. (1995). *Practical Flow Cytometry, 3rd edition*, 3rd edition. John Wiley, New York.
- Shaw, A. D., diCamillo, A., Vlahov, G., Jones, A., Bianchi, G., Rowland, J. & Kell, D. B. (1997). Discrimination of the variety and region of origin of extra virgin olive oils using C-13 NMR and multivariate calibration with variable reduction. *Analytica Chimica Acta* **348**, 357-374.
- Simpson, T. W., Colon, G. E. & Stephanopoulos, G. (1995). 2 Paradigms of metabolic engineering applied to amino acid biosynthesis. *Biochemical Society Transactions* **23**, 381-387.
- Simpson, T. W., Shimizu, H. & Stephanopoulos, G. (1998). Experimental determination of group flux control coefficients in metabolic networks. *Biotechnology and Bioengineering* **58**, 149-153.
- Small, J. R. (1993). Flux control coefficients determined by inhibitor titration: the design and analysis of experiments to minimize errors. *Biochemical Journal* **296**, 423-433.
- Small, J. R. & Kacser, H. (1993a). Responses of Metabolic Systems to Large Changes in Enzyme-Activities and Effectors .1. The Linear Treatment of Unbranched Chains. *European Journal of Biochemistry* **213**, 613-624.
- Small, J. R. & Kacser, H. (1993b). Responses of Metabolic Systems to Large Changes in Enzyme-Activities and Effectors .2. The Linear Treatment of Branched Pathways and Metabolite Concentrations - Assessment of the General Nonlinear Case. *European Journal of Biochemistry* **213**, 625-640.
- Smith, V., Botstein, D. & Brown, P. O. (1995). Genetic Footprinting - a Genomic Strategy For Determining a Gene's Function Given Its Sequence. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 6479-6483.
- Smith, V., Chou, K. N., Lashkari, D., Botstein, D. & Brown, P. O. (1996). Functional analysis of the genes of yeast chromosome V by genetic footprinting. *Science* **274**, 2069-2074.
- Spellman, P. T., Sherlock, G., Zhang, M. Q., Iyer, V. R., Anders, K., Eisen, M. B., Brown, P. O., Botstein, D. & Futcher, B. (1998). Comprehensive identification of cell cycle-regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization. *Molecular Biology of the Cell* **9**, 3273-3297.
- Stephanopoulos, G. & Simpson, T. W. (1997). Flux amplification in complex metabolic networks. *Chemical Engineering Science* **52**, 2607-2627.
- Tamayo, P., Slonim, D., Mesirov, J., Zhu, Q., Kitareewan, S., Dmitrovsky, E., Lander, E. S. & Golub, T. R. (1999). Interpreting patterns of gene expression with self-organizing maps: Methods and application to hematopoietic differentiation. *Proc. Natl. Acad. Sci.* **96**, 2907-2912.
- Taylor, J., Goodacre, R., Wade, W. G., Rowland, J. J. & Kell, D. B. (1998). The deconvolution of pyrolysis mass spectra using genetic programming: application to the identification of some *Eubacterium* species. *FEMS Microbiology Letters* **160**, 237-246.

- Teusink, B., Baganz, F., Westerhoff, H. V. & Oliver, S. G. (1998). Metabolic Control Analysis as a tool in the elucidation of the function of novel genes. In *Methods in Microbiology: Yeast gene analysis* (ed. M. F. Tuite and A. J. P. Brown), pp. 297-336. Academic Press, London.
- Thatcher, J. W., Shaw, J. M. & Dickinson, W. J. (1998). Marginal fitness contributions of nonessential genes in yeast. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 253-257.
- Thomas, S. & Fell, D. A. (1995). Error and bias in control coefficients calculated from elasticities. *Biochem. Soc. Trans.* **23**, S294.
- Thomas, S. & Fell, D. A. (1998). The role of multiple enzyme activation in metabolic flux control. *Advances in Enzyme Regulation* **38**, 65-85.
- Velculescu, V. E., Zhang, L., Zhou, W., Vogelstein, J., Basrai, M. A., Bassett, D. E., Hieter, P., Vogelstein, B. & Kinzler, K. W. (1997). Characterization of the yeast transcriptome. *Cell* **88**, 243-251.
- Wanders, R. J. A., Groen, A. K., van Roermund, C. W. T. & Tager, J. M. (1984). Factors determining the relative contribution of the adenine nucleotide translocator and the ADP-regenerating system to the control of oxidative phosphorylation in isolated rat-liver mitochondria. *European Journal of Biochemistry* **142**, 417-24.
- Wang, J. H. & Hewick, R. M. (1999). Proteomics in drug discovery. *Drug Discovery Today* **4**, 129-133.
- Weiss, S. H. & Kulikowski, C. A. (1991). *Computer systems that learn: classification and prediction methods from statistics, neural networks, machine learning, and expert systems*. Morgan Kaufmann Publishers, San Mateo, CA.
- Welch, G. R. & Kell, D. B. (1986). Not just catalysts; the bioenergetics of molecular machines. In *The Fluctuating Enzyme* (ed. G. Welch, R.), pp. 451-492. John Wiley, New York.
- Westerhoff, H. V., Hellingwerf, K. J. & van Dam, K. (1983). Thermodynamic efficiency of microbial growth is low but optimal for maximal growth rate. *Proc. Natl. Acad. Sci. USA* **80**, 305-309.
- Westerhoff, H. V. & Kell, D. B. (1987). Matrix method for determining the steps most rate-limiting to metabolic fluxes in biotechnological processes. *Biotechnology and Bioengineering* **30**, 101-107.
- Westerhoff, H. V. & Kell, D. B. (1988). A control theoretical analysis of inhibitor titrations of metabolic channelling. *Comments Mol. Cell. Biophys.* **5**, 57-107.
- Westerhoff, H. V. & Kell, D. B. (1996). What BioTechnologists knew all along...? *J. Theoret. Biol.* **182**, 411-420.
- Westerhoff, H. V. & van Dam, K. (1987). *Thermodynamics and control of biological free energy transduction*. Elsevier, Amsterdam.
- Wilkins, M. R., Sanchez, J. C., Gooley, A. A., Appel, R. D., Humphery-Smith, I., Hochstrasser, D. F. & Williams, K. L. (1996). Progress with proteome projects - why all proteins expressed by a genome should be identified and how to do it. *Biotechnology & Genetic Engineering Reviews* **13**, 19-50.
- Wilkins, M. R., Williams, K. L., Appel, R. D. & Hochstrasser, D. F. (1997). Proteome research: new frontiers in functional genomics. Springer, Berlin.
- Williams, R. J. (1956). *Biochemical Individuality*. John Wiley, New York.
- Winson, M. K., Goodacre, R., Timmins, É. M., Jones, A., Alsberg, B. K., Woodward, A. M., Rowland, J. J. & Kell, D. B. (1997). Diffuse reflectance absorbance spectroscopy taking in chemometrics (DRASTIC). A hyperspectral FT-IR-based approach to rapid screening for metabolite overproduction. *Analytica Chimica Acta* **348**, 273-282.
- Winson, M. K., Todd, M., Rudd, B. A. M., Jones, A., Alsberg, B. K., Woodward, A. M., Goodacre, R., Rowland, J. J. & Kell, D. B. (1998). A DRASTIC (Diffuse Reflectance Absorbance Spectroscopy Taking in Chemometrics) Approach for the Rapid Analysis of Microbial Fermentation Products: Quantification of Aristeromycin and Neplanocin A in *Streptomyces citricolor* Broths. In *New frontiers in screening for microbial biocatalysts* (ed. K. Kieslich, C. P. van der Beek, J. A. M. de Bont and W. J. J. van den Tweel), pp. 185-191. Elsevier, Amsterdam.
- Wodicka, L., Dong, H. L., Mittmann, M., Ho, M. H. & Lockhart, D. J. (1997). Genome-wide expression monitoring in *Saccharomyces cerevisiae*. *Nature Biotechnology* **15**, 1359-1367.