



US008383126B2

(12) **United States Patent**
Mukamolova et al.

(10) **Patent No.:** **US 8,383,126 B2**
(45) **Date of Patent:** **Feb. 26, 2013**

(54) **BACTERIAL PHEROMONES AND USES THEREOF**

(75) Inventors: **Galina V. Mukamolova**, Moscow Region (RU); **Arseny S. Kaprelyants**, Moscow (RU); **Danielle I. Young**, Ceredigion (GB); **Douglas B. Kell**, Ceredigion (GB); **Michael Young**, Ceredigion (GB)

(73) Assignee: **University of Wales, Aberystwyth**, Aberystwyth, Ceredigion (GB)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 209 days.

(21) Appl. No.: **11/978,257**

(22) Filed: **Oct. 29, 2007**

(65) **Prior Publication Data**

US 2008/0206269 A1 Aug. 28, 2008

Related U.S. Application Data

(62) Division of application No. 09/445,289, filed as application No. PCT/GB98/01619 on Jun. 3, 1998.

(30) **Foreign Application Priority Data**

Jun. 4, 1997 (GB) 9711389.8
May 27, 1998 (GB) 9811221.2

(51) **Int. Cl.**

A61K 39/02 (2006.01)
A61K 39/04 (2006.01)
A61K 38/00 (2006.01)
C07K 1/00 (2006.01)

(52) **U.S. Cl.** **424/190.1**; 424/234.1; 424/248.1; 530/350; 530/300; 530/825; 514/1.1

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,331,318 B1 12/2001 Milstein
6,649,169 B2 11/2003 Dowling et al.
7,452,869 B2 11/2008 Hellner et al.
7,595,383 B1 * 9/2009 Gennaro et al. 536/23.7
2008/0206269 A1 8/2008 Mukamolova et al.

OTHER PUBLICATIONS

Skolnick et al. Trends in Biotechnology 18: 34-39, 2000.*
Illustrated Stedman's Medical Dictionary. 24th Edition, Williams and Wilkins, London. p. 707 (1982).
Mukamalova et al. Arch. Microbiol. 172: 9-14 (Jul. 1999).
Mukamalova et al. Antonie van Leeuwenhoek 67: 289-295 (1995).
U.S. Appl. No. 09/445,289, Mukamalova et al.
Badcock, K. et al., "Hypothetical 17.7 KD protein," TRPRO Database entry 033049, Jan. 1, 1998, Accession No. 033049.
Cole, S.T. et al. "Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence," Nature 393(6685): 537-44 (1998).

Cole, S.T. et al., "Hypothetical 15.7 KD protein," TRPRO Database entry P71755, Feb. 1, 1997, Accession No. P71755.

Cole, S.T. et al., "Hypothetical 18.8 KD protein," TRPRO Database entry 007747, Jul. 1, 1997, Accession No. 007747.

Cole, S.T. et al., "Hypothetical 38.1 KD protein," TRPRO Database entry 005594, Jul. 1, 1997, Accession No. 005594.

Eiglmeir, K. et al. "Use of an ordered cosmid library to deduce the genomic organization of *Mycobacterium leprae*," Mol. Microbiol. 7(2): 197-206 (1993).

Kaprelyants, Arseny S. and Kell, Douglas B. et al., "Do bacteria need to communicate with each other for growth?" Trends in Microbiology 4(6):237-42 (1996).

Kaprelyants, Arseny S. et al., "Estimation of dormant *Micrococcus luteus* cells by penicillin lysis and by resuscitation in cell-free spent culture medium at high dilution," FEMS Microbiology Letters 115(2-3):347-52 (1994).

Kunst, F. et al., "Cell wall-binding protein homolog yocH-*Bacillus subtilis*" PIR Database entry E69901, Dec. 5, 1997, accession No. E69901.

Kunst, F. et al. "The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*," Nature 390:249-56 (1997).

International Preliminary Examination Report dated Sep. 14, 1999, from International Patent Application No. PCT/GB98/01619 (filed Jun. 3, 1998).

International Search Report dated Nov. 4, 1998, from International Patent Application No. PCT/GB98/01619 (filed Jun. 3, 1998).

Jing, ER; "Study of resuscitating effect for dormant tuberculosis by recombinant resuscitation promoting factor (Rpf) proteins." *China Journal of Modern Medicine*. (19:17) 2640-43. Sep. 2009.

Harlow et al. In: Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory, Chapter 5, 60-71, 1988.

Lyristis, M. et al., "Hypothetical protein 1-*Clostridium perfringens* (fragment)," PIR Database entry S49552, No. 29, 1995, Accession No. S49552.

Lyristis, Michael et al. "Identification and molecular analysis of a locus that regulates extracellular toxin production in *Clostridium perfringens*," Mol. Microbiol. 12(5): 761-77 (1994).

Mukamolova, Galina V. et al. "A bacterial cytokine," Proc. Natl. Acad. Sci. USA 95(15): 8916-21 (1998).

Ogasawara, N. et al. "Hypothetical 47.7 KD protein in meta-ksga intergenic region," SWISSPROT Database entry Yabe-Bacsu, Oct. 1, 1994, Accession No. P37546.

Ogasawara, Naotake et al. "Systemic Sequencing of the 180 Kilobase Region of the *Bacillus subtilis* Chromosome containing the replication Origin," DNA Research 1:1-14 (1994).

Parker, W.D. et al. "Cytochrome oxidase mutations aiding diagnosis of sporadic Alzheimer's disease, Sequence 29 from Patent US 5565323," Genbank Accession No. 127393, 1997.

Votyakova, Tatyana V. et al., "Influence of Viable Cells on the Resuscitation of Dormant Cells in *Micrococcus luteus* cultures held in an Extended Stationary Phase: The Population Effect," Applied and Environmental Microbiology 60(9): 3284-91 (1994).

Skolnick, et al. Trends in Biotechnology. 18: 34-39 (2000).

Rudinger, et al. Peptide Hormones. (Ed) JA Parsons. University Park Press, pp. 1-7 (1976).

Houghten, et al. Vaccines. Cold Spring Harbor Laboratory. 86: 21-25 (1986).

The Webster's New Riverside University Dictionary, p. 933 (1984).

* cited by examiner

Primary Examiner — S. Devi

(74) Attorney, Agent, or Firm — McCarter & English, LLP; Maria Laccotripe Zacharakis

(57) **ABSTRACT**

RP-factors, their cognate receptors, convertases, respective genes and inhibitors or mimetics thereof are described. In particular, antibodies, pharmaceutical compositions and (therapeutic, diagnostic) methods based on the RP-factors and their receptors/convertases are described.

38 Claims, 23 Drawing Sheets

FIG. 1A

FIG. 1A-1
FIG. 1A-2
FIG. 1A-3

SEQ ID NO: 1	MtubZ94752	<u>mlrlvvqallllvlafaggyavaack</u> vtltvdgtamrvttmk sr vidive	50
	MtubZ94752	engfsvddrddllypaagvqvhdadtivlrrsrplqisldghda kgvwtta	100
	MtubZ94752	stvdealaqlamtdtapaaasrasrvpLsgmalpvvsaktvqlndgglvr	150
	MtubZ94752	tvhlpapnvagllsaagvpllqsdhvvpaatapivegmqiqvtrnr ikkv	200
SEQ ID NO: 2	MtubMTV008	----- <u>mpvqwlwrartakgttlknarttli</u> aaiaqt	32
	Mlep104666	----- <u>m</u> sesyrkl	8
SEQ ID NO: 4	MtubMTV043	----- <u>msgrhrkpt</u>	9
	MtubZ94752	terlplppnarvedpemnmsrevvedpgvpgtqdvtfavaevngvetgr	250
SEQ ID NO: 36	MlutZ96935	----- <u>mtlfttsat</u>	9
SEQ ID NO: 5	MlepL01095	----- <u>mpgemldvrklc</u>	12
SEQ ID NO: 6	MtubU38939	----- <u>mhlplpadhgrsrcnrhpi</u> spLslignisatsqdmssmt	38
SEQ ID NO: 7	MtubZ81368	----- <u>mtpgllttagagrprdrca</u>	19
	MtubMTV008	<u>lvttspagianaddagldpnaaagpdavgfdpnlp</u> ppapdaapvdtppape	82

FIG. 1A-1

SEQ ID NO: 8 Scoeli6C12S 25 ----ir~~ta~~avtlvaatalqatqeavaapsaplrtdWD~~AI~~AAC~~ESS~~GNWQAN

Mlep104666 58 ttssiivaki~~it~~ftgamldgsialagqaspatdsEWDQV~~AR~~CE~~SG~~GNWSIN

MtubMTV043 59 tsnvsvaki~~af~~tgavlqggq~~ia~~maaqataatdgdWDQV~~AR~~CE~~SG~~GNWSIN

MtubZ94752 300 lpvanvvt~~pa~~heavvrvgtkpgtevp~~pv~~idgsIWD~~AI~~AG~~CE~~AGGNWAIN

MlutZ96935 59 rsrratasivagmtlagaaa~~v~~gfsapagaatvdtWDR~~LA~~ECESNGTWDIN

MlepL01095 62 klfksavvsqivtas~~mal~~ststqmanavprePNWD~~AV~~AQ~~CE~~SGRNWRAN

MtubU38939 88 riakpliksamaaglv~~tas~~mslstavahagpsPNWD~~AV~~AQ~~CE~~SGGNWAAN

MtubZ81368 69 rivctvfietavvatmf~~val~~lqlst~~iss~~kaddiDWD~~AI~~AQ~~CE~~SGGNWAAN

MtubMTV008 132 dagfdpnlpplapdf~~l~~sp~~pa~~eeappvpvaysVNWD~~AI~~AQ~~CE~~SGGNWSIN

. * ** *

Scoeli6C12\$ 75 TGNGYYGGLQFARSSNIAGGLKYAPRADLATRGEQIAVAERLARLQGMS

Mlep104666 108 TGNGYLGGLQFSQGTWASHGGGEYAPSAQLATREQQIAVAERVLATQGSQ

MtubMTV043 109 TGNGYLGGLQFTQSTWAAHGGGEFAPSAQLASREQQIAVGERVLTQGRG

MtubZ94752 350 TGNGYYGGVQFDQGTWEANGGLRYAPRADLATREEQIAVAEVT~~RL~~RQGWG

MlutZ96935 106 TGNGFYGGVQFTLSSWQAVGEG----YPHQASKAEQIKRAEILQDLQGWG

MlepL01095 109 TGNGFYGGGLQFKPTIWARYGGVG----NPAGASREQQITVANRVLADQGLD

MtubU38939 135 TGNGKYGGGLQFKPATWAAFGGVG----NPAAASREQQIAVANRVLAEQGLD

MtubZ81368 116 TGNGLYGGGLQISQATWDSNGGVG----SPAAASPPQQIEVADNIMKTQGPQ

MtubMTV008 179 TGNGYYGGGLQFTAGTWRANGGSQ----SAANASREEQIRVAENVLRSQGIR

****.***** . * . ** * . ***** *** .. **

FIG. 1A-2

Scoeli6C12\$	AW-----	78
Mlep104666	AWPACGHGLSGPSLQEVLPAG---MGAPW-----	152
MtubMTV043	AWPVCGRGLSNATPREVLPA SaAMDAPldaaVNGEPAPLA-PPPADPAP	158
MtubZ94752	AWPVCAaaraGa-	362
MlutZ96935	AWPLCSQKLgltqadadagdvdateaapvavertatvqrqsaadeaaeq	156
MlepL01095	AWPKCGAASDLPIITLWHPAQGVKQIINDIIfmgdttlaialngl----	155
MtubU38939	AWPTCGAASGLPIALWSKPAQGIKQIINEIIfwagiqasfpr-----	176
MtubZ81368	AWPKCScsqgdaplgslthlftlaaetggcsgrdd-----	154
MtubMTV008	AWPVCGrrg-----	188
	*** *	
Mlep104666	pqqpadnf-----pPTPGDVPSPLarp-----	174
MtubMTV043	pvelaandlpaplgeplpaapadpappadlaFPAPADVAPPvelavndlp	208
MlutZ96935	aaaaeqavvaeaetivvksgdslwtlaneyeveggwtalyeankgavsd	206
MtubMTV043	aplgeplpaapadpappadlappapadlappapadlappvel	258
MlutZ96935	aviyvqqelvlpqa-----	220
MtubMTV043	avndlpaplgeplpaapaelappadlapasadlappapadlappapaela	308
MtubMTV043	ppapadlappaavneqtapgdqpatapggpgvlgatdlelpepdpgpadap	358
MtubMTV043	ppgdrvteapaetpgvsniaYtkklwqairaqdvvcgndaldslagpyvig-	407

FIG. 1A-3

FIG. 1B

FIG. 1B-1
FIG. 1B-2

SEQ ID NO: 9	A	YabEBsubt	mgeregrvdslltdtlynlseekeaffitqmkklfsvklSKSVILVAACLLAGSGTAYAAHELTQSVSVSINGKKHHR	82
SEQ ID NO: 1	A	Mtub294752	-----MLRLVUGALLLVLAFAAGG-YAVAACKTVTLTVDGTAMR--VF	39
SEQ ID NO: 10	A	YochBSsubt	-----mkkktimsfvav	11
	A	YabEBsubt	THANTVGDLLLETLDIKTRDEDKITPAKQTKITADHDVVYEAAKPVKLTING-EETLHSTAKTVGALLDEQDVVDVKEQQQID	163
	A	Mtub294752	TMKSRVIDIVEENGFSVDDRDDLYPAAGVQVHDADTIVLRRSRPLQISLDGHDAKQVHTASTVDEALAQLANTDTPAFAAS	121
SEQ ID NO: 11	A	Caceto506	-----KR? ?AVILMVAVIPTIISSHKKNITVNIIDG-KTSKIITYKKSNEGSIKNNILVGPDKIQ	58
			##	

FIG. 1B-1

FIG. 1B-2

FIG. 1D

FIG. 1D-1
FIG. 1D-2

SEQ ID NO: 4 1 msgrhrkpttsnvsvakiaftgavlpgggiamaaqataatdgewdqvarcesggnwsintgngylgg
lqftqstwaahgggefapsaqilasreqgiavgerlvl atqgrgawpvcgrgl snatprevlpasaamd
apldaaaavngepaplapppadp 156

157 appvelaandlpaplgaplpaapadpappadlappapadv 196

197 appvelavndlpaplgaplpaapadpappadlappapadlappapadl 252

253 appvelavndlpaplgaplpaapaelappadlap-asadlappapadlappapaelappapadlappa
320 -----avne 323

324 qtapgqdqpatapgpgvglatdlelpepdpqpada pppgdvteapaetpgvsnia ytkklwqaira
389 qdvvcgndaldslaqpyvig* 407

Motif	sequence
A	157 appvelaandl 167 SEQ ID NO: 25
B'	168 paplgaplpaapad 181 SEQ ID NO: 28
C	182 pappadl 188 SEQ ID NO: 29
D	189 appapadv 196 SEQ ID NO: 31

FIG. 1D-1

A 197 appvelavndl 207 SEQ ID NO: 26
B' 208 paplgeplpaapad 221 SEQ ID NO: 28
C 222 pappadl 228 SEQ ID NO: 29
D 229 appapadl 236 SEQ ID NO: 30
D 237 appapadl 244 SEQ ID NO: 30
D 245 appapadl 252 SEQ ID NO: 30

A 253 appvelavndl 263 SEQ ID NO: 26
B 264 paplgeplpaapael 278 SEQ ID NO: 27
'C 279 appadl 284 SEQ ID NO: 55
D* 285 apasadl 291 SEQ ID NO: 56
D 292 appapadl 299 SEQ ID NO: 30
D 300 appapael 307 SEQ ID NO: 57
D 308 appapadl 315 SEQ ID NO: 30
D' 316 appa 319 SEQ ID NO: 58

'A' 320 avne 323 SEQ ID NO: 59

A = appvela[av]ndl

B = paplgeplpaapa[de]l

C = pappadl

D = appapa[de][lv]

FIG. 1D-2

SEQ ID NO:34 Lmonocytoγ...
SEQ ID NO:36 MlutFactor

mmkkaiaa¹tagiav²tafaapt³iasastv⁴vveag⁵dtl⁶wgiaqskgt⁷tvdaikkannl⁸tt⁹dkivpggk¹⁰iqv¹¹n¹² 72
mt¹³fttsatrs¹⁴rratasi¹⁵v¹⁶g¹⁷mt¹⁸lagaa¹⁹v²⁰gsapagaat²¹-----v²²tdwrlae²³desng²⁴tw²⁵dintgn²⁶ 62

Lmonocytoγ...
MlutFactor

nevaaa¹ek²teks³vsatw⁴lnvrtgagv⁵dn⁶si⁷tsi⁸kggk⁹vtvettesng¹⁰whk¹¹ityndg¹²ktg¹³f¹⁴vn¹⁵gkyl¹⁶tdka¹⁷ 144
gfyggv¹⁸gf¹⁹tl²⁰sswg²¹lv²²ggeg²³yphq²⁴---askae²⁵qik²⁶raei²⁷lq²⁸dl²⁹ggw³⁰awp³¹lcsq³²kl³³gt³⁴cadadag³⁵----- 125

Lmonocytoγ...
MlutFactor

vstp¹vaptqev²kk³ETTT⁴QQA⁵PVA⁶ETK⁷TEV⁸KQT⁹QA¹⁰TPAP¹¹KVA¹²ETK¹³ETP¹⁴V¹⁵ID¹⁶QNA¹⁷THA¹⁸VKS¹⁹GDT²⁰I²¹VA²²DSV²³ 216
-----D²⁴VDA²⁵TEA²⁶APVA²⁷VERT²⁸ATV²⁹QRQ³⁰SA³¹DEA³²AAEQ³³AAAAEQ³⁴AV³⁵AE³⁶ET³⁷I³⁸VKSG³⁹DSL⁴⁰VT⁴¹DAN⁴² 184

Lmonocytoγ...
MlutFactor

KYGV¹svqdim²sw³nnl⁴-----SSSS⁵IYVG⁶QKDA⁷IKQ⁸Tantat⁹pk¹⁰ae¹¹vk¹²teap¹³aaek¹⁴qa¹⁵pv¹⁶vkent¹⁷nt¹⁸tatt¹⁹ 283
EYEV²⁰eggwt²¹alye²²ankg²³avs²⁴DA²⁵AV²⁶IYVG²⁷QED²⁸VL²⁹PQA³⁰----- 220

Lmonocytoγ...
MlutFactor

ekketat¹qqqtap²kap³tea⁴ak⁵pap⁶ap⁷st⁸nt⁹n¹⁰ank¹¹tn¹²tn¹³tn¹⁴tn¹⁵tn¹⁶tn¹⁷tn¹⁸tn¹⁹tn²⁰tn²¹tn²²tn²³tn²⁴tn²⁵tn²⁶tn²⁷tn²⁸tn²⁹tn³⁰ng³¹ss³² 355
----- 220

Lmonocytoγ...
MlutFactor

nnns¹ssas²ai³ae⁴aq⁵hl⁶g⁷k⁸ays⁹w¹⁰gg¹¹ng¹²pt¹³fd¹⁴c¹⁵g¹⁶ty¹⁷ky¹⁸v¹⁹fak²⁰ag²¹isl²²prt²³sg²⁴aq²⁵ya²⁶st²⁷tr²⁸ises²⁹qak³⁰pg³¹ 427
----- 220

Lmonocytoγ...
MlutFactor

dlv¹ffdy²gsg³ish⁴v⁵gi⁶yv⁷ng⁸g⁹mina¹⁰q¹¹dn¹²g¹³ky¹⁴dn¹⁵ih¹⁶g¹⁷sg¹⁸w¹⁹g²⁰kyl²¹vg²²fgrv²³ 478
----- 220

FIG. 1E

FIG. 2A

FIG. 2A-1
FIG. 2A-2

SEQ ID NO: 35 1 accaaggagaaggacaccccggtgtgcctcggccgcatcagcgaggactcgccatgg 60
61 acaccatgactctcttcaccacttcgccacccgctcccgccgtgccaccgctcgatcg 120
M T L F T T S A T R S R R A T A S I V
SEQ ID NO: 37 g
121 tcgcgggcatgacctcgccggcgccgcgctgggcttctccgccccgagggcg 180
A G M T L A G A A A V G F S A P A Q A A
SEQ ID NO: 38 A
oligo A1>>>
csacsgtsgacacstgggaccgscstsgcsgag
181 ccaccgtggacacctgggaccgctcgccgagtgcgagtcacaacggcacctgggacatca 240
T V D T W D R L A E C E S N G T W D I N
T V D T W D R L A E E X S N G T X D
SEQ ID NO: 39 <<< oligo G2 SEQ ID NO: 40 oligo G1>>>
gttgccgaagatgcggcc agttcacctgtcctcctg
241 acaccgggaacggcttctacggcgggcggtgcagttcacctgtcctccttggcaggccgctg 300
T G N G F Y G G V Q F T L S S W Q A V G
SEQ ID NO: 42 G

FIG. 2A-1

SEQ ID NO: 41 <<< Oligo A2
 ccictycciatrvggigtgtycg
301 gcggcgaaggctaccgcaccaggcctcgaaaggccgagcagatcaagcgcgcgagatcc 360
 G E G Y P H Q A S K A E Q I K R A E I L
 G E G Y P H Q A S K
361 tccaggacctgcagggtggggtggcgctggccgctgtgtgtgcagaagctgggcctgacct 420
 Q D L Q G W G A W P L C S Q K L G L T Q
421 aggtgacgcggacgcgggtgacgtggacgcccaccgagggcccggtcgccgtggagc 480
 A D A D A G D V D A T E A A P V A V E R
481 gcacggccacctgcagcgcagtcctgcgcgcgacgaggtgcgcgagcagcagcgcgctg 540
 T A T V Q R Q S A A D E A A E Q A A
541 ccgcggagcagccgctcgtcgcgcgagggccgagaccatcgtcgaagtcgggtgactccc 600
 A E Q A V V A E A E T I V V K S G D S L
601 tctggacgctcgccaacgagtagtacgaggtggaggtggctggaccgcccctctacgagcca 660
 W T L A N E Y E V E G G W T A L Y E A N
661 acaagggtcgtctccgacgcccgcgtgatctacgtcggccaggagctcgtcctgcgcg 720
 K G A V S D A A V I Y V G Q E L V L P Q
721 aggcctgagacgcctgacggcccccccgaccggtacc 758
 A *

SEQ ID NO: 43 1 ATVDTWDLA ECESNGTWDI NTGNIFYGGV QFTLSSWQAV GGEYPHQAS KAEQIKRAEI 60
 61 LQDLQGWGAW PLCSQXLGLT QADADAGDVD ATEAAPVAVE RTATVQRQSA ADEAAAEQAA 120
 121 AAEQAVVAEA ETIVVXSGDS LWTLANEYEV EGGWTALYEA NKGAVSDAAV IYVGQELVLP QA 182

FIG. 2A-2

SEQ ID NO:44	ggatccgcaccgcccgttaacctggtcgccgcgcacgcgcactcggggcgacccggcggaag	60
SEQ ID NO:45	I R T A A V T L V A A T A L G A T G E A	
	cggtgccgcgccctcggcgccccctgcgcaccgactgggacgcccatcgccgcgtgcgagt	120
	V A A P S A P L R T D W D A I A A C E S	
	ccagcggcaactggcaggcggaacacccggcaacggctactacggggcctgcagttcgcac	180
	S G N W Q A A N T G N G Y Y G G L Q F A R	
	ggtccagctggatcgccgcggcgccctcaagtagcccccgcgcggacctcgccacccc	240
	S S W I A A G G L K Y A P R A D L A T R	
	gcggcgagcagatcgccgtggcggaacgcctcgcccgtctgcagggggatgtccgcctgg	299
	G E Q I A V A E R L A R L Q G M S A W	

FIG. 2B




FIG. 3A




FIG. 3B

FIG. 3C





FIG. 3D




FIG. 4A




FIG. 4B




FIG. 5C




FIG. 5B




FIG. 5A




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 8A




FIG. 8B




FIG. 9A




FIG. 9B




FIG. 10

1

BACTERIAL PHEROMONES AND USES THEREOF

RELATED INFORMATION

This application is a divisional application of U.S. application Ser. No. 09/445,289, filed May 11, 2000, pending, which is the U.S. national phase application, pursuant to 35 U.S.C. §371, of international patent application Ser. No. PCT/GB98/01619, filed Jun. 3, 1998, designating the United States and published in English as WO 98/55624 on Dec. 10, 1998, which claims priority to GB 97113898.8, filed Jun. 4, 1997 and GB 9811221.2, filed May 27, 1998, the disclosures of each of which applications are incorporated herein in their entirety by this reference.

The sequence listing in the 57,763 byte text file "2011-10-04_118160-00302_ST25.txt" that was created Oct. 4, 2011 and filed electronically by EFS-Web on Oct. 4, 2011 is incorporated herein by reference.

FIELD OF THE INVENTION

The present invention relates to RP-factors, their cognate receptors, convertases, respective genes and to inhibitors or mimetics thereof. In particular, the invention relates to antibodies, pharmaceutical compositions and (therapeutic, diagnostic) methods based on the RP-factors and their receptors/convertases.

INTRODUCTION

Bacterial Pheromones

It is known that certain chemicals may mediate intercellular communication in bacterial cultures. Such communication has been shown to be of importance during sporulation, conjugation, changes in virulence and in bioluminescence. It is now clear that a variety of different autocrine and/or paracrine chemical compounds ("pheromones") produced as secondary metabolites are responsible for such social behaviour in prokaryotes (see e.g. 25 Kell et al., 1995, Trends Ecol. Evolution, 10, 126-129).

Pheromones may be distinguished from nutrients inter alia in that: (i) they are produced by the organisms themselves, (ii) they are active at very low concentrations (e.g. at picomolar or nanomolar concentrations), and (iii) with the exception of prohormone processing, their metabolism is not necessary for activity (although they may of course ultimately be degraded).

The chemical nature of these pheromonal compounds varies widely: those associated with Gram-negative organisms tend to be of low molecular weight (e.g. N-acyl homoserine lactone derivatives), whilst a number of Gram-positive organisms use proteins and polypeptides (Kell et al, 1995, ibidem).

Pheromones are also known to play an important role in the development of bacterial cultures. For unstressed (uninjured) bacteria and optimal growth media, the "self-promoting" mode of culture growth is normally masked due to the high rate of production of growth factors and the sensitivity of the cells to these pheromones. Only under unfavourable conditions (for example, poor growth media, small initial inocula and/or starved cells) is this self-promoting behaviour "visible".

For example, a dramatic reduction in the length of the lag phase of cultures of *Nitrosomonas europaea* is mediated by N-(3-oxo-hexanoyl) homoserine lactone, and chorionic gonadotropin-like ligand (a 48 kD protein) had similar growth-stimulating activity for *Xanthomonas maltophilia*. A

2

number of mammalian hormones (including peptide and steroid hormones as well as cytokines) have also been shown to exhibit potent growth-stimulating activities for both Gram-positive and Gram-negative bacteria.

Latency and Resuscitation

The ability of a microbial cell to grow and divide on a nutrient agar plate constitutes the benchmark method for determining the number of living cells in a sample of interest. However, it is also widely recognised that, especially in nature, the distinction between life and non-life is not absolute; many cells may exist in "dormant" or "moribund" forms or states and will not produce colonies on nutrient media (i.e. are "non-culturable"). However, these dormant or latent cells are not dead: they can be returned, by a process known as resuscitation, to a state of viability/culturability.

For example, it is known that cells of the (high-G+C Gram-positive) bacterium *Micrococcus luteus* can enter a state of true dormancy from which they may be resuscitated by culture supernatants, even in the absence of any 'initially viable' cells.

The latent state has profound medical implications: many pathogenic bacteria (including pathogenic mycobacteria such as *M. tuberculosis*) are known to persist for extended periods in latent states in a host organism. Indeed, tuberculosis is a re-emergent infection of great concern, and it is recognised in particular that the causative organism (*Mycobacterium tuberculosis*) can lie dormant (remain latent) in patients and carriers for periods of years.

The latent state also has important commercial implications, since it complicates many laboratory methods for the detection, cultivation and enumeration of bacteria (for example in the food and healthcare industries).

There is therefore a pressing need to understand the physiological bases of latency and resuscitation.

SUMMARY OF THE INVENTION

The present invention is based, at least in part, on the discovery of a new class of pheromones which stimulate the resuscitation of bacteria after true dormancy. This "resuscitation factor" (herein embraced by the term "RP-factor") may exhibit activity at picomolar concentrations (implying a non-nutritional role). The elucidation of the structure of the pheromones at the amino acid sequence level has also permitted the present inventors to describe a larger family of proteins, some members of which act more broadly as regulators of cellular growth or replication and not necessarily as resuscitation promoting factors. Further sequence comparisons have also led to the identification of the cognate receptors, at least some of which share certain sequence similarities with their cognate RP-factors.

Thus, in a first aspect of the present invention there is provided an isolated RP-factor.

RP-Factors

The term "RP-factor" is used herein to encompass any representative of that family of substances the members of which are capable of resuscitating dormant, moribund or latent cells (e.g. dormant bacterial cells). In addition, the RP-factors of the invention may also exhibit growth-stimulatory activity with respect to growing cells (e.g. growing bacterial cells), and/or may be competent to reduce the lag time of cell (e.g. bacterial cell) cultures. The resuscitation activity (and optionally also the growth-stimulatory activity or lag-time reducing activity) of the RP-factor may be specific for a particular (bacterial) cell (e.g. specific for one or more patho-

genic mycobacteria), or may be non-specific. Specificity may be manipulated for example by engineering (e.g. by mutagenesis or chimaerisation, as herein described) of the specificity-determining domain(s) of the RP-factor or by replacement of the signalling domain.

The term "RP-factor" is also used herein in a somewhat broader sense to encompass polypeptides which are expressed by bacteria and which regulate (e.g. promote, trigger, prevent or impair) the growth or multiplication of a cell (the "target cell") by acting as signalling moieties in conjunction with (e.g. by binding to) cognate cellular receptors. Such polypeptides may be referred to herein as bacterial cytokines.

The RP-factors of the invention therefore include bacterial cytokines which may or may not be capable of resuscitating dormant, moribund or latent cells (e.g. dormant bacterial cells) and/or exhibit growth-stimulatory activity with respect to growing cells (e.g. growing bacterial cells). They may or may not also be competent to reduce the lag time associated with the growth of cell (e.g. bacterial cell) cultures. Moreover, some bacterial cytokines which fall within the scope of the term "RP-factor" as defined herein may even prevent or impair the growth of the target cells (particularly where the target cells are eukaryotic (e.g. mammalian) cells).

The RP-factors of the invention may fall into at least two functional classes: aut signalling factors and allosignalling factors. Aut signalling factors act to regulate the growth of the bacterial cell in which they were expressed (i.e. they act as bacterial autocrine factors), while allosignallers act to regulate the growth of other cells (i.e. they act as bacterial paracrine factors). Aut signalling factors therefore act as self-regulators of bacterial cell growth, and may be essential for viability and/or growth. Some RP-factors may function as both auto- and allosignalling cytokines.

Allosignalling factors may exhibit a range of different specificities. Some may act solely on other bacterial cells of the same species as the cell in which they were expressed ("homoactive" factors), while others may act on cells of one or more other species ("heteroactive" factors). Heteroactive factors may exhibit a broad range of specificity: they may act on several different species (for example, in a genus-specific manner), or may be species-specific. Some heteroactive bacterial factors may act on eukaryotic cells, and may be specific for particular cell-types. For example, some heteroactive bacterial cytokines (particularly those produced by certain pathogens) may act on mammalian cells (e.g. mammalian epithelial, endothelial or immune cells), and may be tissue- or cell-type specific.

Notwithstanding the above explanation, it is postulated that the specificity of at least some RP-factors may be concentration dependent. In these cases, the specificity of any given RP-factor falls within a continuum, so that an aut signalling RP-factor may mediate cross-talk and so exhibit allosignalling activity when present at sufficiently high concentrations. Similarly, allosignalling RP-factors may exhibit homo- or heteroactivity depending on concentration.

The RP-factor may be translocated through the cell membrane, whereafter it may be secreted into the surrounding environment or remain associated with the surface of the cell. Thus, at least two classes of RP-factor may exist: secreted and non-secreted. The secreted RP-factors are characterised by the presence of a secretory signal sequence (the presence of which is readily recognised by those skilled in the art on the basis of the presence of DNA and/or amino acid sequence motifs). The non-secreted RP-factors may be cell-associated or cytosolic factors. Both classes of RP-factor may exist in a single cellular source (e.g. in a single bacterial source). Both classes of RP-factor find application in the invention.

Non-secreted RP-factors may act in at least four different ways: (a) as a membrane-anchored juxtacrine factor mediating a growth regulating signal between two different cells in close physical proximity or contact; and/or (b) as an intercellular signalling moiety upon cleavage by an enzyme (e.g. a convertase, as herein defined) which releases a soluble signalling moiety into the extracellular milieu; and/or (c) as an autocrine factor via binding to cognate receptors located on the surface of the cell in which the non-secreted factor is expressed or acting entirely intracellularly; and/or (d) as a cognate receptor for another non-secreted or secreted RP-factor.

Thus, the RP-factors of the invention may include the nine factors identified by the sequences shown in FIG. 1A and the five factors identified by those shown in FIG. 1B, together with their species variants, allelic forms, homologues, derivatives, muteins and corresponding secreted/nonsecreted forms (vide infra).

Preferably, the RP-factors of the invention are species variants, allelic forms, homologues, derivatives, muteins and corresponding secreted/nonsecreted forms of any one of the nine factors identified by the sequences shown in FIG. 1A and the five factors identified by those shown in FIG. 1B.

The RP-factors may be synthesised in the form of a precursor which is processed to produce a mature form. Such processing may proceed via various intermediate (pro-) forms. Such precursors, intermediate forms and mature proteins are all intended to be covered by the term "RP-factor" as used herein, except where indicated otherwise. As used herein, the term "pro-RP-factor" specifically defines any of various precursors (which may or may not be active) of a mature RP-factor.

The processing may comprise proteolytic cleavage and/or secretion. The precursors may be inactive, and become active on processing as a mature form. The precursors may comprise proteins having secretory leader sequences which are removed during secretion (pre-forms). Such forms are herein referred to as "pre-RP-factor or pre-pro-RP-factors". As explained above, such pre- or prepro-forms are also intended to be covered by the term "RP-factor" as used herein, except where indicated otherwise.

Processing may be attendant on the binding of an RP-factor precursor to a cognate receptor. Such receptors may then directly (or indirectly) cleave the precursor to produce a more mature form of the RP-factor. Such processing may occur as a cascade, involving several receptor-processing complexes, and so ultimately result in the production of a mature RP-factor which then acts as a signalling moiety by binding to a terminal (signal transducing) receptor.

In such processing, the proximal (or intermediate) receptors may function as convertases, and the terminal receptor as a signal transducer. However, a receptor may function as both a convertase and a signal transducer. As used herein, the term "convertase" is intended to define a molecule which binds an RP-factor precursor and (directly or indirectly) processes it to produce a more mature form. They may, for example, have protease activity.

The receptors/convertases discussed above may be disposed at the cell surface (e.g. membrane bound), cytosolic or extracellular.

Preferably, the RP-factor is derived from a bacterium (e.g. a pathogenic bacterium). Particularly preferred are RP-factors derived from high G+C Gram-positive bacteria.

However, the inventors have also discovered RP-factor family members in representatives of the low G+C Gram-positive organisms, including *Bacillus subtilis* and clostridia. Thus, RP-factors derived from low G+C Gram-positive bac-

5

teria (e.g. pathogenic low G+C Gram-positive bacteria) are also preferred according to the invention. Examples of the latter include: *Streptococcus* spp., *Staphylococcus* spp., *Listeria* spp., *Bacillus* spp., *Clostridium* spp. and *Lactobacillus* spp.

The invention also contemplates homologues, allelic forms, species variants, derivatives, muteins or equivalents of the RP-factors and RP-factor receptors/convertases of the invention.

Preferably, the homologues, derivatives, muteins or equivalents of the RP-factor of the invention have at least 20% identity with any one of the particular amino acid sequences shown in FIGS. 1A and 1B.

Particularly preferred are homologues, derivatives, muteins or equivalents of the RP-factor of the invention which have at least 30% identity, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity with any one of the particular amino acid sequences shown in FIGS. 1A and 1B.

The homologues, derivatives, muteins or equivalents of the RP-factor of the invention may have at least 25% homology with any one of the particular amino acid sequences shown in FIGS. 1A and 1B.

Particularly preferred are homologues, derivatives, muteins or equivalents of the RP-factor of the invention which have at least 30% homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% homology with any one of the particular amino acid sequences shown in FIGS. 1A and 1B.

The invention also contemplates chimaeric RP-factors. These are factors which comprise one or more heterologous domains. In this context, a heterologous domain is a portion of an RP-factor which is derived from a different RP-factor to that from which the other domain(s) with which it is associated are derived. Such chimaeric RP-factors find particular utility in applications where the specificity and/or activity of the RP-factor is manipulated or altered.

The invention also contemplates all individual functional domains of the RP-factors of the invention as separate and independent entities.

The invention also contemplates recombinant RP-factor. As used herein, the term "recombinant" is intended to define material which has been produced by that body of techniques collectively known as "recombinant DNA technology" (for example, using the nucleic acid, vectors and or host cells described infra).

Cognate Receptors

In some cases, the cognate cellular receptor is a cell surface receptor; in other cases, it is a cytosolic receptor with which the cytokine interacts after uptake by the target cell. The receptors with which the RP-factors and/or bacterial cytokines of the invention interact may share certain structural motifs with the RP-factors/cytokines themselves. In particular, the receptors may contain a ligand binding domain which is structurally similar to the signalling domain of the cognate RP-factor/cytokine.

The receptors may also comprise a membrane anchor domain and a wall spanning domain.

Preferably, the cognate receptor comprises a receptor domain as hereinbelow defined and/or a wall spanning domain as hereinbelow defined and/or a membrane anchor.

Particularly preferred are cognate receptors comprising the amino acid sequence of MtubZ94752 as shown in FIG. 1A or the amino acid sequence of YabE from *B. subtilis* as shown in FIG. 1B.

6

The cognate receptors may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The cognate receptors may also comprise derivative or equivalent sequences of amino acid residues which have at least 20% identity or homology with the amino acid sequence of MtubZ94752 as shown in FIG. 1A or the amino acid sequence of YabE from *B. subtilis* as shown in FIG. 1B, for example at least 30% identity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

RP-Factor/Cognate Receptor Domain Structure

The RP-factors of the invention (including the bacterial cytokines as also defined herein) and their cognate receptors may comprise a plurality of discrete domains. These domains may be functionally and/or structurally distinct.

The RP-factors of the invention may be characterised by the presence of at least two functional domains: a secretory signal sequence (which may be wholly or partially absent in the active form of the factor) and a signalling domain. The signalling domain may fall into one of at least two distinct classes described in more detail infra.

Many RP-factors also comprise a third functional domain which mediates a physical association with the surface of the target cell (hereinafter referred to as the "localizing domain" and described in more detail infra).

The RP-factors of the invention may further comprise a specificity-determining domain, which may function in conjunction with the signalling domain.

Non-secreted RP-factors may further comprise a wall-spanning domain (described in more detail infra) and/or a membrane anchor.

The gross structure and/or amino acid sequence of the aforementioned domains may vary considerably. In particular, the structure of the surface localizing domain may differ according to the structure of the cell-wall of the target cell. For example, the surface localizing domain may fall into one of at least two distinct classes: class I (which may act on peptidoglycan) and class II (which may act on the outer lipid envelope found in mycobacteria).

The cognate receptors of the invention may be characterised by the presence of at least two functional domains: a receptor domain and a wall spanning domain. They may also comprise a membrane anchor. The receptor domain may be structurally similar to the signalling domain of the cognate RP-factor (as described in more detail infra).

Receptor/Signalling Domain, class I

This domain may be associated with RP-factors from high G+C Gram-positive bacteria (such as mycobacteria and *Micrococcus* spp.) and/or their cognate receptors. When present on RP-factors, the domain may be involved in receptor binding, and may for example bind a structurally similar domain on a cognate receptor. Thus, when present as part of an RP-factor of the invention, the domain is termed the "signalling domain", and when present in the cognate receptor, the domain is termed the "receptor domain".

The domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues indexed by asterisks in any one of the 9 sequences set out in FIG. 1A.

In preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues indexed by asterisks and dots in any one of the 9 sequences set out in FIG. 1A.

In particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues highlighted in bold upper case type in any one of the 9 sequences set out in FIG. 1A.

In more particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues presented in upper case type in any one of the 9 sequences set out in FIG. 1A.

The domain may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The domain may also comprise derivative or equivalent sequences of amino acid residues which have at least 20% identity or homology with any one of the particular amino acid sequences defined above, for example at least 30% identity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

Receptor/Signalling Domain, Class II

This domain may be associated with RP-factors from low G+C Gram-positive bacteria (such as bacilli and clostridia) and/or their cognate receptors. When present on RP-factors, the domain may be involved in receptor binding, and may for example bind a structurally similar domain on a cognate receptor. Thus, when present as part of an RP-factor of the invention, the domain is termed the "signalling domain", and when present in the cognate receptor, the domain is termed the "receptor domain".

The domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues indexed by asterisks in any one of the 5 sequences set out in FIG. 1B(B).

In preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues indexed by asterisks and dots in any one of the 5 sequences set out in FIG. 1B(B).

In particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues highlighted in bold upper case type in any one of the 5 sequences set out in FIG. 1B(B).

In more particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues presented in upper case type in any one of the 5 sequences set out in FIG. 1B(B).

The domain may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The domain may also comprise derivative or equivalent sequences of amino acid residues which have at least 20% identity or homology with any one of the particular amino acid sequences defined above, for example at least 30% iden-

tity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

Wall Spanning Domain

This domain may be associated with non-secreted RP-factors (e.g. cell-associated RP-factors or RP-factors which act as juxtacrine factors) and with the cognate receptors of the RP-factors of the invention. When present, the domain is involved in mediating an interaction with the cell wall such that the RP-factor/receptor as a whole may span it.

The wall spanning domain may therefore be bounded by cytosolic and extracellular regions in vivo. The domain is often associated with a membrane anchor, the two structural elements acting in concert to maintain the RP-factor/receptor at the cell surface.

The domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues presented in upper case and indexed by hashes (#) in any one of the 5 sequences set out in FIG. 1B(A).

In preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues presented in upper case and indexed by hashes and dots in any one of the 5 sequences set out in FIG. 1B(A).

In particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues highlighted in bold upper case type in any one of the 5 sequences set out in FIG. 1B(A).

In more particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues presented in upper case type in any one of the 5 sequences set out in FIG. 1B(A).

The domain may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The domain may also comprise derivative or equivalent sequences of amino acid residues which have at least 20% identity or homology with any one of the particular amino acid sequences defined above, for example at least 30% identity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

Localizing Domain, Class I

This domain may be present in secreted RP-factors, and may mediate a physical association with the surface of the target cell by acting to bind peptidoglycan or some other surface component(s). It may therefore act to increase the local concentration of the cytokine at the target cell surface, so promoting activity by increasing the local concentration of RP-factor in the immediate vicinity of the cognate receptor. Localizing domains may therefore be a characteristic feature of allosignalling bacterial cytokines, and may be absent in aut signalling factors or vice versa. For example, when present in aut signalling factors, localizing domains may act to retain the factor at or near the cell surface after secretion through the cell membrane.

When present, the localizing domain may confer important binding properties on the RP-factor, whereby binding to cognate receptor is biphasic and characterised by a primary (relatively unspecific) association with the cell surface followed by a secondary (relatively highly specific) association with the cognate receptor.

The domain may comprise a sequence of amino acid residues, the identity and relative 25 positions of which correspond to those residues indexed by asterisks in any one of the 10 sequences set out in FIG. 1C.

In preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues indexed by 30 asterisks and dots in any one of the 10 sequences set out in FIG. 1C.

In particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues highlighted in bold upper case type in any one of the 10 sequences set out in FIG. 1C.

In more particularly preferred embodiments, the domain may comprise a sequence of amino acid residues, the identity and relative positions of which correspond to those residues set out in any one of the 10 sequences set out in FIG. 1C.

The domain may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The domain may also comprise derivative or equivalent sequences of amino acid 15 residues which have at least 20% identity or homology with any one of the particular amino acid sequences defined above, for example at least 30% identity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

Localizing Domain, Class II

This domain may be present in secreted RP-factors, and may mediate a physical association with the surface of the target cell by acting to bind the outer lipid envelope present in mycobacteria. It may therefore act to increase the local concentration of the cytokine at the target cell surface, so promoting activity by increasing the local concentration of RP-factor in the immediate vicinity of the cognate receptor. Localizing domains may therefore be a characteristic feature of allosignalling bacterial cytokines, and may be absent in autologous factors.

When present, the localizing domain may confer important binding properties on the RP-factor, whereby binding to cognate receptor is biphasic and characterised by a primary (relatively unspecific) association with the cell surface followed by a secondary (relatively highly specific) association with the cognate receptor.

The domain may comprise an alanine plus proline-rich segment, such as one or more of the amino acid motifs depicted in FIG. 1D as 'A' (SEQ ID NO: 59), A, B, B' (SEQ ID NO: 63), C (SEQ ID NO: 29), 'C' (SEQ ID NO: 55), D, D* (SEQ ID NO: 56) and D' (SEQ ID NO: 58) (any one of which may be tandemly repeated). D. Motifs A, B and D are depicted in FIG. 1D with brackets around two amino acids, to indicate that the motifs define sequences that include a choice of one or the other of the two amino acids within the brackets as follows:

A = appv~~e~~la[av]ndl; (SEQ ID NO: 62)
 B = paplgeplpaapa[de]l; (SEQ ID NO: 60)
 and
 D = appapa[de][lv]. (SEQ ID NO: 61)

In preferred embodiments, the domain may comprise a sequence of amino acid residues corresponding to residues 158-322 of MtubMTVO43 as shown in FIG. 1D or to that of residues 45-112 of MtubMTV008 as shown in FIG. 1A.

The domain may also comprise derivative or equivalent sequences of amino acid residues which are as defined above but in which amino acids have been added, deleted or substituted (e.g. conservatively substituted), provided that biological activity (e.g. signalling or ligand-binding activity) is substantially retained.

The domain may also comprise derivative or equivalent sequences of amino acid residues which have at least 20% identity or homology with any one of the particular amino acid sequences defined above, for example at least 30% identity or homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity or homology therewith.

The term "isolated" is used herein to indicate that the factor exists in a physical milieu distinct from that in which it occurs in nature. For example, the isolated factor may be substantially isolated with respect to the complex cellular milieu in which it naturally occurs. The absolute level of purity is not critical, and those skilled in the art can readily determine appropriate levels of purity according to the use to which the factor is to be put.

In many circumstances, the isolated factor will form part of a composition (for example a more or less crude extract containing many other proteins and substances), buffer system or pharmaceutical excipient, which may for example contain other components (including other proteins, such as albumin).

In other circumstances, the isolated protein may be purified to essential homogeneity, for example as determined by PAGE or column chromatography (for example HPLC or mass spectrometry). In preferred embodiments, the isolated RP-factor of the invention is essentially the sole active RP-factor in a given composition. Particularly preferred are compositions in which an RP-factor (or a particular species, homologue, mutein, derivative or equivalent thereof) is present as the sole active ingredient in a pharmaceutical composition.

The RP-factor for use in the invention need not be isolated in the sense defined above, however. For example, more or less crude culture supernatants (e.g. "spent" medium) may contain sufficient concentrations of RP-factor for use in several applications. Preferably, such supernatants are fractionated and/or extracted (see below), but in many circumstances they may be used without pretreatment. They are preferably derived from spent media used to culture RP-factor-producing microorganisms (for example, the bacterial sources described *infra*). The supernatants are preferably sterile. They may be treated in various ways, for example by concentration, filtration, centrifugation, spray drying, dialysis and/or lyophilisation. Conveniently, the culture supernatants are simply centrifuged to remove cells/cell debris and filtered.

Such supernatants find utility in diagnostic kits and methods, for example in the diagnostic kits and methods described *infra*. They also find utility in the recovery from various samples of culturable microorganisms (e.g. from soil, food, marine, freshwater, or tissue samples) or from samples taken from an organism (e.g. a human or animal).

The culture supernatants may also be used as supplements in various culturing substrates, for example in culture or transport media. The culture medium may take any convenient form, such as for example agar plates, broths, slopes, coated dipsticks, coated probes, membranes, coated or filled wells or films. The medium may be a defined or complex

medium, and may contain indicator dyes to facilitate identification of cultured microorganisms. Preferably, the medium is suitable for the culturing or transport of bacteria, for example *Mycobacterium* spp.

The term "isolated" as applied to the other materials of the invention (for example, the genes and other nucleic acids encoding the RP-factor and their cognate receptors/convertases) is to be interpreted mutatis mutandis. Thus, as applied to nucleic acid (e.g. RNA or DNA or (structural) genes), the isolated nucleic acid may be present in any of a wide variety of vectors and in any of a wide variety of host cells (or other milieu, such as buffers, viruses or cellular extracts).

The term "family", as applied to the proteins of the invention, is used herein to indicate a group of proteins which share substantial sequence similarities, either at the level of the primary sequence of the proteins themselves, or at the level of the DNA encoding them. The sequence similarities may extend over the entire protein/gene, or may be limited to particular regions or domains. Similarities may be based on nucleotide/amino acid sequence identity as well as similarity (for example, those skilled in the art recognise certain amino acids as similar, and identify differences based on switches of similar amino acids as conservative changes). Some members of a protein family may be related in the sense that they share a common evolutionary ancestry, and such related proteins may herein be referred to as homologues. The members of a protein family do not necessarily share the same biochemical properties or biological functions, though their similarities are usually reflected in common functional features (such as effector binding sites and substrates).

The criteria by which protein families are recognised are well-known in the art, and include computer analysis of large collections of sequences at the level of DNA and protein as well as biochemical techniques such as hybridisation analysis and enzymatic assays (see for example Pearson and Lipman (1988), PNAS USA, 85: 2444).

Thus, the RP-factors of the invention include the factors shown in FIGS. 1 A and 1B, together with their species variants, allelic forms, homologues, derivatives, muteins and corresponding secreted/nonsecreted forms (vide infra). Preferably, the RP-factors of the invention are species variants, allelic forms, homologues, derivatives, muteins and 5 corresponding secreted/nonsecreted forms of any one of the proteins represented in FIG. 1A and FIG. 1B.

The RP-factors may be synthesised in the form of a precursor which is processed to produce a mature form. Such processing may proceed via various intermediate (pro-) forms. Such precursors, intermediate forms and mature proteins are all intended to be covered by the term "RP-factor" as used herein, except where indicated otherwise. As used herein, the term "pro-RP-factor" specifically defines any of various precursors (which may or may not be active) of a mature RP-factor.

The processing may comprise proteolytic cleavage and/or secretion. The precursors may be inactive, and become active on processing as a mature form. The precursors may comprise proteins having secretory leader sequences which are removed during secretion (pre-forms). Such forms are herein referred to as "pre-RP-factor or pre-pro-RP-factors". As explained above, such pre- or prepro-forms are also intended to be covered by the 20 term "RP-factor" as used herein, except where indicated otherwise.

Processing may be attendant on the binding of an RP-factor precursor to a cognate receptor. Such receptors may then directly (or indirectly) cleave the precursor to produce a more mature form of the RP-factor. Such processing may occur as a cascade, involving several receptor-processing complexes,

and so ultimately result in the production of a mature RP-factor which then acts as a signalling moiety by binding to a terminal (signal transducing) receptor.

In such processing, the proximal (or intermediate) receptors may function as convertases, and the terminal receptor as a signal transducer. However, a receptor may function as both a convertase and a signal transducer. As used herein, the term "convertase" is intended to define a molecule which binds an RP-factor precursor and (directly or indirectly) processes it to produce a more mature form. They may, for example, have protease activity.

The receptors/convertases discussed above may be disposed at the cell surface (e.g. membrane bound), cytosolic or extracellular.

Preferably, the RP-factor is derived from a bacterium (e.g. a pathogenic bacterium). Particularly preferred are RP-factors derived from high G+C Gram-positive bacteria.

The term "derived from" as applied to a defined source is intended to define not only a source in the sense of it being the physical origin for the material, but also to define material which has structural and/or functional characteristics which correspond to those of material which does originate from the reference source. Thus, a protein "derived from" a given source need not necessarily have been purified from that source.

The term "high G+C Gram-positive bacterium" is a term of art defining a particular class of evolutionarily related bacteria. The class includes *Micrococcus* spp. (e.g. *M. luteus*), *Mycobacterium* spp. (for example a fast- or slow-growing mycobacterium, e.g. *M. tuberculosis*, *M. leprae*, *M. smegmatis* or *M. bovis*), *Streptomyces* spp. (e.g. *S. rimosus* and *S. coelicolor*) and *Corynebacterium* spp. (e.g. *C. glutamicum*). Preferred according to the invention are RP-factors/cognate receptors/convertases derived from mycobacteria ("mycobacterial RP-factors/RP-factor receptors/convertases").

The invention also contemplates homologues, allelic forms, species variants, derivatives, muteins or equivalents of the RP-factors and RP-factor receptors/convertases of the invention.

The term "homologue" is used herein in two distinct senses. It is used sensu stricto to define the corresponding protein from a different organism (i.e. a species variant), in which case there is a direct evolutionary relationship between the protein and its homologue. This may be reflected in a structural and functional equivalence, the protein and its homologue performing the same role in each organism.

The term is also used herein sensu lato to define a protein which is structurally similar (i.e. not necessarily related and/or structurally and functionally equivalent) to a given (reference) RP-factor. In this sense, homology is recognised on the basis of purely structural criteria by the presence of amino acid sequence identities and/or conservative amino acid changes (as set out by Dayhoff et alia, *Atlas of protein structure* vol. 5, National BioMed Fd'n, Washington D.C., 1979).

For the purposes of the invention, homologues may be recognised as those proteins the corresponding DNAs of which are capable of specifically or selectively cross-hybridising, or which can cross-hybridise under selective, appropriate and/or appropriately stringent hybridisation conditions.

The term "selectively or specifically (cross)hybridisable" in this context indicates that the sequences of the corresponding ssDNAs are such that binding to a unique (or small class) of homologous sequences can be obtained under more or less stringent hybridisation conditions. This method of the invention is not dependent on any particular hybridisation condi-

tions, which can readily be determined by the skilled worker (e.g. by routine trial and error or on the basis of thermodynamic considerations).

Preferably, the homologues, derivatives, muteins or equivalents of the RP-factor of the invention have at least 20% identity with any one of the particular amino acid sequences shown in FIG. 1A or FIG. 1B.

Particularly preferred are homologues, derivatives, muteins or equivalents of the RP-factor of the invention which have at least 30% identity, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% identity with any one of the particular amino acid sequences shown in FIG. 1A or FIG. 1B.

The homologues, derivatives, muteins or equivalents of the RP-factor of the invention may have at least 25% homology with any one of the particular amino acid sequences shown in FIG. 1A or FIG. 1B.

Particularly preferred are homologues, derivatives, muteins or equivalents of the RP-factor of the invention which have at least 30% homology, for example at least 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 98% homology with any one of the particular amino acid sequences shown in FIG. 1A or FIG. 1B.

The term "derivative" as applied herein to the proteins (e.g. the RP-factors or RP-factor receptors/convertases) of the invention is used to define proteins which are modified versions of the proteins of the invention. Such derivatives may include fusion proteins, in which the proteins of the invention have been fused to one or more different proteins or peptides (for example an antibody or a protein domain conferring a biochemical activity, to act as a label, or to facilitate purification).

The derivatives may also be products of synthetic processes which use a protein of the invention as a starting material or reactant.

The term "mutein" is used herein to define proteins that are mutant forms of the proteins of the invention, i.e. proteins in which one or more amino acids have been added, deleted or substituted. The muteins of the invention therefore include fragments, truncates and fusion proteins (e.g. comprising fused immunoglobulin, receptor, convertase or enzyme moieties).

The muteins of the invention also include proteins in which mutations have been introduced which effectively promote or impair one or more activities of the protein, for example mutations which promote or impair the function of a receptor, a recognition sequence or an effector binding site.

Muteins may be produced by any convenient method. Conveniently, site-directed mutagenesis with mutagenic oligonucleotides may be employed using a double stranded template (pBluescript KS II construct containing the RP-factor or RP-factor receptor/convertase gene), (e.g. Chameleon™ or QuikChange™—Stratagene™). After verifying each mutant derivative by sequencing, the mutated gene is excised and inserted into a suitable vector so that the modified protein can be over-expressed and purified.

Preferred mutant forms are truncates consisting (or consisting essentially) of the RP-factor signalling domain or the RP-factor specificity-determining factor, or of the ligand binding domain of the RP-factor receptor, or combinations of two or more of the foregoing.

The invention also contemplates chimaeric RP-factors. These are factors which comprise one or more heterologous domains. In this context, a heterologous domain is a portion of an RP-factor which is derived from a different RP-factor to that from which the other domain(s) with which it is associated are derived. Such chimaeric RP-factors find particular

utility in applications where the specificity and/or activity of the RP-factor is manipulated or altered.

Useful in the construction of such chimaeric RP-factors are DNA fragments or cassettes consisting essentially of DNA encoding selected domains (for example, the signalling domain or the specificity-determining domain), the fragment or cassette optionally being bounded by one or more restriction endonuclease cleavage sites or cloning sites. The invention also contemplates concatenated domain cassettes, as well as mutant RP-factor structural genes which have cloning sites (e.g. one or more restriction endonuclease cleavage sites) located in one or more interdomain regions.

The term equivalent as used herein and applied to the materials of the invention defines materials (e.g. proteins, DNA etc.) which exhibit substantially the same functions as those of the materials of the invention while differing in structure (e.g. nucleotide or amino acid sequence). Such equivalents may be generated for example by identifying sequences of functional importance (e.g. by identifying conserved or canonical sequences or by mutagenesis followed by functional assay), selecting an amino acid sequence on that basis and then synthesising a peptide based on the selected amino acid sequence. Such synthesis can be achieved by any of many different methods known in the art, including solid phase peptide synthesis (to generate synthetic peptides) and the assembly (and subsequent cloning) of oligonucleotides.

The homologues, fragments, muteins, equivalents or derivatives of the proteins of the invention may also be defined inter alia as those proteins which cross-react with antibodies to the proteins of the invention, and in particular which cross-react with antibodies directed against any of the specific proteins shown FIG. 1A or FIG. 1B.

The invention also contemplates all individual functional domains of the RP-factors of the invention as separate and independent entities.

The invention also contemplates recombinant RP-factor. As used herein, the term "recombinant" is intended to define material which has been produced by that body of techniques collectively known as "recombinant DNA technology" (for example, using the nucleic acid, vectors and or host cells described infra).

The invention also contemplates a pharmaceutical composition (e.g. a vaccine) comprising the RP-factor or RP-factor receptor/convertase (or homologue, species variant, allelic form, derivative, mutein or equivalent thereof) of the invention.

A pharmaceutical composition is a solid or liquid composition in a form, concentration and level of purity suitable for administration to a patient (e.g. a human or animal patient) upon which administration it can elicit the desired physiological changes. The vaccines of the invention may include any suitable adjuvant (e.g. Freund's adjuvant, BCG or BCG extracts).

In another aspect, the invention relates to a pharmaceutical composition comprising the material of the invention which is: (a) for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or (b) in a pharmaceutical excipient, a unit dosage form or in a form suitable for local or systemic administration.

In another aspect, the invention relates to an antibody (or antibody derivative) specific for the RP-factor (or homologue, derivative, mutein or equivalent thereof) of the invention.

The antibody is preferably in a form suitable for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or formulated in a pharmaceutical excipient, a unit dosage form or in a form suitable for local or systemic administration.

tion. The antibody may be labelled and/or immortalised and/or conjugated to another moiety, and such embodiments find particular utility in diagnostic applications.

According to another aspect of the invention there is provided an isolated or recombinant RP-factor receptor.

The receptor/convertase may be derived from any of the sources hereinbefore described, for example from a bacterial source (e.g. a pathogenic bacterial source). Such sources include high G+C Gram-positives, *Micrococcus* spp. (e.g. *M. luteus*); or *Mycobacterium* spp. (for example a fast- or slow-growing mycobacterium, e.g. *M. tuberculosis*, *M. leprae*, *M. smegmatis* or *M. bovis*); or *Streptomyces* spp. (e.g. *S. rimosus* and *S. coelicolor*); or *Corynebacterium* spp. (e.g. *C. glutamicum*).

The invention also contemplates homologues, derivatives, muteins or equivalents of the receptors/convertases of the invention, as well as recombinant RP-factor receptors/convertases (as hereinbefore defined).

The invention also contemplates a pharmaceutical composition (e.g. a vaccine) comprising the receptor/convertase (or homologue, derivative, mutein or equivalent thereof) of the invention.

Preferably, the receptor/convertase (or homologue, derivative, mutein or equivalent thereof) or pharmaceutical composition is: (a) for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or (b) in a pharmaceutical excipient, a unit dosage form or in a form suitable for local or systemic administration.

Also contemplated is an antibody (or antibody derivative) specific for the receptor/convertase (or homologue, derivative, mutein or equivalent thereof) of the invention. The antibody may be: (a) for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or (b) in a pharmaceutical excipient, a unit dosage form or in a form suitable for local or systemic administration.

Also contemplated is an RP-factor antagonist or inhibitor.

Preferably, the antagonist or inhibitor comprises: (a) the antibody of the invention; and/or (b) the receptor/convertase of the invention; and/or (c) an RP-factor mutein comprising an RP-factor specificity-determining domain, which for example lacks a functional signalling domain. The receptor may function as an antagonist or inhibitor if administered in soluble form, where it may act as a sink for soluble RP-factor. Preferably, modified receptors consisting of the receptor domain (and lacking the membrane anchor and wall spanning domain) are used as inhibitors or antagonists. Such derivatives may exhibit higher solubility.

The antagonist or inhibitor of the invention is preferably: (a) for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or (b) in a pharmaceutical excipient, a unit dosage form or in a form suitable for local or systemic administration.

Also contemplated by the invention is an RP-factor agonist, activator or mimetic. Preferably, the agonist, activator or mimetic comprises: (a) the RP-factor receptor/convertase antibody as herein described; and/or (b) an RP-factor mutein comprising (or consisting of) an RP-factor specificity-determining domain; and/or (c) an RP-factor mutein comprising (or consisting of) an RP-factor signalling domain; and/or (d) operably coupled combinations of any of (a)-(c).

The agonist, activator or mimetic may be: (a) for use in therapy (e.g. immunotherapy), diagnosis or prophylaxis; and/or (b) formulated in a pharmaceutical excipient, a unit dosage form, in a form suitable for local or systemic administration or in admixture with an antibiotic.

Preferably, the agonist, activator or mimetic may be for use in adjunctive therapy (for example formulated or presented in combination with an antimicrobial agent, e.g. an antibiotic).

The invention also contemplates isolated nucleic acid encoding the RP-factor (or homologue, derivative, allelic form, species variant, mutein or equivalent thereof) or RP-factor receptor/convertase (or homologue, derivative, allelic form, species variant, mutein or equivalent thereof) of the invention. The nucleic acids of the invention therefore embrace DNA having any sequence so long as it encodes the proteins of the invention. It will be appreciated by those skilled in the art that as a result of degeneracy in the genetic code, any particular amino acid sequence of the invention may be encoded by many different sequences. Thus, the nucleic acid sequence may be selected or optimised, e.g. with respect to the codon usage in any particular host cell.

The invention also contemplates vectors (e.g. an expression vector) comprising the nucleic acid of the invention. The nature of the vector is not critical to the invention. Any suitable vector may be used, including plasmid, virus, bacteriophage, transposon, minichromosome, liposome or mechanical carrier.

The expression vectors of the invention are DNA constructs suitable for expressing DNA which encodes the desired protein product (e.g. RP-factor or RP-factor receptor) which may include: (a) a regulatory element (e.g. a promoter, operator, activator, repressor and/or enhancer), (b) a structural or coding sequence which is transcribed into mRNA and (c) appropriate transcription, translation, initiation and termination sequences. They may also contain sequence encoding any of various tags (e.g. to facilitate subsequent purification of the expressed protein, such as affinity (e.g. His) tags).

Particularly preferred are vectors which comprise an expression element or elements operably linked to the DNA of the invention to provide for expression thereof at suitable levels. Any of a wide variety of expression elements may be used, and the expression element or elements may for example be selected from promoters, enhancers, ribosome binding sites, operators and activating sequences. Such expression elements may comprise an enhancer, and for example may be regulatable, for example being inducible (via the addition of an inducer).

As used herein, the term "operably linked" refers to a condition in which portions of a linear DNA sequence are capable of influencing the activity of other portions of the same linear DNA sequence. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is operably linked to a coding sequence if it controls the transcription of the sequence; a ribosome binding site is operably linked to a coding sequence if it is positioned so as to permit translation.

The vector may further comprise a positive selectable marker and/or a negative selectable marker. The use of a positive selectable marker facilitates the selection and/or identification of cells containing the vector.

Also contemplated by the invention are host cells comprising the vector of the invention. Any suitable host cell may be used, including prokaryotic host cells (such as *Escherichia coli*, *Streptomyces* spp. and *Bacillus subtilis*) and eukaryotic host cells.

In another aspect, the invention provides a culture or transport medium comprising the RP-factor (or homologue, derivative, mutein or equivalent thereof) of the invention. The culture medium may take any convenient form, such as for example agar plates, broths, slopes, coated dipsticks, coated probes, membranes, coated or filled wells or films. The

medium may be a defined or complex medium, and may contain indicator dyes to facilitate identification of cultured microorganisms. Preferably, the medium is suitable for the culturing or transport of bacteria, for example *Mycobacterium* spp. *Streptomyces* spp. and *Corynebacterium* spp.

The invention also contemplates a nucleic acid probe comprising nucleic acid complementary to the nucleic acids of the invention. Such probes are preferably selectively hybridisable with nucleic acid encoding the proteins (e.g. the RP-factors of RP-factor receptors/convertases) of the invention. They are conveniently single stranded DNA or RNA probes.

The invention also contemplates a diagnostic kit comprising the factor (or homologue, derivative, mutein or equivalent thereof), receptor, antibody, probe or culture medium of the invention.

In another aspect, the invention contemplates antisense DNA corresponding to the nucleic acid encoding the RP-factor or RP-factor receptor/convertase of the invention.

The invention also contemplates a process for producing an antimicrobial drug comprising the steps of: (a) providing an RP-factor receptor; (b) providing candidate drugs; (c) screening the candidate drugs by contacting the RP-factor receptor/convertase with one of the candidate drugs and determining the affinity of the candidate drug for the RP-factor receptor, wherein the affinity is an index of antimicrobial activity, and optionally (d) synthesising or purifying a drug having antimicrobial activity on the basis of the identity of the candidate drug screened in step (c).

Preferably, the process for producing an antimicrobial drug comprises the steps of: (a) providing an RP-factor receptor/convertase; (b) providing a candidate drug; (c) providing an RP-factor; (d) screening the candidate drugs by contacting the RP-factor receptor/convertase with one of the candidate drugs in the presence of the RP-factor, and then determining the ability of the candidate drug to compete non-productively with the RP-factor for binding to the RP-factor receptor, wherein the competitive binding ability is an index of antimicrobial activity, and optionally (e) synthesising or purifying a drug having antimicrobial activity on the basis of the identity of the candidate drug screened in step (d).

The invention also covers an antimicrobial drug produced by (or obtainable by) the processes of the invention, and also derivatives thereof.

Also contemplated by the invention is a method for determining the microbiological quality of a product (e.g. a food-stuff, pharmaceutical preparation or medical product) comprising the step of contacting a sample of the product with an RP-factor (for example, an RP-factor as hereinbefore defined). In such methods, the RP-factor preferably forms part of a nutrient composition (e.g. a plate, broth, film or dipstick).

In another aspect, the invention relates to a method of culturing bacterial (e.g. mycobacterial) cells, comprising the step of incubating the cells in a culture medium containing an RP-factor (for example, an RP-factor as hereinbefore defined).

Also contemplated by the invention is an ex vivo method of diagnosis, comprising the step of contacting a biological sample with an RP-factor (for example, an RP-factor as hereinbefore defined).

The diagnostic method of the invention preferably includes the step of incubating the culture or transport medium of the invention to permit growth of cells in the biological sample (e.g. bacterial cells).

Also contemplated by the invention is a method of: (a) stimulating the growth of a microorganism; and/or (b) resuscitating a dormant, moribund or latent microorganism; com-

prising the step of contacting the microorganism with an RP-factor (for example, an RP-factor as hereinbefore defined).

The invention also contemplates a process for producing the recombinant RP-factor or RP-factor receptor/convertase of the invention comprising the steps of: (a) culturing the host cell of the invention, and (b) purifying the factor or receptor/convertase from the cultured host cells (e.g. from a culture supernatant or cell fraction).

Also contemplated by the invention is a process for producing the recombinant RP-factor or receptor/convertase of the invention comprising the steps of: (a) probing a gene library with a nucleic acid probe which is selectively hybridisable with the cognate structural gene to produce a signal which identifies a gene that selectively hybridises to the probe; (b) expressing the gene identified in step (a) (for example by cloning into a host cell, e.g. according to the process as hereinbefore defined) to produce the factor or receptor.

Also covered is a recombinant RP-factor or receptor/convertase obtainable by the above-described process.

Medical Applications

The invention permits the isolation, synthesis and rational design of a wide range of novel medicaments and pharmaceuticals for use in therapy, prophylaxis and diagnosis.

The various forms of therapy, prophylaxis and diagnosis in which the materials of the invention find application may involve changing, breaking or perturbing the resuscitation (RP-factor) signal transduction pathway of one or more infecting pathogens.

Thus, the materials of the invention find general application as antimicrobial agents, for example as antibacterial agents. They may therefore be used in the treatment, prophylaxis or diagnosis of microbial (e.g. bacterial) infections, particularly those infections associated with latency (e.g. mycobacterial infections).

Thus, the invention may for example be used to prevent, reduce or interfere with: (a) the resuscitation of a latent (or dormant) pathogen, and/or (b) the growth of a pathogen, and/or (c) the multiplication and spread of a pathogen; and/or (d) the activation of a latent infection (for example a latent bacterial (e.g. mycobacterial) infection).

In general, the materials of the invention may be used to treat conditions in which changing, breaking or perturbing the resuscitation (RP-factor) signal transduction pathway or blockading the RP-factor receptor/convertase associated with an infecting pathogen is indicated.

Particularly useful materials for use in such therapies/prophylactic methods include RP-factor antagonists or inhibitors. Such antagonists or inhibitors may comprise antibodies to the RP-factor or to the RP-factor receptor/convertase as herein defined; the RP-factor receptor/convertase of the invention; an RP-factor mutein, e.g. which comprises an altered RP-factor specificity-determining domain and/or which lacks a functional signalling domain.

RP-factor antibodies act to sequester and ultimately eliminate endogenous RP-factors in a patient bearing a latent microbial infection.

RP-factor receptor antibodies bind non-productively to the receptors associated with the infecting pathogen. Antibodies to the convertase inactivate (e.g. by steric inhibition) the convertase activity and so prevent maturation of the RP-factor. The antibodies may therefore competitively inhibit the binding of endogenous RP-factor to the receptors/convertases associated with the infecting pathogen. Alternatively, they

may bind with high affinity (and/or essentially irreversibly) to the RP-factor receptors/convertases and so block RP-factor-ligand binding or RP-factor maturation. A similar activity is displayed by the RP-factor muteins having altered specificity and/or signalling activity.

In either case, the RP-factor-RP-receptor/convertase binding required for resuscitation of latent pathogens, growth of the pathogen and/or progression of the disease state is perturbed, reduced or abolished.

RP-factor receptors for use as therapeutics in such methods are uncoupled from the signal transduction pathway with which they are normally associated. Thus, they are preferably free (i.e. in soluble or dispersible) form and/or not membrane bound. In this way, effective circulating or systemic concentrations of the free RP-factor receptor can be established and maintained in a patient. In this form, the RP-factor receptors act as RP-factor sinks, and titrate out (and preferably ultimately eliminate) endogenous RP-factors in a patient bearing a latent microbial infection. The receptors therefore reduce or prevent activation of the (latent) pathogen and/or stimulation of pathogen growth, so slowing or halting the progression of the infection.

In another aspect, the invention may be used to resuscitate or assist in resuscitating (or activate or assist in activating) a latent (dormant) pathogenic microbe in vivo thereby to potentiate adjunctive antimicrobial therapy. The adjunctive antimicrobial therapies for use in such applications are those which depend for full efficacy on a non-latent or active (e.g. growing or replicating) target pathogen population (for example adjunctive therapies based on certain types of antibiotic). Thus, the materials of the invention may act synergistically with various antimicrobial compounds in antimicrobial therapy.

In a preferred embodiment, the invention is used to potentiate the antimicrobial therapy of tuberculosis, for example involving co-administration of one or more of isoniazid, rifampicin, pyrazinamide and/or ethambutol (or streptomycin).

Particularly useful materials for use in such therapies include for example the RP-factors of the invention, RP-factor agonists, activators and mimetics. Such agonists, activators or mimetics may comprise: the RP-factor receptor antibodies as hereinbefore described; the RP-factor convertase as hereinbefore defined; an RP-factor mutein comprising (or consisting of) an RP-factor specificity-determining domain; an RP-factor mutein comprising (or consisting of) an RP-factor signalling domain; and/or operably coupled combinations thereof.

The RP-factor receptor antibodies for use in such methods are those which serve to trigger an efferent signal transduction pathway at the RP-factor receptor. They may therefore act as RP-factor mimetics, breaking latency/dormancy and acting to resuscitate the pathogen.

Particularly useful in such methods are mutant RP-factors having altered specificity (e.g. in which the specificity-determining domain has been mutated or modified). Such mutant RP-factors may be active against a broad range of pathogens (e.g. against substantially all pathogenic or infective mycobacteria) or targeted against specific pathogens (for example, *M. tuberculosis* and *M. leprae*).

The antibodies, RP-factors, receptors and convertases discussed above may be administered directly or via a live vaccine vehicle. Such live vaccine vehicles comprise microorganisms which have been genetically engineered to express (and preferably secrete) the therapeutically active antibodies, RP-factors, receptors and convertases of the invention in vivo.

The invention therefore finds application in the treatment of a wide variety of microbial infections, and finds particular application in the treatment of latent microbial (e.g. bacterial) infections.

In preferred embodiments, the invention finds application in the treatment of actinomycete or mycobacterial infections, for example those involving *M. tuberculosis*, *M. leprae*, *M. bovis*, *M. kansasii* and *M. avium*.

Other infections which may be treated according to the invention include those involving *Corynebacterium* spp. (including *Corynebacterium diphtheriae*), *Tropheryma whippelii*, *Nocardia* spp. (including *Nocardia asteroides* and *Nocardia brasiliensis*), *Streptomyces* spp. (including *Streptomyces griseus*, *Streptomyces paraguayensis* and *Streptomyces somaliensis*), *Actinomyces* spp., *Nocardiopsis* spp., *Rhodococcus* spp., *Gordonia* spp., *Tsukamurella* spp. and *Oerskovia* spp. as well as other pathogenic organisms from the group referred to as high G+C Gram-positive bacteria. Other infections which may be treated include those involving pathogenic low G+C Gram-positive bacteria (e.g. *Streptococcus* spp., *Staphylococcus* spp., *Listeria* spp., *Bacillus* spp., *Clostridium* spp. and *Lactobacillus* spp.).

The invention may also be embodied in various vaccines or immunotherapeutic agents.

Such vaccines or agents target one or more elements of the RP-factor mediated signal transduction pathway described herein (and in particular, the RP-factor or RP-factor receptors/convertases themselves). Thus, the RP-factors may be administered as part of a vaccine or immunotherapeutic composition to elicit an immune response directed against endogenous RP-factor in the patient, so reducing, preventing activation of the pathogen and so slowing or halting the progression of the infection.

Alternatively (or in addition), the RP-factor receptors/convertases may be administered as part of a vaccine or immunotherapeutic composition to elicit an immune response directed against receptors for pathogen-borne RP-factor in the patient. In this way, cellular and/or humoral immune responses may be stimulated against the pathogen(s) and/or activation of a latent pathogen (or its continued growth or multiplication) via the RP-factor signal transduction pathway may be reduced or prevented, so slowing or halting the progression of the infection.

The invention also finds application in the preparation of live vaccines: attenuated microbial strains can be constructed in which the gene(s) encoding (or regulating the expression or activity of) one or more RP-factors are mutated. Such attenuated vaccines may be based on mutant strains of actinomycetes, mycobacteria (for example *M. tuberculosis*, *M. leprae*, *M. bovis* (such as *M. bovis* BCG), *M. kansasii* and *M. avium*), *Corynebacterium* spp. (including *Corynebacterium diphtheriae*), *Tropheryma whippelii*, *Nocardia* spp. (including *Nocardia asteroides* and *Nocardia brasiliensis*), *Streptomyces* spp. (including *Streptomyces griseus*, *Streptomyces paraguayensis* and *Streptomyces somaliensis*), *Actinomyces* spp., *Nocardiopsis* spp., *Rhodococcus* spp., *Gordonia* spp., *Tsukamurella* spp. and *Oerskovia* spp. as well as other pathogenic organisms from the group referred to as high G+C Gram-positive bacteria.

Particularly useful in such attenuated vaccines are strains bearing mutated RP-factor-encoding genes. Such mutations may be frameshift, deletion, insertion and/or substitution mutations. In preferred embodiments the mutations are null mutations (e.g. non-reverting null mutations), and may prevent growth of the microbe (i.e. "attenuate" it). In other embodiments the mutations may result in the expression of mutant RP-factors having altered specificity (e.g. in which the

specificity-determining domain has been mutated or modified) and/or which lack a functional signalling domain. Such mutant RP-factors may bind with high affinity (and/or essentially irreversibly) and non-productively to the RP-factor receptors/convertases and so block RP-factor-ligand binding or RP-factor maturation. The attenuated microbial strains of the invention may also bear mutations in other genes (for example, in other genes essential to growth), and may also bear one or more genetic marker elements.

Biotechnological Applications

It is widely recognised that the great majority (probably well in excess of 99%) of soil organisms have not yet been cultured. Hitherto uncultured organisms are also expected to exist in other sources. The present invention may be used to permit the recovery of such organisms by culture from any source. Thus, the invention provides a way of unlocking an immense reservoir of biodiversity that is known to exist, but is presently inaccessible.

Thus, the present invention provides an unprecedented resource from which libraries of potentially useful microorganisms and biomolecules can be generated. Such libraries can then be used in screening methods to search for medically or industrially useful products.

Thus, in another aspect the invention provides a process for producing a library of biomolecules comprising the steps of: (a) providing a sample (e.g. a soil, marine, food, freshwater, tissue or organism-derived); (b) incubating the sample in a culture medium comprising an RP-factor (for example, an RP-factor as defined in the preceding claims or a culture supernatant comprising an RP-factor) to produce a microbial culture; and (c) isolating microorganisms from the culture of step (b).

The process may further comprise the step of screening the isolated microorganisms for those which elaborate one or more biomolecules of interest (for example a metabolite, enzyme, antibiotic (e.g. antiviral, antibacterial or antifungal agent) or toxin).

Also contemplated is a biomolecule produced by (or obtainable by) the above process, or a derivative thereof.

In another aspect, the invention provides a process for producing a library of microorganisms (e.g. bacteria) comprising the steps of: (a) providing a sample (e.g. a soil, marine, food, freshwater, tissue or organism-derived sample); (b) incubating the sample in a culture medium comprising an RP-factor (for example, an RP-factor as defined in the preceding claims or a culture supernatant comprising an RP-factor) to produce a microbial culture; (c) isolating microorganisms from the culture of step (b).

Also contemplated is a microbe produced by (or obtainable by) the above process, or a 30 derivative (e.g. mutant) thereof.

EXEMPLIFICATION

The invention will now be described in more detail with reference to several Examples. These are for exemplary purposes only and are not intended to limit the invention in any way.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1E: Part A. Multiple sequence alignment of the predicted amino-acid sequences of RP-factor-like gene products from *M. luteus*, *M. tuberculosis*, *M. leprae* and *Streptomyces coelicolor*. Proteins similar to the RP-factor are derived from *M. tuberculosis* (accession nos. U38939, nt

2406-2765, and Z81368, nt 33932-34396) and *M. leprae* (accession nos. L01095, nt 12292-12759, and L04666, nt 25446-24921). The DNA sequences of interest in accession Z81 368 are also encompassed by accession AD000010.

5 N-terminal residues corresponding to predicted Gram-positive signal sequences are underlined. The *M. leprae* L04666 sequence may also contain a short, 32 aa signal peptide.

Part B. Multiple sequence alignment of gene products related to YabE of *Bacillus subtilis*. The alignment is given in two parts (A and B), with aligned residues in upper case. Those residues which are conserved (or conservatively substituted) in two or more sequences are in bold. In Part A, perfectly conserved residues are marked with a hash (#) and conservative substitutions with a dot (.). Cperfring is an incomplete ORF 1 from *Clostridium perfringens* (Acc. No. U04966); Caceto506 is an incomplete ORF from contig 506, *Clostridium acetobutylicum* genome sequencing project. YocH from *B. subtilis* and YabE from *B. subtilis* are YocH and YabE predicted gene products from the *B. subtilis* genome sequencing project (Acc. Nos. BG13521 and P37456).

Part C. Alignment of the RP-factor C-terminal domain with known and hypothetical wall-associated proteins from other organisms. Perfectly conserved residues are marked with an asterisk, those conserved in at least 7 sequences are marked with a dot (.).

Part D. Motifs in the C-terminus (residues 158-322) of MtubMTVO43.

Part E. Alignment between the predicted amino acid sequence of the *M. luteus* RP-factor and p60 proteins from *Listeria* spp. Many of the residues that are conserved in the alignment between the C-terminal portion of the *M. luteus* RP-factor (residues 125-220) and the *L. monocytogenes* EGD p60 protein (residues 158-245), are also conserved in the p60 protein from six other *Listeria* spp.

FIGS. 2A- and 2B: Part A. The sequence of the RP-factor-encoding gene and its predicted product. The nucleotide sequence is in lower case with PCR primers in bold. The predicted protein sequence is in upper case bold (single letter code). Protein and peptide microsequence data used for oligonucleotide design are in upper case italics.

Part B. The sequence of a 299 base pair DNA fragment encoding part of an RP-factor from *Streptomyces coelicolor*. The deduced amino acid sequence is given below the DNA sequence using the single letter amino acid code.

FIGS. 3A-3D: The elution profile of the resuscitation activity. Fractions eluted from the DEAE Sepharose™ column (see Materials and Methods) with 0.25 M KCl were applied to a Mono Q® column which was developed with a 20 ml linear gradient from 0.08 to 0.28 M KCl in 10 mM Tris-Cl buffer supplemented by 10% glycerol, pH 7.4. 10 ml of a diluted suspension of starved cells (CFU $3 \cdot 10^6$ cells·ml⁻¹, total count $1.2 \cdot 10^9$ cells·ml⁻¹) were added to 200 ml of LMM supplemented with 0.5% w/v lactate and 0.05% yeast extract containing of 2 µl of each fraction in 5-10 replicates in the Bioscreen instrument. For details see Materials and Methods.

A: absorbance at 280 nm and magnitude of KCl concentration. B: resuscitation activity. C: SDS-PAGE profile of the fractions following DEAE-cellulose and Mono Q® chromatography. Lanes 1, markers (94,000, 67,000, 43,000, 30,000, 20, 100, 14,400); 2, fraction from DEAE-cellulose column; 3, purified preparation (fraction number 8 from the Mono Q®-column). D: Reduction of apparent lag phase of viable cells. 10 µl of a diluted suspension of viable, stationary phase cells (viable count 20 cells) was added to 200 ml of LMM supplemented with 0.5% w/v L-lactate and containing 2 µl of each fraction (from a different experiment to that shown in parts A and B) in 5-10 replicates in the Bioscreen instrument.

The apparent lag phase was estimated by extrapolating the exponential growth line to the abscissa.

FIGS. 4A-4B: Effect of purified RP-factor on *M. luteus*.

A. Concentration dependence of RP-factor activity for resuscitation: resuscitation of dormant cells with different concentrations of RP-factor. 10 μ l of a diluted suspension of starved cells (CFU $3 \cdot 10^6$ cells \cdot ml $^{-1}$, total count $5 \cdot 10^9$ cells \cdot ml $^{-1}$) was added to 200 μ l of LMM supplemented with 0.5% w/v L-lactate, 0.05% yeast extract and RP-factor in concentrations shown in 5-10 replicates in the Bioscreen instrument. For details see Materials and Methods.

B. Growth of washed cells. Stationary phase cells of *M. luteus* grown in LMM were washed five times by suspension and centrifugation in LMM from which lactate had been omitted. Bacteria were finally suspended in the same medium by repeatedly passing them through a syringe, diluted and inoculated into a 20 ml flask containing LMM or LMM plus 31 pM RP-factor. The initial cell density was 250 viable cells per ml and incubation was at 30° C. with intensive shaking. Growth was monitored by plating 0.1 ml samples on plates containing broth E solidified with agar.

FIGS. 5A-5C: Detection of RP-factor-like genes in *Micrococcus luteus*, *Mycobacterium smegmatis* and *Streptomyces rimosus*.

Part A	Part B	
<i>M. luteus</i>	<i>M. luteus</i>	Part C
Lane 1	λ BstEII	λ PstI
Lane 2	ClaI	<i>S. rimosus</i> XhoI
Lane 3	SaII	<i>S. rimosus</i> StuI
Lane 4	SacII	<i>S. rimosus</i> SmaI
Lane 5	PstI	<i>S. rimosus</i> PvuII
Lane 6	NcoI	<i>S. rimosus</i> PstI
Lane 7	NheI	<i>S. rimosus</i> BamHI
Lane 8	MluI	<i>M. smegmatis</i> XhoI
Lane 9	AatII	<i>M. smegmatis</i> StuI
Lane 10	λ PstI	<i>M. smegmatis</i> SmaI
Lane 11		<i>M. smegmatis</i> PvuII
Lane 12		<i>M. smegmatis</i> PstI
Lane 13		<i>M. smegmatis</i> BamHI
Lane 14		λ PvuII

FIGS. 6A and B Effect of *M. luteus* RP-factor on the growth of *Mycobacterium smegmatis* (A) and *Mycobacterium bovis* (B) in batch culture as observed turbidimetrically. *M. smegmatis* was grown in broth E, to which was added RP-factor at 31 pMol/L. Cells were inoculated at a level of circa 200 per well, and growth was monitored in the Bioscreen instrument. *M. bovis* was grown in Sauton medium, as described in the Materials and Methods section, to which RP-factor (620 pMol/L) was either added or not. The inoculum was circa $1 \cdot 10^5$ cells \cdot ml $^{-1}$, and the OD shown is the average of 10 separate determinations of 10 separate tubes.

FIGS. 7A-7C: A: Purification of His-tagged RP-factor. RP-factor was expressed in *E. coli* HSM174(DE3) and purified as described infra. Shown is the SDS-PAGE profile of fractions following Ni $^{2+}$ -chelation chromatography. The molecular weight (kDa) markers (SIGMA) were bovine serum albumin (67), ovalbumin (43), glyceraldehyde 3-phosphate dehydrogenase (36), carbonic anhydrase (30), soya bean trypsin inhibitor (20.1), and lactalbumin (14.4). Lane: 1, markers; 2, crude extract from *E. coli* containing pET19b vector; 3, crude extract from *E. coli* containing pRPf1; 4, purified recombinant RP-factor.

B: Reduction of the apparent lag phase of viable cells of *M. luteus* by purified 5 recombinant RP-factor. For experimental details see the legend for FIG. 3C. A dilution factor of 10^0 corresponds to 33 ig RP-factor/ml.

C: Stimulation of the growth of washed cells of *M. luteus* by purified recombinant RP-factor. Stationary phase cells of *M. luteus* grown in LMM were washed 5 times by suspension and centrifugation in LMM from which lactate had been omitted. Bacteria were finally suspended in the same medium by repeatedly passing them through a syringe, diluted, and inoculated into a 20 ml flask with LMM or LMM in the presence of RP-factor (230 pMol/L). The initial cell density was ca. 10^2 viable cells per ml and incubation was at 30° C. with intensive shaking. Growth was monitored by plating 0.1 ml samples on plates containing nutrient broth E solidified with agar.

FIGS. 8A and 8B: A: Anti-RP-factor serum inhibits the growth of *Micrococcus luteus*. Bacteria were inoculated at an initial density of $5 \cdot 10^5$ per ml into lactate minimal medium (LMM) and the OD $_{600nm}$ was monitored at intervals. Growth of the cultures was monitored over 140 hours at intervals. The samples labelled LMM+Ab and LMM+control Ab contain equivalent amounts of immune and pre-immune serum, respectively. Immune serum (Ab) and pre-immune serum (control Ab) were employed at a 1:1000 dilution.

B: RP-factor overcomes the inhibitory effect of anti-RP-factor serum on growth of *Micrococcus luteus*. Bacteria were inoculated at an initial density of 10^7 cells per ml and growth was monitored by measuring the OD $_{600nm}$ at intervals. Immune serum (Ab) and pre-immune serum (control Ab) were employed at a 1:1000 dilution and RP-factor was added at a final concentration of 50 ng/ml.

FIGS. 9A and 9B: Part A. Blocked alignment of nine RP-factors (as explained infra, MtubZ94752 may be a cognate receptor). Areas of sequence identity/similarity are indicated by the shaded areas. The *S. coelicolor* gene product shown is a fragment. Part B. Schematic showing the domain structure of some gene products in the RP-factor family.

FIG. 10: Effect of recombinant RP-factor on growth of *M. tuberculosis* in Sauton medium. Sauton medium containing 0.05% Tween-80™ and 100 μ Mol/L Na oleate+10% (v/v) supplement (which contains, per liter, 50 g bovine serum albumin, 20 g glucose, 8.5 g NaCl) was inoculated to an initial cell density of $31 \cdot 10^3$ cfu/ml (viable count determined by plating on agar-solidified Middlebrook 7H9 medium containing 10% v/v supplement, composition as detailed above) [total count by microscopy= 10^6 cells per ml] with a 2.5 month-old culture of *M. tuberculosis* strain H37Ra grown in the same medium. Growth of tube cultures at 37° C. was measured by determining the OD $_{600nm}$ at intervals for 28 days. The undiluted concentrations of the RP-factors, Rpf (*M. luteus*) and Rpf2 (*M. tuberculosis*), employed for these 15 experiments were ca. 10 μ g/ml.

EXAMPLES

Material and Methods

Organisms and Media.

Micrococcus luteus NCIMB 13267 (previously described as "Fleming strain 2665") was grown aerobically at 30° C. in shake flasks in lactate minimal medium (LMM) containing L-lactate as described previously. When the culture had reached stationary phase agitation was continued at 30° C. for up to 2 months. Cultures were then held aerobically at room temperature without agitation for period for up to a further 2-3

months. The apparent initial viability of these cultures at this point (measured by comparing the plate count with the microscopic count) was less than 10^{-3} .

Mycobacterium smegmatis ("fast" strain, All-Russia State Institute for Control of Veterinary Preparations, Moscow) was grown in either Sauton medium or nutrient broth E (LabM). Overnight pre-cultures were used to inoculate cultures to an initial density of 10^3 cells/ml. *Mycobacterium bovis* (BCG), *Mycobacterium tuberculosis* H37RV and *Mycobacterium avium* were grown in Sauton medium.

M. luteus Spent Medium Preparation.

Supernatant was obtained after the centrifugation of late logarithmic phase *M. luteus* cultures (200-1000 ml) grown in lactate minimal medium or in the same medium in which lactate was replaced by succinate plus 0.01% yeast extract from which macromolecules had been removed by dialysis. The inoculum consisted of 2% of cells grown in rich medium (Broth E, LabM) and then washed in LMM lacking lactate. The supernatants were passed through a 0.22 μ m filter (Whatman) before use.

M. luteus Cell Viability by Plating.

Plates consisting of 1.3% Nutrient Broth E (LabM) or lactate minimal medium were used. Cell dilutions were made in quadruplicate with centrifuged and autoclaved spent 20 medium taken from the starved culture. Plates were incubated at 30° C. for 3-5 d.

M. luteus Cell Viability by MPN.

The MPN assay was performed in a Bioscreen C optical growth analyzer (Labsystems, Finland) using lactate minimal medium supplemented by 0.5% lactate and 0.05% of yeast extract as a resuscitation medium. Dilutions of starved cells were made as described. 10 μ l of each dilution (5-10 replicates) were added to a well containing 200 μ l of either lactate minimal medium supplemented by 0.5% lactate and 0.05% of yeast extract or the same medium with fraction tested (2-20 μ l). Growth (optical density) was monitored using a 600 nm filter. Plates were incubated at 30° C. with intensive continuous shaking. The overall measurement period was 120 h, each well being measured hourly.

The fractions obtained after chromatography were dialysed against elution buffer 2 (see below), diluted in resuscitation medium in various proportions (1:10, 1:100, 1:500, 1:1000, 1:5,000, 1:10,000) and filtered through 0.22 μ m Gelman filters before testing. The calculation of the MPN was based on published Tables.

Total Cell Counts

Unstained cells were counted with a phase-contrast microscope and an improved Neubauer counting chamber. In long-term experiments with mycobacteria, organisms 10 were stained with Ziehl-Neelsen reagent before counting.

Chromatography

Pre-wetted DEAE cellulose was added to culture supernatant (1:10 v/v) and incubated at 4° C. for 1 h with slow stirring. The cellulose was loaded into a column, and washed with 5 volumes of buffer 1 consisting of 10 mM Tris-Cl, 1 mM EDTA, 1 mM DTT, 10% (v/v) glycerol, pH 7.4 with 10 mM KCl. The column was eluted stepwise with 2-3 bed volumes of 0.3M KCl in buffer 1. The fraction obtained was slowly diluted with buffer 1 on ice to give a final KCl concentration of 0.08M. Forty column volumes of this fraction was then loaded onto a DEAE Sepharose™ fast flow column (1 part of Sepharose™ pre-equilibrated with buffer 1 containing 0.08M KCl). The column was washed with 5 bed volumes buffer 1 containing 0.08M KCl and eluted stepwise with 3 volumes of 0.25M KCl in buffer 1. The fraction obtained was again slowly diluted with buffer 1 on ice to a final KCl concentration of 0.08M, filtered through a 0.22 μ m Gelman filter and

loaded onto a Mono a® column (model HR5/5, pre-packed, Pharmacia) equilibrated with buffer 2 consisting of 10 mM Tris-Cl, 10% glycerol, pH 7.4 containing 0.08M KCl. The Mono Q® column was eluted by a linear gradient from 0.08 M to 0.28 M KCl in buffer 2 (the total volume of the elution was 20 ml). The flow rate and fraction size were 1 ml/min and 1 ml/tube respectively. All manipulations except the Mono Q® chromatography step were performed at 4° C. The fractions obtained were dialysed against 10 mM Tris-Cl containing 10% glycerol (dialysis is important for the retention of activity) and stored at 4° C. for up to 5 days without loss of activity. For prolonged storage in a deep freeze, fractions were dialysed in the same way and glycerol added to a final concentration of 20-30% w/v. The protein content in purified preparations was estimated by tryptophan fluorescence using lysozyme as a standard.

Trypsin Treatment:

Trypsin was added to the active, dialysed fraction obtained from the Mono Q® column and diluted by LMM supplemented with 0.5% w/v lactate and 0.05% yeast extract (1:100) (the final concentration of trypsin was 50 ug/ml). The mixture was incubated for 30 min at 37° C. The reaction was stopped by the addition of trypsin inhibitor (100 ug/ml). In control experiments trypsin inhibitor was added to the mixture (100 ug/ml) prior to incubation.

PAGE Electrophoresis.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was performed according to Laemmli. Chromatographic fractions were dialysed against 10 mM Tris HCl, pH 7.4 for 4-5 h, dried in a speed-vacuum apparatus (1.5 h), dissolved in sample buffer (Sigma, S-3401), loaded onto 15% acrylamide gel and run at a constant voltage of 200V. The gel was stained with colloidal Coomassie™ G (Sigma). Chemicals.

Nutrient Broth E, yeast extract and agar were obtained from Lab M, whilst L-lactate (Li salt), succinate, trypsin, soybean trypsin inhibitor and DEAE-Sepharose™ fast flow were obtained from Sigma. DEAE cellulose DE52 was obtained from Whatman, and Mono S® and Mono Q® from Pharmacia. Other chemicals were of analytical grade and were obtained from Sigma or BDH.

Protein microsequence data from the N-terminus (ATVDTWDRLEExSNGTxD) (SEQ ID NO: 38) and an internal peptide (VGEGYPHQASK) (SEQ ID NO: 42) obtained from the purified RP-factor were used to design two oligonucleotides, denoted A1 [GCSACSGTSGACAC-STGGGACCGSCTSGCSGAG] (SEQ ID NO: 37) and A2 [GCYTGRGTGIGGRTAICCYTCICC] (SEQ ID NO: 41), respectively. Taq polymerase was employed under standard conditions to amplify a 147 bp PCR product from *M. luteus* DNA with these primers. The PCR product obtained from *M. luteus* DNA with these two primers was labeled with digoxigenin and used as a probe for Southern hybridization experiments. SmaI-digested genomic DNA was size-fractionated by agarose gel electrophoresis and circa 1.4 kbp fragments were cloned in pMTL20 and established in *E. coli* strain DH5a. Two recombinant plasmids carrying the desired insert were detected by hybridization, confirmed by PCR using oligonucleotides A1 and A2, and one of them was manually sequenced on both strands using the dideoxy chain termination method.

Standard procedures were employed to isolate DNA from *M. luteus* and *M. smegmatis*. *Streptomyces rimosus* DNA was kindly supplied by Dr. D. Hranueli. Southern hybridisations with *M. smegmatis* and *S. rimosus* DNA were initially carried out under non-stringent conditions (0.5 SSC, 37° C.). Stringent conditions (0.1 SSC, 65° C.) were subsequently

employed for screening an ordered cosmid library of *Streptomyces coelicolor* A3(2) DNA.

Purification of RP-Factor

RP-factor purified from culture supernatants of cells grown in lactate minimal medium, according to the protocol described in Materials and Methods, revealed the presence of a significant amount of polymeric material eluted from all types of columns used, which inhibited both the resuscitation of dormant cells and the growth of viable cells of *M. luteus*. Moreover, elevated concentrations of this material could even cause the lysis of cells (not shown). This inhibitory material appears to be a polymer derived from lactate, as lactate-containing LMM stored for 10 hours at room temperature without cells and subjected to the same procedure of purification revealed inhibitory properties similar to those of this spent medium. To avoid this problem we replaced lactate in the growth medium with succinate, although for good growth it proved necessary to add a small amount (0.01% w/v) of yeast extract dialysed to remove macromolecules.

Using succinate-grown cultures, the active fraction was purified by a combination of anion exchange media (see Material and Methods). The final activity was eluted at around 180 mM KCl from a linear KCl gradient (from 0.08 to 0.28M KCl) on a Mono Q® column in 3 adjacent fractions (FIG. 3). It is worth mentioning that it proved important to dialyse the fractions before testing their activity because some fractions were inactive before dialysis. Active fractions did not change their resuscitation activity after dilution up to 400 times (v/v).

Interestingly, those fractions which were active in causing resuscitation could also increase the growth rate of viable cells.

The resuscitation-promoting material from the final purification step was checked by SDS-PAGE. The final product (FIG. 3C) proved to consist of a single protein with a molecular weight estimated to be ca 16 kD. All active fractions consist of single band with maximum content of protein in fraction N9.

Cloning of the RP-Factor Gene

Two primers were designed from protein microsequence data obtained for the N-terminus of the purified RP-factor and for an internal peptide. They were used to amplify a 147 bp fragment of *M. luteus* DNA, which was cloned and sequenced. The complete gene was then obtained by a combination of inverse PCR using oligonucleotides G1 and G2 and isolation of a 1.4 kbp SmaI restriction fragment. Sequencing revealed that the original PCR product was part of a gene capable of encoding a protein having a signal sequence (FIG. 2A). The predicted size of the secreted form of the gene product is 19,148 Dal, and its predicted N-terminal amino acid sequence agrees with the protein microsequence data, including residues that were not used in primer design (FIG. 2A). The fact that the predicted gene product is larger than the RP-factor purified from culture supernatants suggests that it may, for example, be secreted as a precursor which is converted to its biologically active form upon contact with its cognate receptor/convertase.

Identification of RP-Factor Homologues

A BLAST search was undertaken using the predicted amino acid sequence of the ORF from *M. luteus* as query. Seven genes with substantial similarity have been sequenced previously. Five are found in *M. tuberculosis* and two in *Mycobacterium leprae* (FIG. 1A). One or more gene products in each organism appear to have a secretory signal sequence (underlined in FIG. 1A). The functions of the predicted products of these mycobacterial genes are unknown; they were found by genome sequencing projects. The BLAST search

also revealed similarity between residues 126-220 of the RP-factor and a conserved segment of the (major extracellular) p60 proteins that have been implicated in adherence of *Listeria* spp. to 3T6 mouse fibroblasts suggesting, perhaps, a possible role for the RP-factor or a proteolytic product thereof in adhesion in *M. luteus* (FIG. 1E).

In common with *M. tuberculosis* and *M. leprae*, *M. luteus* contains a second gene similar to that encoding the RP-factor. Southern hybridisation experiments, using DNA samples cleaved with a range of different restriction enzymes, and the cloned 147 bp fragment as probe (FIGS. 5A & B), reveal two hybridising bands. The stronger hybridisation signal arises from the gene encoding the secreted RP-factor. The other gene may correspond to one of the other mycobacterial genes identified above.

Southern hybridisation experiments, using the 147 bp fragment as probe, as well as PCR experiments, using two oligonucleotides based on highly conserved amino acid motifs as primers, indicate that genes encoding proteins similar to the RP-factor are of widespread occurrence, at least throughout Gram-positive bacteria whose DNA has a high G+C content. Similar genes are detectable by either or both of these methods in all six *Streptomyces* species we have tested, including *Streptomyces rimosus* (FIG. 5C) as well as in other mycobacteria, including *Mycobacterium* I (four similar genes—FIG. 5C), *Mycobacterium bovis* (BCG) and *Corynebacterium glutamicum* (2 similar genes).

Domain Structure

The sequence information shows that the RP-factor gene and all of its mycobacterial homologues share a secretory signal sequence and a particularly highly conserved, ca. 70-residue segment. One (MTubZ94752) also has a membrane anchoring motif. The conserved 70-residue segment is a candidate for a signalling domain. Most of this segment is weakly hydrophilic (Kyte-Doolittle) and is predicted to form amphipathic α -helical (Garner-Robson; Chou-Fasman) or β -sheet regions (Eisenberg). Overall, the segment has a low surface probability (Emini). The C-terminal section, by contrast, is much less highly conserved and might be considered a better candidate for determining localization or specificity (i.e. be a cellular compartment-targeting or specificity-determining domain). By analogy with other protein signalling systems (e.g. many pro-hormones in animals, and systemin in plants) it is possible that the proximate signalling molecule is a proteolytically cleaved product.

Two acidic residues, D7 & E13 (numbering according to the *M. luteus* secreted protein), within this segment are absolutely conserved. The KAEQIKRAE segment (residues 51-59; SEQ ID NO: 64) represents an island of particularly high surface probability. These elements may form part of functional domains within the RP-factor protein.

The conserved domain contains four conserved tryptophan residues (one of which is in a region of high surface probability DTWDR-residues 4-8; SEQ ID NO: 65). In the complex between human growth hormone and its first bound receptor, interactions involving two surface-located tryptophan residues in the receptor account for more than 75% of the binding free energy of the complex (Clackson and Wells, Science 267, 383-386, 1995). The two conserved cysteine residues may form a disulphide bridge.

Alignments showing the domain structures of the various proteins are shown in FIGS. 9A 5 and 9B.

RP-Factor Activity

As well as resuscitating dormant cells, the purified RP-factor from *M. luteus* has been tested for growth-stimulatory activity against *M. luteus* and several other organisms. It strongly stimulates the growth of *M. luteus* and *M. smegmatis*

and it appears to have weaker activity on *M. tuberculosis*, *M. bovis* (BCG) and *M. avium* (see FIG. 6). In all cases, there is a shortening of the apparent lag phase in batch culture (see FIGS. 3D, 4B, 6B and Table 1). The factor is active in poor media and in poor media supplemented with yeast extract and it loses activity after boiling or treatment with trypsin.

When ca. 40 pMol/L RP-factor was added to washed cells of *Mycobacterium smegmatis*, growth occurred after 20-24 hr, whereas the control lacking RP-factor showed no growth after 6 days. Experiments with slowly growing mycobacteria yielded similar results.

Growth of *M. bovis* (BCG) was also strongly stimulated by 40 pMol/L RP-factor: growth occurred after 14 days whereas the control lacking RP-factor showed no growth after 90 days. Finally, RP-factor also stimulated the growth of *Mycobacterium tuberculosis*, *Mycobacterium smegmatis*, *Mycobacterium avium* and *Mycobacterium kansasii* (see Table 1).

TABLE 1

Purified <i>M. luteus</i> RP-factor stimulates growth of mycobacteria		
Organism	Bacterial growth [§]	
	RP-factor omitted	RP-factor added
<i>Mycobacterium tuberculosis</i> H37Ra	1.3 ± 1.9 (5)	110 ± 32 (5)
<i>Mycobacterium tuberculosis</i> H37Rv	1.5 ± 2 (4)	45 ± 28 (4)
<i>Mycobacterium avium</i>	0 (3)	>300 (3)
<i>M. bovis</i> (BCG)	0 (5)	54 ± 38 (5)
<i>M. smegmatis</i> *	0 (8)	225 ± 44 (8)
<i>Mycobacterium kansasii</i>	2.5 ± 2.5 (3)	90 ± 77 (3)

[§]Growth was estimated microscopically (magnification times 600) after 14 days of 5 incubation; ca. 50 µl of each culture was fixed, stained using Ziehl-Neelsen reagent and counted. Values in the body of the Table are average numbers of cells in a microscope field (10-20 fields counted) ± standard deviation with the number of determinations in parentheses. RP-factor (after elution from the Mono Q column and dialysis) was used at a concentration of circa 40 pMol/L; activity was lost after either trypsin treatment, heating (autoclaving) or filtration through a 12 kDa cutoff membrane.

*Washed cells of *M. smegmatis* were used for this experiment.

Isolation and Characterisation of the Gene Encoding the Second Homologue from *M. luteus*

A combination of inverse PCR using oligos G1 and G2 (see FIG. 2A) as primers, and cloning of suitably sized genomic restriction fragments, can be employed to isolate the gene encoding the second homologue from *M. luteus*. The sequence of the gene can then be determined, taking care to eliminate any possible PCR errors by analysis of genomic clones and direct sequencing of PCR fragments obtained by combining the products of multiple, independent PCR reactions. Comparative sequence analyses of the proteins from *M. luteus*, *M. leprae* and *M. tuberculosis* can then be used to refine predictions concerning residues, sequence motifs and structural motifs which may be important for biological function.

Over-Expression and Purification of *M. luteus* and *M. tuberculosis* Gene Products in *E. coli*

PCR primers can be designed, incorporating suitable restriction sites such that sequences encoding the secreted forms of the *M. luteus* and the *M. tuberculosis* RP-factors can be amplified and inserted, in the correct reading frame, into commercially available plasmids (pET or pCAL vectors). The PCR-amplified fragments can first be cloned in a pBlue-script KS II vector (Stratagene) so that their entire sequence can be verified, to eliminate possible PCR errors. (This material can also be employed for site-directed mutagenesis—vide infra.) The pET or pCAL constructs can then be employed to obtain controlled expression of large quantities of histidine- or calmodulin binding peptide-tagged proteins

that can be purified, essentially to homogeneity, in a single step. Finally, the tags used in protein purification can be removed (using enterokinase or thrombin, as appropriate).

Expression of RP-Factor from *Micrococcus luteus* in *E. coli*

Two primers [5'-GTCAGAATTCATATGGCCACCGTG-GACACCTGGG-3'] (SEQ ID NO: 46) and [5'-TGACG-GATCCTATTAGGCCTGCGGCAGGACGAG-3'] (SEQ ID NO: 47) were employed to amplify (5 cycles of 30 s at 94° C., 30 s at 60° C., 30 s at 72° C., followed by 15 cycles of 30 s at 94° C., 60 s at 72° C.) the RP-factor coding sequence (i.e. lacking the signal sequence) from the cloned 1.4 kbp SmaI fragment of genomic DNA. It was first established in *E. coli* DH5α as a 567 bp EcoRI-BamHI fragment in pMTL20 and then excised as a 562 bp NdeI-BamHI fragment, inserted into pET19b (Novagen) and re-established in *E. coli* DH5 α. The sequence of the PCR product and vector-insert junction in this plasmid, denoted pRPF1, was verified. RP-factor was expressed from RPF1 after transforming it into *E. coli* HSM174(DE3). The protein, containing a His10-tag at the N-terminus, was isolated by sonicating bacteria, previously grown to an OD_{600nm} = 0.6 and induced with 0.4 mM IPTG for 4 h, in a modified binding buffer (MBB—5 mM imidazole pH7.9/0.5M NaCl/20 mM Tris-HCl/8M urea) containing 5 mM DTT and 2 mM EDTA. After low speed centrifugation, low MW compounds, including EDTA and DTT, were removed by elution through a Sephadex™ G10 column pre-equilibrated with MBB. A Ni2+-chelation column (Ni2+-coordinated iminodiacetic acid immobilized on Sepharose™ 6B), was loaded with the G10 eluate, washed with 20 vol MBB and then successively eluted with four 10 vol aliquots of MBB containing 0.01 M, 0.05 M, 0.2 M and 1 M imidazole, respectively. The column was finally eluted with strip buffer (20 mM Tris-HCl, pH 7.9/100 mM EDTA/0.5 M NaCl). Monoclonal anti-(polyHis) antibodies (Sigma, clone His-1) were employed for immunoblot analysis of fractions subjected to SDS PAGE electrophoresis and electroblotted using standard methods. Fractions were dialysed against buffer 2 and assayed for biological activity as indicated above.

Analysis of Recombinant RP-Factor

The coding sequence corresponding to the secreted form of RP-factor, starting at residue A₃₉, was inserted into pET 19b to generate plasmid pRPF1 (vide infra). Extracts of IPTG-induced *E. coli* strain HSM174(DE3) containing pRPF 1 were challenged with a poly-His antibody. A strong signal was associated with a protein (apparent size 29 kDa, predicted size 22 kDa) which was eluted from the affinity column by 1M imidazole (FIG. 7A). The His-tagged protein from HSM174(DE3) reduced the apparent lag phase of viable cells of *M. luteus*, whereas the control (material eluted from the same column under the same conditions when an extract from cells containing plasmid vector only was applied) showed no activity (FIG. 7B). The association of biological activity with the recombinant protein, produced in *E. coli* containing pRPF, and the absence of biological activity in the isogenic control containing pET19b, demonstrates unequivocally that the active molecule is indeed a product of the *rpf* gene.

Antibody Preparation

A rabbit was immunized three times at one week intervals using recombinant RP-factor (the recombinant protein prepared as described above). The protein was administered at 300 µg of protein per injection in incomplete Freud's adjuvant (0.5 ml protein and 0.5 ml adjuvant) Blood was collected before administration was started and on the 11th day after the last injection. The immunoglobulin fraction was obtained by standard procedures using PEG. Antibodies were additionally purified on a protein G-Superose™ column according to

the standard (Pharmacia) protocol. The final protein concentration to was adjusted spectrophotometrically to 1 mg/ml.

Alternatively, monoclonal antibodies can be produced using established techniques.

Use of Anti-RP-Factor Antibody to Inhibit Bacterial Growth

Micrococcus luteus was inoculated at an initial density of 5×10^5 per ml into lactate minimal medium (LMM) and the OD_{600nm} was monitored at intervals. Growth of the cultures was monitored over 140 hours, and the presence of the anti-RP-factor serum (prepared as described above under "Antibody preparation") completely inhibited bacterial growth (see FIG. 8).

Expression of a *M. tuberculosis* RP-Factor in *E. coli*

Two primers [5'-ATCAGAAATTCATATGGACGACATC-GATTGGGACGC-3'] (SEQ ID NO: 48) and [5'-CGCAG-GATCCCTCAATCGTCCCTGCTCC-3'] (SEQ ID NO: 49) were employed to amplify (5 cycles of 30 s at 94° C., 30 s at 58° C., 30 s at 72° C., followed by 25 cycles of 30 s at 94° C., 60 s at 72° C.) the RP-factor coding sequence (i.e., lacking the signal sequence) from *M. tuberculosis* H37Rv genomic DNA. The PCR product was first established in *E. coli* DH5a as a 336 bp EcoRI-BamHI fragment in pMTL20 and then excised as a 331 bp NdeI-BamHI fragment, inserted into pET19b (Novagen) and re-established in *E. coli* DH5a. The sequence of the PCR product and vector-insert junction in this plasmid, denoted pRPF2, was verified. The *M. tuberculosis* RP-factor was expressed from pRPF2 after transforming it into *E. coli* HSM174(DE3). The protein, containing a His10-tag at the N-terminus, was isolated by sonicating bacteria, previously grown to an OD_{600 nm}=0.9 and induced with 0.4 mM IPTG for 4 h, in binding buffer (BB—5 mM imidazole pH7.9 /0.5M NaCl/20 mM Tris-HCl/8M urea). After low speed centrifugation, a Ni2+-chelation column (Ni2+-coordinated iminodiacetic acid immobilised on Sepharose™ 6B), was loaded with the supernatant, washed with 20 vol 13B, 20 vol BB containing 100 mM imidazole, and then eluted with 10 vol BB containing 0.5 M imidazole. Additional purification was achieved by Mono Q® column chromatography (vide infra, save that the salt gradient was from 0.1 M to 1M NaCl). Monoclonal anti-(polyHis) antibodies (Sigma, clone His-1) were employed for immunoblot analysis of fractions subjected to SDS PAGE electrophoresis and electroblotted using standard methods. Fractions were dialysed against buffer 2 and assayed for biological activity as indicated above.

Analysis of a Recombinant *M. tuberculosis* RP-Factor

The coding sequence corresponding to the secreted form of the *M. tuberculosis* RP-factor (g1655671; acc. no. Z81368), starting at residue D₅₀, was inserted into pET19b to generate plasmid pRPF2 (vide infra). Extracts of IPTG-induced *E. coli* strain HSM174(DE3) containing pRPF2 were challenged with a poly-His antibody. A strong signal was associated with a protein which was eluted from the affinity column by 0.5M imidazole. The histidine-tagged protein from HSM174(DE3) caused a slight but significant enhancement of the growth of *M. tuberculosis* H37Rv, as shown in FIG. 10. It also stimulated the growth of *M. luteus* in LMM. The control culture attained a final OD_{600nm} of 1.0, whereas cultures containing the RP-factor (1:100,000 dilution) attained a final OD_{600nm} of between 2.0 and 6.0.

Effect of *M. luteus* RP-Factor on Growth of *Mycobacterium tuberculosis* Cells Isolated from Macrophages

In three independent experiments, dormant/latent *M. tuberculosis* cells isolated from cultured murine peritoneal macrophages were resuscitated by the *M. luteus* RP-factor. The total number of *M. tuberculosis* cells in the heterogeneous suspension obtained from murine macrophages was

determined microscopically. The viable cell count was determined by plating on agar-solidified Sauton medium containing 10% (v/v) supplement (which contains, per liter, 50 g bovine serum albumin, 20 g glucose, 8.5 g NaCl) or by the MPN method, using liquid Sauton medium containing 10% (v/v) supplement (see above).

The viable count (MPN) of these cell suspensions was enhanced between 25 and 2,500 times by the presence of the *M. luteus* RP-factor (added at a final concentration of 10 ng/ml) (see Table 2). All values in the body of the table are numbers of bacteria per ml suspension

Peritoneal macrophages were obtained from white mice (wild type) by a standard protocol. Infection of macrophages by *M. tuberculosis* "Academiya" (laboratory strain) was performed in vivo by intraperitoneal injection of 10^6 cells (total count) per mouse followed by incubation for 6 days (1st passage). For the second and third passages macrophage cells in monolayers were infected using *M. tuberculosis* cells isolated from macrophages from the previous passage.

TABLE 2

Effect of <i>M. luteus</i> RP-factor on growth of <i>Mycobacterium tuberculosis</i> cells isolated from macrophages				
Experiment	Total count [x] (determined microscopically)	Viable count (determined by plating)	Viable count (MPN)	MPN in presence of RP-factor
I	$10^6 > x > 10^5$	90	70	4.10^3
II	$10^6 > x > 10^5$	9	40	1.10^3
III	2.10^6	<1	<1	24.10^3

Macrophages were grown as a monolayer on plastic petri dishes (10^6 cells/5 cm²) in standard RPMI medium containing gentamicin and penicillin (10 ig/ml, each) under standard conditions (CO₂/O₂ mixture in a 37° C. incubator). *M. tuberculosis* cells were recovered from macrophages by passing them repeatedly through a thin syringe needle. Macrophage cell debris was removed by low speed centrifugation and *M. tuberculosis* cells were then collected by centrifugation at higher speed.

Effect of yabE and yocH Knockout Mutations on Growth of *Bacillus subtilis*

The entire yabE coding region together with flanking sequences was amplified from *B. subtilis* genomic DNA using primers D11 [5'-GAAGAGAATTCCTTCCAT-CACGA-3'] (SEQ ID NO: 50) and D12 [5'-CCAAACGAAT-TCGGTCAATCAC-31 (SEQ ID NO: 51) as a 1803 bp product. A 1186 bp HindIII-BclI fragment encompassing the 3' end of the coding sequence was excised from the PCR product, ligated with HindIII+BamHI-digested pMTL20, and used to transform *E. coli* strain DH5a with selection for ampicillin-resistance. Plasmid pYABE was isolated from one of the transformants. A 763 bp HindIII-BamHI fragment from entirely within the yabE coding sequence was excised from the pYABE, ligated with HindIII+BamHI-digested pMUTIN4, an integrating plasmid that may be employed for generating knockout mutations in *B. subtilis* (Edwards & Errington, 1997, Molecular Microbiology, 24, 905-915) and used to transform *E. coli* strain XL 1-Blue with selection for ampicillin-resistance. Plasmid pYAB2, containing an internal segment of the yabE coding sequence, was isolated from one of the transformants. A 1207 bp HindIII-EcoRI fragment encompassing the 3' end of the yabE coding sequence was excised from pYABE, ligated with HindIII+EcoRI digested pMUTIN4 and used to transform *E. coli* strain XL1-Blue with

selection for ampicillin-resistance. Plasmid pYAB3, containing the 3' end of the yabE coding sequence, was isolated from one of the transformants.

The entire yocH coding region together with flanking sequences was amplified from *B. subtilis* genomic DNA using primers D10 [5'-GCAAGGATCCCAGAC-TAAAAAACAG-3'] (SEQ ID NO: 52) and D9 [5'-ATCAG-GATCCATATTATTAGTTTAAGA-3'] (SEQ ID NO: 53) as a 1145 bp product. A 358 bp HpaI fragment from entirely within the yocH coding sequence was excised from the PCR product, ligated with SmaI-digested pMTL20, and used to transform *E. coli* strain XL1-Blue with selection for ampicillin-resistance. Plasmid pYOC2a, containing an internal segment of the yocH coding sequence, was isolated from one of the transformants. The insert in this plasmid was then excised from pYOC2a as a 385 bp EcoRI-HindIII fragment and inserted into pMUTIN4, to yield pYOC2. A 307 bp HindIII-BamHI fragment encompassing the 3' end of the yocH coding sequence was excised from the 1145 bp PCR product, ligated with HindIII+BamHI digested pMUTIN4, and used to trans-

form *E. coli* strain DH5 α with selection for ampicillin-resistance. Plasmid pYOC3, containing a DNA segment encompassing the 3' end of the yocH coding sequence, was isolated from one of the transformants.

Plasmids pYAB2, pYAB3, pYOC2 and pYOC3 were linearised with ApaI, which cleaves once in the pMUTIN4 vector sequences, ligated with T4 DNA ligase and employed to transform *Bacillus subtilis* strain SA253 nonA nonB leuA8 arg-15 with selection for resistance to erythromycin on a rich nutrient medium (LB+1 μ g Em/ml). Em^R transformants were then picked and verified by Southern hybridization. Using the integrating plasmid as probe, and digesting the chromosomal DNA with ApaI, strains harbouring a single copy of the integrated plasmid gave two hybridising bands whereas the wild type (and any spontaneous Em^R mutants that were present) gave a single hybridising band.

Analysis of the products of transformation with each of the four plasmids indicates that yabE and yocH gene products are required for growth (at least under certain conditions) in *B. subtilis*.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 65

<210> SEQ ID NO 1

<211> LENGTH: 362

<212> TYPE: PRT

<213> ORGANISM: *Mycobacterium tuberculosis*

<400> SEQUENCE: 1

```

Met Leu Arg Leu Val Val Gly Ala Leu Leu Leu Val Leu Ala Phe Ala
1           5           10           15
Gly Gly Tyr Ala Val Ala Ala Cys Lys Thr Val Thr Leu Thr Val Asp
20           25           30
Gly Thr Ala Met Arg Val Thr Thr Met Lys Ser Arg Val Ile Asp Ile
35           40           45
Val Glu Glu Asn Gly Phe Ser Val Asp Asp Arg Asp Asp Leu Tyr Pro
50           55           60
Ala Ala Gly Val Gln Val His Asp Ala Asp Thr Ile Val Leu Arg Arg
65           70           75           80
Ser Arg Pro Leu Gln Ile Ser Leu Asp Gly His Asp Ala Lys Gln Val
85           90           95
Trp Thr Thr Ala Ser Thr Val Asp Glu Ala Leu Ala Gln Leu Ala Met
100          105          110
Thr Asp Thr Ala Pro Ala Ala Ala Ser Arg Ala Ser Arg Val Pro Leu
115          120          125
Ser Gly Met Ala Leu Pro Val Val Ser Ala Lys Thr Val Gln Leu Asn
130          135          140
Asp Gly Gly Leu Val Arg Thr Val His Leu Pro Ala Pro Asn Val Ala
145          150          155          160
Gly Leu Leu Ser Ala Ala Gly Val Pro Leu Leu Gln Ser Asp His Val
165          170          175
Val Pro Ala Ala Thr Ala Pro Ile Val Glu Gly Met Gln Ile Gln Val
180          185          190
Thr Arg Asn Arg Ile Lys Lys Val Thr Glu Arg Leu Pro Leu Pro Pro
195          200          205
Asn Ala Arg Arg Val Glu Asp Pro Glu Met Asn Met Ser Arg Glu Val
210          215          220

```

-continued

Val	Glu	Asp	Pro	Gly	Val	Pro	Gly	Thr	Gln	Asp	Val	Thr	Phe	Ala	Val
225					230					235					240
Ala	Glu	Val	Asn	Gly	Val	Glu	Thr	Gly	Arg	Leu	Pro	Val	Ala	Asn	Val
				245					250					255	
Val	Val	Thr	Pro	Ala	His	Glu	Ala	Val	Val	Arg	Val	Gly	Thr	Lys	Pro
			260					265					270		
Gly	Thr	Glu	Val	Pro	Pro	Val	Ile	Asp	Gly	Ser	Ile	Trp	Asp	Ala	Ile
		275					280					285			
Ala	Gly	Cys	Glu	Ala	Gly	Gly	Asn	Trp	Ala	Ile	Asn	Thr	Gly	Asn	Gly
	290					295					300				
Tyr	Tyr	Gly	Gly	Val	Gln	Phe	Asp	Gln	Gly	Thr	Trp	Glu	Ala	Asn	Gly
305					310					315					320
Gly	Leu	Arg	Tyr	Ala	Pro	Arg	Ala	Asp	Leu	Ala	Thr	Arg	Glu	Glu	Gln
				325					330					335	
Ile	Ala	Val	Ala	Glu	Val	Thr	Arg	Leu	Arg	Gln	Gly	Trp	Gly	Ala	Trp
			340					345					350		
Pro	Val	Cys	Ala	Ala	Arg	Ala	Gly	Ala	Arg						
		355					360								

<210> SEQ ID NO 2
 <211> LENGTH: 188
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 2

Met	Pro	Val	Gly	Trp	Leu	Trp	Arg	Ala	Arg	Thr	Ala	Lys	Gly	Thr	Thr
1				5					10					15	
Leu	Lys	Asn	Ala	Arg	Thr	Thr	Leu	Ile	Ala	Ala	Ala	Ile	Ala	Gly	Thr
			20				25						30		
Leu	Val	Thr	Thr	Ser	Pro	Ala	Gly	Ile	Ala	Asn	Ala	Asp	Asp	Ala	Gly
		35				40					45				
Leu	Asp	Pro	Asn	Ala	Ala	Ala	Gly	Pro	Asp	Ala	Val	Gly	Phe	Asp	Pro
	50					55				60					
Asn	Leu	Pro	Pro	Ala	Pro	Asp	Ala	Ala	Pro	Val	Asp	Thr	Pro	Pro	Ala
65				70					75						80
Pro	Glu	Asp	Ala	Gly	Phe	Asp	Pro	Asn	Leu	Pro	Pro	Pro	Leu	Ala	Pro
			85					90						95	
Asp	Phe	Leu	Ser	Pro	Pro	Ala	Glu	Glu	Ala	Pro	Pro	Val	Pro	Val	Ala
		100					105						110		
Tyr	Ser	Val	Asn	Trp	Asp	Ala	Ile	Ala	Gln	Cys	Glu	Ser	Gly	Gly	Asn
	115					120						125			
Trp	Ser	Ile	Asn	Thr	Gly	Asn	Gly	Tyr	Tyr	Gly	Gly	Leu	Arg	Phe	Thr
	130				135						140				
Ala	Gly	Thr	Trp	Arg	Ala	Asn	Gly	Gly	Ser	Gly	Ser	Ala	Ala	Asn	Ala
145				150					155					160	
Ser	Arg	Glu	Glu	Gln	Ile	Arg	Val	Ala	Glu	Asn	Val	Leu	Arg	Ser	Gln
			165				170						175		
Gly	Ile	Arg	Ala	Trp	Pro	Val	Cys	Gly	Arg	Arg	Gly				
		180					185								

<210> SEQ ID NO 3
 <211> LENGTH: 174
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium leprae

<400> SEQUENCE: 3

Met	Ser	Glu	Ser	Tyr	Arg	Lys	Leu	Thr	Thr	Ser	Ser	Ile	Ile	Val	Ala
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

-continued

1	5	10	15
Lys Ile Thr Phe Thr Gly Ala Met Leu Asp Gly Ser Ile Ala Leu Ala	20	25	30
Gly Gln Ala Ser Pro Ala Thr Asp Ser Glu Trp Asp Gln Val Ala Arg	35	40	45
Cys Glu Ser Gly Gly Asn Trp Ser Ile Asn Thr Gly Asn Gly Tyr Leu	50	55	60
Gly Gly Leu Gln Phe Ser Gln Gly Thr Trp Ala Ser His Gly Gly Gly	65	70	75
Glu Tyr Ala Pro Ser Ala Gln Leu Ala Thr Arg Glu Gln Gln Ile Ala	85	90	95
Val Ala Glu Arg Val Leu Ala Thr Gln Gly Ser Gly Ala Trp Pro Ala	100	105	110
Cys Gly His Gly Leu Ser Gly Pro Ser Leu Gln Glu Val Leu Pro Ala	115	120	125
Gly Met Gly Ala Pro Trp Ile Asn Gly Ala Pro Ala Pro Leu Ala Pro	130	135	140
Pro Pro Pro Ala Glu Pro Ala Pro Pro Gln Pro Pro Ala Asp Asn Phe	145	150	155
Pro Pro Thr Pro Gly Asp Val Pro Ser Pro Leu Ala Arg Pro	165	170	

<210> SEQ ID NO 4

<211> LENGTH: 407

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 4

Met Ser Gly Arg His Arg Lys Pro Thr Thr Ser Asn Val Ser Val Ala	1	5	10	15
Lys Ile Ala Phe Thr Gly Ala Val Leu Gly Gly Gly Gly Ile Ala Met	20	25	30	
Ala Ala Gln Ala Thr Ala Ala Thr Asp Gly Glu Trp Asp Gln Val Ala	35	40	45	
Arg Cys Glu Ser Gly Gly Asn Trp Ser Ile Asn Thr Gly Asn Gly Tyr	50	55	60	
Leu Gly Gly Leu Gln Phe Thr Gln Ser Thr Trp Ala Ala His Gly Gly	65	70	75	80
Gly Glu Phe Ala Pro Ser Ala Gln Leu Ala Ser Arg Glu Gln Gln Ile	85	90	95	
Ala Val Gly Glu Arg Val Leu Ala Thr Gln Gly Arg Gly Ala Trp Pro	100	105	110	
Val Cys Gly Arg Gly Leu Ser Asn Ala Thr Pro Arg Glu Val Leu Pro	115	120	125	
Ala Ser Ala Ala Met Asp Ala Pro Leu Asp Ala Ala Ala Val Asn Gly	130	135	140	
Glu Pro Ala Pro Leu Ala Pro Pro Pro Ala Asp Pro Ala Pro Pro Val	145	150	155	160
Glu Leu Ala Ala Asn Asp Leu Pro Ala Pro Leu Gly Glu Pro Leu Pro	165	170	175	
Ala Ala Pro Ala Asp Pro Ala Pro Pro Ala Asp Leu Ala Pro Pro Ala	180	185	190	
Pro Ala Asp Val Ala Pro Pro Val Glu Leu Ala Val Asn Asp Leu Pro	195	200	205	
Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala Asp Pro Ala Pro				

-continued

210	215	220
Pro Ala Asp Leu Ala	Pro Pro Ala Pro Ala	Asp Leu Ala Pro Pro Ala
225	230	235 240
Pro Ala Asp Leu Ala	Pro Pro Ala Pro Ala	Asp Leu Ala Pro Pro Val
245	250	255
Glu Leu Ala Val Asn Asp Leu Pro	Ala Pro Leu Gly Glu Pro Leu Pro	
260	265	270
Ala Ala Pro Ala Glu Leu Ala Pro	Pro Ala Asp Leu Ala Pro Ala Ser	
275	280	285
Ala Asp Leu Ala Pro Pro Ala Pro	Ala Asp Leu Ala Pro Pro Ala Pro	
290	295	300
Ala Glu Leu Ala Pro Pro Ala Pro	Ala Asp Leu Ala Pro Pro Ala Ala	
305	310	315 320
Val Asn Glu Gln Thr Ala Pro Gly Asp Gln Pro	Ala Thr Ala Pro Gly	
325	330	335
Gly Pro Val Gly Leu Ala Thr Asp Leu Glu Leu Pro	Glu Pro Asp Pro	
340	345	350
Gln Pro Ala Asp Ala Pro Pro Pro Gly Asp Val Thr Glu Ala Pro	Ala	
355	360	365
Glu Thr Pro Gln Val Ser Asn Ile Ala Tyr Thr Lys Lys Leu Trp Gln		
370	375	380
Ala Ile Arg Ala Gln Asp Val Cys Gly Asn Asp Ala Leu Asp Ser Leu		
385	390	395 400
Ala Gln Pro Tyr Val Ile Gly		
405		

<210> SEQ ID NO 5
 <211> LENGTH: 155
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium leprae

<400> SEQUENCE: 5

Met Pro Gly Glu Met Leu Asp Val Arg Lys Leu Cys Lys Leu Phe Val	
1 5 10 15	
Lys Ser Ala Val Val Ser Gly Ile Val Thr Ala Ser Met Ala Leu Ser	
20 25 30	
Thr Ser Thr Gly Met Ala Asn Ala Val Pro Arg Glu Pro Asn Trp Asp	
35 40 45	
Ala Val Ala Gln Cys Glu Ser Gly Arg Asn Trp Arg Ala Asn Thr Gly	
50 55 60	
Asn Gly Phe Tyr Gly Gly Leu Gln Phe Lys Pro Thr Ile Trp Ala Arg	
65 70 75 80	
Tyr Gly Gly Val Gly Asn Pro Ala Gly Ala Ser Arg Glu Gln Gln Ile	
85 90 95	
Thr Val Ala Asn Arg Val Leu Ala Asp Gln Gly Leu Asp Ala Trp Pro	
100 105 110	
Lys Cys Gly Ala Ala Ser Asp Leu Pro Ile Thr Leu Trp Ser His Pro	
115 120 125	
Ala Gln Gly Val Lys Gln Ile Ile Asn Asp Ile Ile Gln Met Gly Asp	
130 135 140	
Thr Thr Leu Ala Ala Ile Ala Leu Asn Gly Leu	
145 150 155	

<210> SEQ ID NO 6
 <211> LENGTH: 176
 <212> TYPE: PRT

-continued

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 6

Met His Pro Leu Pro Ala Asp His Gly Arg Ser Arg Cys Asn Arg His
 1 5 10 15

Pro Ile Ser Pro Leu Ser Leu Ile Gly Asn Ile Ser Ala Thr Ser Gly
 20 25 30

Asp Met Ser Ser Met Thr Arg Ile Ala Lys Pro Leu Ile Lys Ser Ala
 35 40 45

Met Ala Ala Gly Leu Val Thr Ala Ser Met Ser Leu Ser Thr Ala Val
 50 55 60

Ala His Ala Gly Pro Ser Pro Asn Trp Asp Ala Val Ala Gln Cys Glu
 65 70 75 80

Ser Gly Gly Asn Trp Ala Ala Asn Thr Gly Asn Gly Lys Tyr Gly Gly
 85 90 95

Leu Gln Phe Lys Pro Ala Thr Trp Ala Ala Phe Gly Gly Val Gly Asn
 100 105 110

Pro Ala Ala Ala Ser Arg Glu Gln Gln Ile Ala Val Ala Asn Arg Val
 115 120 125

Leu Ala Glu Gln Gly Leu Asp Ala Trp Pro Thr Cys Gly Ala Ala Ser
 130 135 140

Gly Leu Pro Ile Ala Leu Trp Ser Lys Pro Ala Gln Gly Ile Lys Gln
 145 150 155 160

Ile Ile Asn Glu Ile Ile Trp Ala Gly Ile Gln Ala Ser Ile Pro Arg
 165 170 175

<210> SEQ ID NO 7

<211> LENGTH: 154

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 7

Met Thr Pro Gly Leu Leu Thr Thr Ala Gly Ala Gly Arg Pro Arg Asp
 1 5 10 15

Arg Cys Ala Arg Ile Val Cys Thr Val Phe Ile Glu Thr Ala Val Val
 20 25 30

Ala Thr Met Phe Val Ala Leu Leu Gly Leu Ser Thr Ile Ser Ser Lys
 35 40 45

Ala Asp Asp Ile Asp Trp Asp Ala Ile Ala Gln Cys Glu Ser Gly Gly
 50 55 60

Asn Trp Ala Ala Asn Thr Gly Asn Gly Leu Tyr Gly Gly Leu Gln Ile
 65 70 75 80

Ser Gln Ala Thr Trp Asp Ser Asn Gly Gly Val Gly Ser Pro Ala Ala
 85 90 95

Ala Ser Pro Gln Gln Gln Ile Glu Val Ala Asp Asn Ile Met Lys Thr
 100 105 110

Gln Gly Pro Gly Ala Trp Pro Lys Cys Ser Ser Cys Ser Gln Gly Asp
 115 120 125

Ala Pro Leu Gly Ser Leu Thr His Ile Leu Thr Phe Leu Ala Ala Glu
 130 135 140

Thr Gly Gly Cys Ser Gly Ser Arg Asp Asp
 145 150

<210> SEQ ID NO 8

<211> LENGTH: 99

<212> TYPE: PRT

<213> ORGANISM: Streptomyces coelicolor

-continued

<400> SEQUENCE: 8

Ile Arg Thr Ala Ala Val Thr Leu Val Ala Ala Thr Ala Leu Gly Ala
 1 5 10 15
 Thr Gly Glu Ala Val Ala Ala Pro Ser Ala Pro Leu Arg Thr Asp Trp
 20 25 30
 Asp Ala Ile Ala Ala Cys Glu Ser Ser Gly Asn Trp Gln Ala Asn Thr
 35 40 45
 Gly Asn Gly Tyr Tyr Gly Gly Leu Gln Phe Ala Arg Ser Ser Trp Ile
 50 55 60
 Ala Ala Gly Gly Leu Lys Tyr Ala Pro Arg Ala Asp Leu Ala Thr Arg
 65 70 75 80
 Gly Glu Gln Ile Ala Val Ala Glu Arg Leu Ala Arg Leu Gln Gly Met
 85 90 95
 Ser Ala Trp

<210> SEQ ID NO 9

<211> LENGTH: 438

<212> TYPE: PRT

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 9

Met Gly Glu Arg Glu Gly Arg Val Asp Ser Leu Leu Asp Thr Leu Tyr
 1 5 10 15
 Asn Leu Ser Glu Glu Lys Glu Ala Phe Phe Ile Thr Gln Lys Met Lys
 20 25 30
 Lys Leu Phe Ser Val Lys Leu Ser Lys Ser Lys Val Ile Leu Val Ala
 35 40 45
 Ala Cys Leu Leu Leu Ala Gly Ser Gly Thr Ala Tyr Ala Ala His Glu
 50 55 60
 Leu Thr Lys Gln Ser Val Ser Val Ser Ile Asn Gly Lys Lys Lys His
 65 70 75 80
 Ile Arg Thr His Ala Asn Thr Val Gly Asp Leu Leu Glu Thr Leu Asp
 85 90 95
 Ile Lys Thr Arg Asp Glu Asp Lys Ile Thr Pro Ala Lys Gln Thr Lys
 100 105 110
 Ile Thr Ala Asp Met Asp Val Val Tyr Glu Ala Ala Lys Pro Val Lys
 115 120 125
 Leu Thr Ile Asn Gly Glu Glu Lys Thr Leu Trp Ser Thr Ala Lys Thr
 130 135 140
 Val Gly Ala Leu Leu Asp Glu Gln Asp Val Asp Val Lys Glu Gln Asp
 145 150 155 160
 Gln Ile Asp Pro Ala Ile Asp Thr Asp Ile Ser Lys Asp Met Lys Ile
 165 170 175
 Asn Ile Glu Pro Ala Phe Gln Val Thr Val Asn Asp Ala Gly Lys Gln
 180 185 190
 Lys Lys Ile Trp Thr Thr Ser Thr Thr Val Ala Asp Phe Leu Lys Gln
 195 200 205
 Gln Lys Met Asn Ile Lys Asp Glu Asp Lys Ile Lys Pro Ala Leu Asp
 210 215 220
 Ala Lys Leu Thr Lys Gly Lys Ala Asp Ile Thr Ile Thr Arg Ile Glu
 225 230 235 240
 Lys Val Thr Asp Val Val Glu Glu Lys Ile Ala Phe Asp Val Lys Lys
 245 250 255
 Gln Glu Asp Ala Ser Leu Glu Lys Gly Lys Glu Lys Val Val Gln Lys

-continued

260	265	270
Gly Lys Glu Gly Lys Leu Lys Lys His Phe Glu Val Val Lys Glu Asn		
275	280	285
Gly Lys Glu Val Ser Arg Glu Leu Val Lys Glu Glu Thr Ala Glu Gln		
290	295	300
Ser Lys Asp Lys Val Ile Ala Val Gly Thr Lys Gln Ser Ser Pro Lys		
305	310	315
Phe Glu Thr Val Ser Ala Ser Gly Asp Ser Lys Thr Val Val Ser Arg		
325	330	335
Ser Asn Glu Ser Thr Gly Lys Val Met Thr Val Ser Ser Thr Ala Tyr		
340	345	350
Thr Ala Ser Cys Ser Gly Cys Ser Gly His Thr Ala Thr Gly Val Asn		
355	360	365
Leu Lys Asn Asn Pro Asn Ala Lys Val Ile Ala Val Asp Pro Asn Val		
370	375	380
Ile Pro Leu Gly Ser Lys Val His Val Glu Gly Tyr Gly Tyr Ala Ile		
385	390	395
Ile Ala Ala Asp Thr Gly Ser Ala Ile Lys Gly Asn Lys Ile Asp Val		
405	410	415
Phe Phe Pro Ser Lys Ser Asp Ala Ser Asn Trp Gly Val Lys Thr Val		
420	425	430
Ser Val Lys Val Leu Asn		
435		

<210> SEQ ID NO 10
 <211> LENGTH: 288
 <212> TYPE: PRT
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 10

Met Lys Lys Thr Ile Met Ser Phe Val Ala Val Ala Ala Leu Ser Thr		
1	5	10
Thr Ala Phe Gly Ala His Ala Ser Ala Lys Glu Ile Thr Val Gln Lys		
20	25	30
Gly Asp Thr Leu Trp Gly Ile Ser Gln Lys Asn Gly Val Asn Leu Lys		
35	40	45
Asp Leu Lys Glu Trp Asn Lys Leu Thr Ser Asp Lys Ile Ile Ala Gly		
50	55	60
Glu Lys Leu Thr Ile Ser Ser Glu Glu Thr Thr Thr Thr Gly Gln Tyr		
65	70	75
Thr Ile Lys Ala Gly Asp Thr Leu Ser Lys Ile Ala Gln Lys Phe Gly		
85	90	95
Thr Thr Val Asn Asn Leu Lys Val Trp Asn Asn Leu Ser Ser Asp Met		
100	105	110
Ile Tyr Ala Gly Ser Thr Leu Ser Val Lys Gly Gln Ala Thr Ala Ala		
115	120	125
Asn Thr Ala Thr Glu Asn Ala Gln Thr Asn Ala Pro Gln Ala Ala Pro		
130	135	140
Lys Gln Glu Ala Val Gln Lys Glu Gln Pro Lys Gln Glu Ala Val Gln		
145	150	155
Gln Gln Pro Lys Gln Glu Thr Lys Ala Glu Ala Glu Thr Ser Val Asn		
165	170	175
Thr Glu Glu Lys Ala Val Gln Ser Asn Thr Asn Asn Gln Glu Ala Ser		
180	185	190
Lys Glu Leu Thr Val Thr Ala Thr Ala Tyr Thr Ala Asn Asp Gly Gly		

-continued

195	200	205
Ile Ser Gly Val Thr Ala Thr Gly Ile Asp Leu Asn Lys Asn Pro Asn		
210	215	220
Ala Lys Val Ile Ala Val Asp Pro Asn Val Ile Pro Leu Gly Ser Lys		
225	230	235
Val Tyr Val Glu Gly Tyr Gly Glu Ala Thr Thr Ala Ala Asp Thr Gly		
245	250	255
Gly Ala Ile Lys Gly Asn Lys Ile Asp Val Phe Val Pro Glu Lys Ser		
260	265	270
Ser Ala Tyr Arg Trp Gly Asn Lys Thr Val Lys Ile Lys Ile Leu Asn		
275	280	285

<210> SEQ ID NO 11
 <211> LENGTH: 320
 <212> TYPE: PRT
 <213> ORGANISM: Clostridium acetobutylicum
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (3)..(4)
 <223> OTHER INFORMATION: Variable amino acid

<400> SEQUENCE: 11

Lys Arg Xaa Xaa Ala Val Ile Leu Met Val Ala Val Ile Phe Thr Ile		
1	5	10
Ile Ser Ser Met Lys Lys Asn Ile Thr Val Asn Ile Asp Gly Lys Thr		
20	25	30
Ser Lys Ile Ile Thr Tyr Lys Ser Asn Glu Gly Ser Ile Leu Ser Lys		
35	40	45
Asn Asn Ile Leu Val Gly Pro Lys Asp Lys Ile Gln Pro Ala Leu Asp		
50	55	60
Thr Asn Leu Lys Asn Gly Asp Lys Ile Tyr Ile Lys Lys Ala Ile Ser		
65	70	75
Val Glu Val Ala Val Asp Gly Lys Val Arg Arg Val Lys Ser Ser Glu		
85	90	95
Glu Thr Val Ser Lys Met Leu Lys Ala Glu Lys Ile Pro Leu Ser Lys		
100	105	110
Val Asp Lys Val Asn Ile Ser Arg Asn Ala Ala Ile Lys Lys Asn Met		
115	120	125
Lys Ile Ser Ile Thr Arg Val Asn Ser Gln Ile Thr Lys Glu Asn Gln		
130	135	140
Gln Val Asp Phe Pro Thr Glu Val Ile Ser Asp Asp Ser Met Gly Asn		
145	150	155
Asp Glu Lys Gln Val Ile Gln Gln Gly Gln Ala Gly Glu Lys Glu Val		
165	170	175
Phe Thr Lys Ile Val Tyr Glu Asp Gly Lys Ala Val Ser Lys Glu Ile		
180	185	190
Val Gly Glu Val Ile Lys Lys Glu Pro Thr Lys Gln Val Phe Lys Val		
195	200	205
Gly Thr Leu Gly Val Leu Lys Pro Asp Arg Gly Gly Arg Val Leu Tyr		
210	215	220
Lys Lys Ser Leu Gln Val Leu Ala Thr Ala Tyr Thr Asp Asp Phe Ser		
225	230	235
Phe Gly Ile Thr Ala Ser Gly Thr Lys Val Lys Arg Asp Ser Asp Gly		
245	250	255
Tyr Ser Ser Ile Ala Val Asp Pro Thr Val Ile Pro Leu Gly Thr Lys		
260	265	270

-continued

Leu Tyr Val Pro Gly Tyr Gly Tyr Gly Val Val Ala Glu Asp Thr Gly
 275 280 285

Gly Ala Ile Lys Gly Asn Arg Leu Asp Leu Phe Phe Thr Ser Glu Arg
 290 295 300

Glu Cys Tyr Asp Trp Gly Ala Lys Asn Val Thr Val Tyr Ile Leu Lys
 305 310 315 320

<210> SEQ ID NO 12
 <211> LENGTH: 81
 <212> TYPE: PRT
 <213> ORGANISM: Clostridium perfringens

<400> SEQUENCE: 12

Ala Glu Ala Tyr Thr Ala Ser Gly Met His Val Leu Arg Asp Pro Asn
 1 5 10 15

Gly Tyr Ser Thr Ile Ala Val Asp Pro Ser Val Ile Pro Leu Gly Thr
 20 25 30

Lys Leu Tyr Val Glu Gly Tyr Gly Tyr Ala Ile Ile Ala Ala Asp Thr
 35 40 45

Gly Gly Ala Ile Lys Gly Asn Arg Val Asp Leu Phe Phe Asn Thr Glu
 50 55 60

Ala Glu Ala Ser Asn Trp Gly Val Arg Asn Leu Asp Val Tyr Ile Leu
 65 70 75 80

Asn

<210> SEQ ID NO 13
 <211> LENGTH: 51
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: RP-factor
 C-terminal domain peptide

<400> SEQUENCE: 13

Thr Ile Val Val Lys Ser Gly Asp Ser Leu Trp Thr Leu Ala Asn Glu
 1 5 10 15

Tyr Glu Val Glu Gly Gly Trp Thr Ala Leu Tyr Glu Ala Asn Lys Gly
 20 25 30

Ala Val Ser Asp Ala Ala Val Ile Tyr Val Gly Gln Glu Leu Val Leu
 35 40 45

Pro Gln Ala
 50

<210> SEQ ID NO 14
 <211> LENGTH: 46
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 14

Thr Ile Lys Val Lys Ser Gly Asp Ser Leu Trp Lys Leu Ser Arg Gln
 1 5 10 15

Tyr Asp Thr Thr Ile Ser Ala Leu Lys Ser Glu Asn Lys Leu Lys Ser
 20 25 30

Thr Val Leu Tyr Val Gly Gln Ser Leu Lys Val Pro Glu Ser
 35 40 45

<210> SEQ ID NO 15
 <211> LENGTH: 44

-continued

<212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 15

Thr Ile Lys Val Lys Ser Gly Asp Ser Leu Trp Lys Leu Ala Gln Thr
 1 5 10 15

Tyr Asn Thr Ser Val Ala Ala Leu Thr Ser Ala Asn His Leu Ser Thr
 20 25 30

Thr Val Leu Ser Ile Gly Gln Thr Leu Thr Ile Pro
 35 40

<210> SEQ ID NO 16
 <211> LENGTH: 43
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 16

Thr Tyr Thr Val Lys Ser Gly Asp Ser Leu Trp Val Ile Ala Gln Lys
 1 5 10 15

Phe Asn Val Thr Ala Gln Gln Ile Arg Glu Lys Asn Asn Leu Lys Thr
 20 25 30

Asp Val Leu Gln Val Gly Gln Lys Leu Val Ile
 35 40

<210> SEQ ID NO 17
 <211> LENGTH: 43
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 17

Lys Tyr Thr Val Lys Ser Gly Asp Ser Leu Trp Lys Ile Ala Asn Asn
 1 5 10 15

Ile Asn Leu Thr Val Gln Gln Ile Arg Asn Ile Asn Asn Leu Lys Ser
 20 25 30

Asp Val Leu Tyr Val Gly Gln Val Leu Lys Leu
 35 40

<210> SEQ ID NO 18
 <211> LENGTH: 45
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 18

Thr Tyr Thr Val Lys Ser Gly Asp Thr Ile Trp Ala Leu Ser Ser Lys
 1 5 10 15

Tyr Gly Thr Ser Val Gln Asn Ile Met Ser Trp Asn Asn Leu Ser Ser
 20 25 30

Ser Ser Ile Tyr Val Gly Gln Val Leu Ala Val Lys Gln
 35 40 45

<210> SEQ ID NO 19
 <211> LENGTH: 45

-continued

<212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 19

Thr His Ala Val Lys Ser Gly Asp Thr Ile Trp Ala Leu Ser Val Lys
 1 5 10 15
 Tyr Gly Val Ser Val Gln Asp Ile Met Ser Trp Asn Asn Leu Ser Ser
 20 25 30
 Ser Ser Ile Tyr Val Gly Gln Lys Leu Ala Ile Lys Gln
 35 40 45

<210> SEQ ID NO 20
 <211> LENGTH: 46
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 20

Ser Val Lys Val Lys Ser Gly Asp Thr Leu Trp Ala Leu Ser Val Lys
 1 5 10 15
 Tyr Lys Thr Ser Ile Ala Gln Leu Lys Ser Trp Asn His Leu Ser Ser
 20 25 30
 Asp Thr Ile Tyr Ile Gly Gln Asn Leu Ile Val Ser Gln Ser
 35 40 45

<210> SEQ ID NO 21
 <211> LENGTH: 43
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 21

Thr Tyr Thr Val Lys Ser Gly Asp Thr Leu Trp Gly Ile Ser Gln Arg
 1 5 10 15
 Tyr Gly Ile Ser Val Ala Gln Ile Gln Ser Ala Asn Asn Leu Lys Ser
 20 25 30
 Thr Ile Ile Tyr Ile Gly Gln Lys Leu Leu Leu
 35 40

<210> SEQ ID NO 22
 <211> LENGTH: 60
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 22

Thr Tyr Thr Val Lys Lys Gly Asp Thr Leu Trp Asp Ile Ala Gly Arg
 1 5 10 15
 Phe Tyr Gly Asn Ser Thr Gln Trp Arg Lys Ile Trp Asn Ala Asn Lys
 20 25 30
 Thr Ala Met Ile Lys Arg Ser Lys Arg Asn Ile Arg Gln Pro Gly His
 35 40 45
 Trp Ile Phe Pro Gly Gln Lys Leu Lys Ile Pro Gln
 50 55 60

-continued

<210> SEQ ID NO 23
 <211> LENGTH: 60
 <212> TYPE: PRT
 <213> ORGANISM: Unknown
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Unknown Organism: Hypothetical
 wall-associated protein fragment

<400> SEQUENCE: 23

Thr Tyr Thr Val Lys Lys Gly Asp Thr Leu Trp Asp Leu Ala Gly Lys
 1 5 10 15
 Phe Tyr Gly Asp Ser Thr Lys Trp Arg Lys Ile Trp Lys Val Asn Lys
 20 25 30
 Lys Ala Met Ile Lys Arg Ser Lys Arg Asn Ile Arg Gln Pro Gly His
 35 40 45
 Trp Ile Phe Pro Gly Gln Lys Leu Lys Ile Pro Gln
 50 55 60

<210> SEQ ID NO 24
 <211> LENGTH: 167
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 24

Ala Pro Pro Val Glu Leu Ala Ala Asn Asp Leu Pro Ala Pro Leu Gly
 1 5 10 15
 Glu Pro Leu Pro Ala Ala Pro Ala Asp Pro Ala Pro Pro Ala Asp Leu
 20 25 30
 Ala Pro Pro Ala Pro Ala Asp Val Ala Pro Pro Val Glu Leu Ala Val
 35 40 45
 Asn Asp Leu Pro Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala
 50 55 60
 Asp Pro Ala Pro Pro Ala Asp Leu Ala Pro Pro Ala Pro Ala Asp Leu
 65 70 75 80
 Ala Pro Pro Ala Pro Ala Asp Leu Ala Pro Pro Ala Pro Ala Asp Leu
 85 90 95
 Ala Pro Pro Val Glu Leu Ala Val Asn Asp Leu Pro Ala Pro Leu Gly
 100 105 110
 Glu Pro Leu Pro Ala Ala Pro Ala Glu Leu Ala Pro Pro Ala Asp Leu
 115 120 125
 Ala Pro Ala Ser Ala Asp Leu Ala Pro Pro Ala Pro Ala Asp Leu Ala
 130 135 140
 Pro Pro Ala Pro Ala Glu Leu Ala Pro Pro Ala Pro Ala Asp Leu Ala
 145 150 155 160
 Pro Pro Ala Ala Val Asn Glu
 165

<210> SEQ ID NO 25
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 25

Ala Pro Pro Val Glu Leu Ala Ala Asn Asp Leu
 1 5 10

<210> SEQ ID NO 26
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

-continued

<400> SEQUENCE: 26

Ala Pro Pro Val Glu Leu Ala Val Asn Asp Leu
1 5 10

<210> SEQ ID NO 27

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 27

Pro Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala Glu Leu
1 5 10 15

<210> SEQ ID NO 28

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 28

Pro Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala Glu Leu
1 5 10 15

<210> SEQ ID NO 29

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 29

Pro Ala Pro Pro Ala Asp Leu
1 5

<210> SEQ ID NO 30

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 30

Ala Pro Pro Ala Pro Ala Asp Leu
1 5

<210> SEQ ID NO 31

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 31

Ala Pro Pro Ala Pro Ala Asp Val
1 5

<210> SEQ ID NO 32

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 32

Ala Pro Pro Ala Pro Ala Glu Leu
1 5

<210> SEQ ID NO 33

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 33

-continued

Ala Pro Pro Ala Pro Ala Glu Val
1 5

<210> SEQ ID NO 34
<211> LENGTH: 478
<212> TYPE: PRT
<213> ORGANISM: *Listeria monocytogenes*

<400> SEQUENCE: 34

Met Asn Met Lys Lys Ala Thr Ile Ala Ala Thr Ala Gly Ile Ala Val
1 5 10 15

Thr Ala Phe Ala Ala Pro Thr Ile Ala Ser Ala Ser Thr Val Val Val
20 25 30

Glu Ala Gly Asp Thr Leu Trp Gly Ile Ala Gln Ser Lys Gly Thr Thr
35 40 45

Val Asp Ala Ile Lys Lys Ala Asn Asn Leu Thr Thr Asp Lys Ile Val
50 55 60

Pro Gly Gln Lys Leu Gln Val Asn Asn Glu Val Ala Ala Ala Glu Lys
65 70 75 80

Thr Glu Lys Ser Val Ser Ala Thr Trp Leu Asn Val Arg Thr Gly Ala
85 90 95

Gly Val Asp Asn Ser Ile Ile Thr Ser Ile Lys Gly Gly Thr Lys Val
100 105 110

Thr Val Glu Thr Thr Glu Ser Asn Gly Trp His Lys Ile Thr Tyr Asn
115 120 125

Asp Gly Lys Thr Gly Phe Val Asn Gly Lys Tyr Leu Thr Asp Lys Ala
130 135 140

Val Ser Thr Pro Val Ala Pro Thr Gln Glu Val Lys Lys Glu Thr Thr
145 150 155 160

Thr Gln Gln Ala Ala Pro Val Ala Glu Thr Lys Thr Glu Val Lys Gln
165 170 175

Thr Thr Gln Ala Thr Thr Pro Ala Pro Lys Val Ala Glu Thr Lys Glu
180 185 190

Thr Pro Val Ile Asp Gln Asn Ala Thr Thr His Ala Val Lys Ser Gly
195 200 205

Asp Thr Ile Trp Ala Leu Ser Val Lys Tyr Gly Val Ser Val Gln Asp
210 215 220

Ile Met Ser Trp Asn Asn Leu Ser Ser Ser Ser Ile Tyr Val Gly Gln
225 230 235 240

Lys Leu Ala Ile Lys Gln Thr Ala Asn Thr Ala Thr Pro Lys Ala Glu
245 250 255

Val Lys Thr Glu Ala Pro Ala Ala Glu Lys Gln Ala Ala Pro Val Val
260 265 270

Lys Glu Asn Thr Asn Thr Asn Thr Ala Thr Thr Glu Lys Lys Glu Thr
275 280 285

Ala Thr Gln Gln Gln Thr Ala Pro Lys Ala Pro Thr Glu Ala Ala Lys
290 295 300

Pro Ala Pro Ala Pro Ser Thr Asn Thr Asn Ala Asn Lys Thr Asn Thr
305 310 315 320

Asn Thr Asn Thr Asn Asn Thr Asn Thr Pro Ser Lys Asn Thr Asn Thr
325 330 335

Asn Ser Asn Thr Asn Thr Asn Thr Asn Ser Asn Thr Asn Ala Asn Gln
340 345 350

Gly Ser Ser Asn Asn Asn Ser Asn Ser Ser Ala Ser Ala Ile Ile Ala
355 360 365

-continued

Glu Ala Gln Lys His Leu Gly Lys Ala Tyr Ser Trp Gly Gly Asn Gly
 370 375 380

Pro Thr Thr Phe Asp Cys Ser Gly Tyr Thr Lys Tyr Val Phe Ala Lys
 385 390 395 400

Ala Gly Ile Ser Leu Pro Arg Thr Ser Gly Ala Gln Tyr Ala Ser Thr
 405 410 415

Thr Arg Ile Ser Glu Ser Gln Ala Lys Pro Gly Asp Leu Val Phe Phe
 420 425 430

Asp Tyr Gly Ser Gly Ile Ser His Val Gly Ile Tyr Val Gly Asn Gly
 435 440 445

Gln Met Ile Asn Ala Gln Asp Asn Gly Val Lys Tyr Asp Asn Ile His
 450 455 460

Gly Ser Gly Trp Gly Lys Tyr Leu Val Gly Phe Gly Arg Val
 465 470 475

<210> SEQ ID NO 35
 <211> LENGTH: 758
 <212> TYPE: DNA
 <213> ORGANISM: Micrococcus luteus
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (66)..(728)

<400> SEQUENCE: 35

accaaggaga aggacgaccc cggtgtgcct cggccgcccga tcagcgagga ctgcgccatgg 60

acacc atg act ctc ttc acc act tcc gcc acc cgc tcc cgc cgt gcc acc 110
 Met Thr Leu Phe Thr Thr Ser Ala Thr Arg Ser Arg Arg Ala Thr
 1 5 10 15

gcc tcg atc gtc gcg ggc atg acc ctc gcc ggc gcc gcc gcc gtg ggc 158
 Ala Ser Ile Val Ala Gly Met Thr Leu Ala Gly Ala Ala Ala Val Gly
 20 25 30

ttc tcc gcc ccg gcc cag gcc gcc acc gtg gac acc tgg gac cgc ctc 206
 Phe Ser Ala Pro Ala Gln Ala Ala Thr Val Asp Thr Trp Asp Arg Leu
 35 40 45

gcc gag tgc gag tcc aac ggc acc tgg gac atc aac acc ggc aac ggc 254
 Ala Glu Cys Glu Ser Asn Gly Thr Trp Asp Ile Asn Thr Gly Asn Gly
 50 55 60

ttc tac ggc ggc gtg cag ttc acc ctg tcc tcc tgg cag gcc gtc ggc 302
 Phe Tyr Gly Gly Val Gln Phe Thr Leu Ser Ser Trp Gln Ala Val Gly
 65 70 75

ggc gaa ggc tac ccg cac cag gcc tcg aag gcc gag cag atc aag cgc 350
 Gly Glu Gly Tyr Pro His Gln Ala Ser Lys Ala Glu Gln Ile Lys Arg
 80 85 90 95

gcc gag atc ctc cag gac ctg cag ggc tgg ggc gcg tgg ccg ctg tgc 398
 Ala Glu Ile Leu Gln Asp Leu Gln Gly Trp Gly Ala Trp Pro Leu Cys
 100 105 110

tcg cag aag ctg ggc ctg acc cag gct gac gcg gac gcc ggt gac gtg 446
 Ser Gln Lys Leu Gly Leu Thr Gln Ala Asp Ala Asp Ala Gly Asp Val
 115 120 125

gac gcc acc gag gcc gcc ccg gtc gcc gtg gag cgc acg gcc acc gtg 494
 Asp Ala Thr Glu Ala Ala Pro Val Ala Val Glu Arg Thr Ala Thr Val
 130 135 140

cag cgc cag tcc gcc gcg gac gag gct gcc gcc gag cag gcc gct gcc 542
 Gln Arg Gln Ser Ala Ala Asp Glu Ala Ala Ala Gln Ala Ala Ala
 145 150 155

gcg gag cag gcc gtc gtc gcc gag gcc gag acc atc gtc gtc aag tcc 590
 Ala Glu Gln Ala Val Val Ala Glu Ala Glu Thr Ile Val Val Lys Ser
 160 165 170 175

ggg gac tcc ctc tgg acg ctc gcc aac gag tac gag gtg gag ggt ggc 638
 Gly Asp Ser Leu Trp Thr Leu Ala Asn Glu Tyr Glu Val Glu Gly Gly

-continued

180	185	190	
tgg acc gcc ctc tac gag gcc aac aag ggc gcc gtc tcc gac gcc gcc			686
Trp Thr Ala Leu Tyr Glu Ala Asn Lys Gly Ala Val Ser Asp Ala Ala			
195	200	205	
gtg atc tac gtc ggc cag gag ctc gtc ctg ccg cag gcc tga			728
Val Ile Tyr Val Gly Gln Glu Leu Val Leu Pro Gln Ala			
210	215	220	
gacgcctgac cggccccccg gaccggtacc			758

<210> SEQ ID NO 36
 <211> LENGTH: 220
 <212> TYPE: PRT
 <213> ORGANISM: Micrococcus luteus

<400> SEQUENCE: 36

Met Thr Leu Phe Thr Thr Ser Ala Thr Arg Ser Arg Arg Ala Thr Ala		
1	5	10
Ser Ile Val Ala Gly Met Thr Leu Ala Gly Ala Ala Ala Val Gly Phe		
20	25	30
Ser Ala Pro Ala Gln Ala Ala Thr Val Asp Thr Trp Asp Arg Leu Ala		
35	40	45
Glu Cys Glu Ser Asn Gly Thr Trp Asp Ile Asn Thr Gly Asn Gly Phe		
50	55	60
Tyr Gly Gly Val Gln Phe Thr Leu Ser Ser Trp Gln Ala Val Gly Gly		
65	70	75
Glu Gly Tyr Pro His Gln Ala Ser Lys Ala Glu Gln Ile Lys Arg Ala		
85	90	95
Glu Ile Leu Gln Asp Leu Gln Gly Trp Gly Ala Trp Pro Leu Cys Ser		
100	105	110
Gln Lys Leu Gly Leu Thr Gln Ala Asp Ala Asp Ala Gly Asp Val Asp		
115	120	125
Ala Thr Glu Ala Ala Pro Val Ala Val Glu Arg Thr Ala Thr Val Gln		
130	135	140
Arg Gln Ser Ala Ala Asp Glu Ala Ala Ala Glu Gln Ala Ala Ala Ala		
145	150	155
Glu Gln Ala Val Val Ala Glu Ala Glu Thr Ile Val Val Lys Ser Gly		
165	170	175
Asp Ser Leu Trp Thr Leu Ala Asn Glu Tyr Glu Val Glu Gly Gly Trp		
180	185	190
Thr Ala Leu Tyr Glu Ala Asn Lys Gly Ala Val Ser Asp Ala Ala Val		
195	200	205
Ile Tyr Val Gly Gln Glu Leu Val Leu Pro Gln Ala		
210	215	220

<210> SEQ ID NO 37
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 37

gcsacsgtsg acacstggga ccgscstsgcs gag	33
---------------------------------------	----

<210> SEQ ID NO 38
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Micrococcus luteus

-continued

```

<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (13)..(13)
<223> OTHER INFORMATION: Variable amino acid
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (18)..(18)
<223> OTHER INFORMATION: Variable amino acid

<400> SEQUENCE: 38

Ala Thr Val Asp Thr Trp Asp Arg Leu Ala Glu Glu Xaa Ser Asn Gly
1           5           10           15

Thr Xaa Asp

<210> SEQ ID NO 39
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

<400> SEQUENCE: 39

ccgccgtaga agccgttg                               18

<210> SEQ ID NO 40
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

<400> SEQUENCE: 40

agttcacccct gtctctctg                               19

<210> SEQ ID NO 41
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (9)..(9)
<223> OTHER INFORMATION: i
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (15)..(15)
<223> OTHER INFORMATION: i
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)..(21)
<223> OTHER INFORMATION: i

<400> SEQUENCE: 41

gcytgrtgng grtanccytc ncc                               23

<210> SEQ ID NO 42
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Micrococcus luteus

<400> SEQUENCE: 42

Val Gly Gly Glu Gly Tyr Pro His Gln Ala Ser Lys
1           5           10

<210> SEQ ID NO 43

```

-continued

<211> LENGTH: 182
 <212> TYPE: PRT
 <213> ORGANISM: *Micrococcus luteus*

<400> SEQUENCE: 43

Ala Thr Val Asp Thr Trp Asp Arg Leu Ala Glu Cys Glu Ser Asn Gly
 1 5 10 15
 Thr Trp Asp Ile Asn Thr Gly Asn Gly Phe Tyr Gly Gly Val Gln Phe
 20 25 30
 Thr Leu Ser Ser Trp Gln Ala Val Gly Gly Glu Gly Tyr Pro His Gln
 35 40 45
 Ala Ser Lys Ala Glu Gln Ile Lys Arg Ala Glu Ile Leu Gln Asp Leu
 50 55 60
 Gln Gly Trp Gly Ala Trp Pro Leu Cys Ser Gln Lys Leu Gly Leu Thr
 65 70 75 80
 Gln Ala Asp Ala Asp Ala Gly Asp Val Asp Ala Thr Glu Ala Ala Pro
 85 90 95
 Val Ala Val Glu Arg Thr Ala Thr Val Gln Arg Gln Ser Ala Ala Asp
 100 105 110
 Glu Ala Ala Ala Glu Gln Ala Ala Ala Glu Gln Ala Val Val Ala
 115 120 125
 Glu Ala Glu Thr Ile Val Val Lys Ser Gly Asp Ser Leu Trp Thr Leu
 130 135 140
 Ala Asn Glu Tyr Glu Val Glu Gly Gly Trp Thr Ala Leu Tyr Glu Ala
 145 150 155 160
 Asn Lys Gly Ala Val Ser Asp Ala Ala Val Ile Tyr Val Gly Gln Glu
 165 170 175
 Leu Val Leu Pro Gln Ala
 180

<210> SEQ ID NO 44
 <211> LENGTH: 299
 <212> TYPE: DNA
 <213> ORGANISM: *Streptomyces coelicolor*
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (3)..(299)

<400> SEQUENCE: 44

gg atc cgc acc gcc gcg gta acc ctg gtc gcc gcg acc gca ctc ggg 47
 Ile Arg Thr Ala Ala Val Thr Leu Val Ala Ala Thr Ala Leu Gly
 1 5 10 15
 gcg acc gcc gaa gcg gtg gcc gcg ccc tcg gcg ccc ctg cgc acc gac 95
 Ala Thr Gly Glu Ala Val Ala Ala Pro Ser Ala Pro Leu Arg Thr Asp
 20 25 30
 tgg gac gcc atc gcc gcg tgc gag tcc agc ggc aac tgg cag gcg aac 143
 Trp Asp Ala Ile Ala Ala Cys Glu Ser Ser Gly Asn Trp Gln Ala Asn
 35 40 45
 acc gcc aac gcc tac tac gcc gcc ctg cag ttc gca cgg tcc agc tgg 191
 Thr Gly Asn Gly Tyr Tyr Gly Gly Leu Gln Phe Ala Arg Ser Ser Trp
 50 55 60
 atc gcc gcc gcc gcc ctc aag tac gcc ccg cgc gcg gac ctc gcc acc 239
 Ile Ala Ala Gly Gly Leu Lys Tyr Ala Pro Arg Ala Asp Leu Ala Thr
 65 70 75
 cgc gcc gag cag atc gcc gtg gcg gaa cgc ctc gcc cgt ctg cag ggg 287
 Arg Gly Glu Gln Ile Ala Val Ala Glu Arg Leu Ala Arg Leu Gln Gly
 80 85 90 95
 atg tcc gcc tgg 299
 Met Ser Ala Trp

-continued

<210> SEQ ID NO 45
 <211> LENGTH: 99
 <212> TYPE: PRT
 <213> ORGANISM: Streptomyces coelicolor

<400> SEQUENCE: 45

```
Ile Arg Thr Ala Ala Val Thr Leu Val Ala Ala Thr Ala Leu Gly Ala
1          5          10          15
Thr Gly Glu Ala Val Ala Ala Pro Ser Ala Pro Leu Arg Thr Asp Trp
          20          25          30
Asp Ala Ile Ala Ala Cys Glu Ser Ser Gly Asn Trp Gln Ala Asn Thr
          35          40          45
Gly Asn Gly Tyr Tyr Gly Gly Leu Gln Phe Ala Arg Ser Ser Trp Ile
          50          55          60
Ala Ala Gly Gly Leu Lys Tyr Ala Pro Arg Ala Asp Leu Ala Thr Arg
65          70          75          80
Gly Glu Gln Ile Ala Val Ala Glu Arg Leu Ala Arg Leu Gln Gly Met
          85          90          95
Ser Ala Trp
```

<210> SEQ ID NO 46
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 46

gtcagaattc atatggccac cgtggacacc tggg 34

<210> SEQ ID NO 47
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 47

tgacggatcc tattaggcct gcggcaggac gag 33

<210> SEQ ID NO 48
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 48

atcagaattc atatggacga catcgattgg gacgc 35

<210> SEQ ID NO 49
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 49

cgcaggatcc cctcaatcgt ccctgctcc 29

<210> SEQ ID NO 50
 <211> LENGTH: 23
 <212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 50

gaagagaatt ccttccatca cga                                     23

<210> SEQ ID NO 51
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 51

ccaaacgaat tcggtcaatc ac                                     22

<210> SEQ ID NO 52
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 52

gcaaggatcc cagactaaaa aaacag                                 26

<210> SEQ ID NO 53
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer

<400> SEQUENCE: 53

atcaggatcc atattattag tttaaga                                27

<210> SEQ ID NO 54
<211> LENGTH: 663
<212> TYPE: DNA
<213> ORGANISM: Micrococcus luteus
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(663)

<400> SEQUENCE: 54

atg act ctc ttc acc act tcc gcc acc cgc tcc cgc cgt gcc acc gcc      48
Met Thr Leu Phe Thr Thr Ser Ala Thr Arg Ser Arg Arg Ala Thr Ala
1      5      10      15

tcg atc gtc gcg ggc atg acc ctc gcc ggc gcc gcc gcc gtg ggc ttc      96
Ser Ile Val Ala Gly Met Thr Leu Ala Gly Ala Ala Ala Val Gly Phe
20     25     30

tcc gcc ccg gcc cag gcc gcc acc gtg gac acc tgg gac cgc ctc gcc      144
Ser Ala Pro Ala Gln Ala Ala Thr Val Asp Thr Trp Asp Arg Leu Ala
35     40     45

gag tgc gag tcc aac ggc acc tgg gac atc aac acc ggc aac ggc ttc      192
Glu Cys Glu Ser Asn Gly Thr Trp Asp Ile Asn Thr Gly Asn Gly Phe
50     55     60

tac ggc ggc gtg cag ttc acc ctg tcc tcc tgg cag gcc gtc ggc ggc      240
Tyr Gly Gly Val Gln Phe Thr Leu Ser Ser Trp Gln Ala Val Gly Gly
65     70     75     80

gaa ggc tac ccg cac cag gcc tcg aag gcc gag cag atc aag cgc gcc      288
Glu Gly Tyr Pro His Gln Ala Ser Lys Ala Glu Gln Ile Lys Arg Ala
85     90     95

gag atc ctc cag gac ctg cag ggc tgg ggc gcg tgg ccg ctg tgc tcg      336

```

-continued

Glu	Ile	Leu	Gln	Asp	Leu	Gln	Gly	Trp	Gly	Ala	Trp	Pro	Leu	Cys	Ser		
			100					105					110				
cag	aag	ctg	ggc	ctg	acc	cag	gct	gac	gcg	gac	gcc	ggt	gac	gtg	gac	384	
Gln	Lys	Leu	Gly	Leu	Thr	Gln	Ala	Asp	Ala	Asp	Ala	Gly	Asp	Val	Asp		
		115					120					125					
gcc	acc	gag	gcc	gcc	ccg	gtc	gcc	gtg	gag	cgc	acg	gcc	acc	gtg	cag	432	
Ala	Thr	Glu	Ala	Ala	Pro	Val	Ala	Val	Glu	Arg	Thr	Ala	Thr	Val	Gln		
		130					135					140					
cgc	cag	tcc	gcc	gcg	gac	gag	gct	gcc	gcc	gag	cag	gcc	gct	gcc	gcg	480	
Arg	Gln	Ser	Ala	Ala	Asp	Glu	Ala	Ala	Ala	Glu	Gln	Ala	Ala	Ala	Ala		
		145			150					155				160			
gag	cag	gcc	gtc	gtc	gcc	gag	gcc	gag	acc	atc	gtc	gtc	aag	tcc	ggt	528	
Glu	Gln	Ala	Val	Val	Ala	Glu	Ala	Glu	Thr	Ile	Val	Val	Lys	Ser	Gly		
			165						170					175			
gac	tcc	ctc	tgg	acg	ctc	gcc	aac	gag	tac	gag	gtg	gag	ggt	ggc	tgg	576	
Asp	Ser	Leu	Trp	Thr	Leu	Ala	Asn	Glu	Tyr	Glu	Val	Glu	Gly	Gly	Trp		
		180						185					190				
acc	gcc	ctc	tac	gag	gcc	aac	aag	ggc	gcc	gtc	tcc	gac	gcc	gcc	gtg	624	
Thr	Ala	Leu	Tyr	Glu	Ala	Asn	Lys	Gly	Ala	Val	Ser	Asp	Ala	Ala	Val		
		195					200					205					
atc	tac	gtc	ggc	cag	gag	ctc	gtc	ctg	ccg	cag	gcc	tga				663	
Ile	Tyr	Val	Gly	Gln	Glu	Leu	Val	Leu	Pro	Gln	Ala						
		210				215					220						

<210> SEQ ID NO 55
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 55

Ala Pro Pro Ala Asp Leu
 1 5

<210> SEQ ID NO 56
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 56

Ala Pro Ala Ser Ala Asp Leu
 1 5

<210> SEQ ID NO 57
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 57

Ala Pro Pro Ala Pro Ala Glu Leu
 1 5

<210> SEQ ID NO 58
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 58

Ala Pro Pro Ala
 1

<210> SEQ ID NO 59
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Mycobacterium tuberculosis

-continued

<400> SEQUENCE: 59

Ala Val Asn Glu
1<210> SEQ ID NO 60
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mycobacterium tuberculosis
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (14)..(14)
<223> OTHER INFORMATION: Asp or Glu

<400> SEQUENCE: 60

Pro Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala Xaa Leu
1 5 10 15<210> SEQ ID NO 61
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mycobacterium tuberculosis
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Asp or Glu
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Leu or Val

<400> SEQUENCE: 61

Ala Pro Pro Ala Pro Ala Xaa Xaa
1 5<210> SEQ ID NO 62
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mycobacterium tuberculosis
<220> FEATURE:
<221> NAME/KEY: MOD_RES
<222> LOCATION: (8)..(8)
<223> OTHER INFORMATION: Ala or Val

<400> SEQUENCE: 62

Ala Pro Pro Val Glu Leu Ala Xaa Asn Asp Leu
1 5 10<210> SEQ ID NO 63
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Mycobacterium tuberculosis

<400> SEQUENCE: 63

Pro Ala Pro Leu Gly Glu Pro Leu Pro Ala Ala Pro Ala Asp
1 5 10<210> SEQ ID NO 64
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Micrococcus luteus

<400> SEQUENCE: 64

Lys Ala Glu Gln Ile Lys Arg Ala Glu
1 5<210> SEQ ID NO 65
<211> LENGTH: 5

-continued

<212> TYPE: PRT
 <213> ORGANISM: *Micrococcus luteus*

<400> SEQUENCE: 65

Asp Thr Trp Asp Arg
 1 5

The invention claimed is:

1. A composition comprising a polypeptide purified to essential homogeneity, wherein the polypeptide is selected from the group consisting of:

- i) a polypeptide having at least 95% sequence identity with amino acid residues 285 to 355 of SEQ ID NO: 1, amino acid residues 117 to 184 of SEQ ID NO: 2, amino acid residues 44 to 114 of SEQ ID NO: 4, amino acid residues 73 to 140 of SEQ ID NO: 6, amino acid residues 54 to 121 of SEQ ID NO: 7 or amino acid residues 224 to 318 of SEQ ID NO: 11;
- ii) a polypeptide having at least 95% sequence identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 36 or SEQ ID NO: 43;
- iii) a polypeptide comprising at least amino acid residues 285 to 355 of SEQ ID NO: 1, amino acid residues 117 to 184 of SEQ ID NO: 2, amino acid residues 44 to 114 of SEQ ID NO: 4, amino acid residues 73 to 140 of SEQ ID NO: 6, amino acid residues 54 to 121 of SEQ ID NO: 7 or amino acid residues 224 to 318 of SEQ ID NO: 11; and
- iv) a polypeptide comprising the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 36 or SEQ ID NO: 43.

2. The composition of claim 1, wherein the composition is in a unit dosage form or a form suitable for local or systemic administration.

3. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 1.

4. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

5. A kit comprising a polypeptide purified to essential homogeneity, wherein the polypeptide is selected from the group consisting of:

- i) a polypeptide having at least 95% sequence identity with amino acid residues 285 to 355 of SEQ ID NO: 1, amino acid residues 117 to 184 of SEQ ID NO: 2, amino acid residues 44 to 114 of SEQ ID NO: 4, amino acid residues 73 to 140 of SEQ ID NO: 6, amino acid residues 54 to 121 of SEQ ID NO: 7 or amino acid residues 224 to 318 of SEQ ID NO: 11;
- ii) a polypeptide having at least 95% sequence identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 36 or SEQ ID NO: 43;
- iii) a polypeptide comprising at least amino acid residues 285 to 355 of SEQ ID NO: 1, amino acid residues 117 to 184 of SEQ ID NO: 2, amino acid residues 44 to 114 of SEQ ID NO: 4, amino acid residues 73 to 140 of SEQ ID NO: 6, amino acid residues 54 to 121 of SEQ ID NO: 7 or amino acid residues 224 to 318 of SEQ ID NO: 11; and
- iv) a polypeptide comprising the amino acid sequence of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 11, SEQ ID NO: 36 or SEQ ID NO: 43; and

instructions for use.

6. The kit of claim 5, wherein the polypeptide is present in a composition comprising a pharmaceutical excipient.

7. The kit of claim 5, wherein the composition is in a unit dosage form or a form suitable for local or systemic administration.

8. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 1.

9. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

10. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 2.

11. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 4.

12. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 6.

13. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 7.

14. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 11.

15. The composition of claim 1, wherein the isolated polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 43.

16. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 2.

17. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 4.

18. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 6.

19. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 7.

20. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 11.

21. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 43.

22. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 2.

23. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 4.

24. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 6.

25. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 7.

26. The kit of claim 5, wherein the isolated polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 11.

79

27. The kit of claim 5, wherein the isolated polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 43.

28. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 2.

29. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID: 4.

30. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 6.

31. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 7.

32. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 11.

33. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 43.

80

34. The composition of claim 1, further comprising a pharmaceutical excipient.

35. The composition of claim 1, wherein the polypeptide has at least 95% sequence identity with the amino acid sequence of SEQ ID NO: 36.

36. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 36.

37. The kit of claim 5, wherein the polypeptide has at least 95% sequence identity the amino acid-sequence of SEQ ID NO: 36.

38. The kit of claim 5, wherein the polypeptide comprises the amino acid sequence of SEQ ID NO: 36.

* * * * *