

A flavour of omics approaches for the detection of food fraud

David I Ellis¹, Howbeer Muhamadali^{1,2}, David P Allen¹,
Christopher T Elliott² and Royston Goodacre¹



Food fraud has been identified as an increasing problem on a global scale with wide-ranging economic, social, health and environmental impacts. Omics and their related techniques, approaches, and bioanalytical platforms incorporate a significant number of scientific areas which have the potential to be applied to and significantly reduce food fraud and its negative impacts. In this overview we consider a selected number of very recent studies where omics techniques were applied to detect food authenticity and could be implemented to ensure food integrity. We postulate that significant reductions in food fraud, with the assistance of omics technologies and other approaches, will result in less food waste, decreases in energy use as well as greenhouse gas emissions, and as a direct consequence of this, increases in quality, productivity, yields, and the ability of food systems to be more resilient and able to withstand future food shocks.

Addresses

¹ Manchester Institute of Biotechnology, School of Chemistry, University of Manchester, 131 Princess Street, Manchester M1 7ND, UK

² Queen's University Belfast, School of Biological Sciences, Institute for Global Food Security, Belfast BT9 5AG, County Antrim, Northern Ireland, UK

Corresponding author: Ellis, David I (D.Ellis@manchester.ac.uk)

Current Opinion in Food Science 2016, 10:7–15

This review comes from a themed issue on **Innovations in Food Science**

Edited by **Daniel Cozzolino**

<http://dx.doi.org/10.1016/j.cofs.2016.07.002>

2214-7993/© 2016 Elsevier Ltd. All rights reserved.

Introduction

In the spring of 2016 the latest report from Operation Opson (Opson V) was released detailing the largest ever seizures of hazardous fake food and drink ever recorded [1]. This joint international Interpol/Europol operation originally began in 2011 and initially included 10 European countries, with it now expanding to include 57 countries across the world. Perhaps not surprisingly the release of the report led to a slew of media headlines involving terms such as monkey meat, copper sulfate painted olives,

fertiliser contaminated sugar, and tonnes of locusts and caterpillars seized, to name but a few; alarmist headlines perhaps, but all of them true (see [Table 1](#) for a summary of Opson V seizures). It should also be pointed out that these large-scale and *record seizures* of fake and counterfeit foods and beverages, carried out at shops, markets, industrial estates, air- and seaports, all occurred during a relatively short period from November 2015 to February 2016, and are only a snapshot of the severity of the problem. The news of Opson V appeared to coincide with the release of the first report [2] from the UK's newly formed National Food Crime Unit (NFCU) which covered the period November 2014 to July 2015. Whilst informative, the only seizures mentioned within the NFCU report involved counterfeit and adulterated alcohol, including 35 000 bottles of counterfeit vodka originating from Ukraine, and 8 000 litres of vodka from Lithuania with forged duty stamps. More worryingly, these seizures also included 20 000 counterfeit branded vodka bottles and material suggesting adulteration with anti-freeze, as well as 130 000 litres of potentially toxic spirits alongside bottling and labelling materials from another raid [2]. More recently, reports of organised international food fraud involved seven countries and thousands of tonnes of wheat, corn, soybeans, rapeseed and sunflower seed imported from multiple non-EU countries, mislabelled as organic and shipped to EU countries via Malta or Italy [3]. These reports readily illustrate and reinforce the fact that food fraud can be international in scope, with no country being immune from its reach and impacts, and that this transboundary criminal activity can be both opportunistic, as well as highly organised. Here though, as analytical scientists, we are primarily concerned with the single greatest, or what could be termed *grand challenge* of food adulteration; its unequivocal detection. Therefore, we have selected a very small number of recently reported omics and related technologies that are being developed to enable the detection of food authenticity and integrity.

Omics technologies

During the past two decades molecular-based technologies have rightly proven themselves as an invaluable option for the detection of food authenticity and integrity [4]. Such DNA-based methodologies generally rely on specific DNA sequences (markers) that can be used for detection of food adulterants and/or approving the authenticity (i.e. quality and origin) of raw ingredients [5].

Table 1

A summary of seizures of fake food and drink undertaken between November 2015 and February 2016 from Opson V

Country	Summary of seizures
Australia	450 kg of honey found to be blended or adulterated. Peanuts repackaged and relabeled as pine nuts.
Belgium	Several kilos of monkey meat found at Zaventem airport.
Bolivia	Warehouse containing thousands of cans of sardines, and fake labels from a famous Peruvian brand of sardines.
Burundi	36 000 liters of illicit alcohol seized during the operation, as well as nine Kalashnikov rifles, ammunition, and three grenades
France	11 kg of locusts and 20 kg of caterpillars seized and destroyed.
Greece	Three illicit factories producing counterfeit alcohol. Equipment, fake labels, caps, empty bottles and 7 400 bottles of fake alcohol seized.
Hungary	Counterfeit non-alcoholic sparkling wine, chocolates, sweets.*
India	See Thailand below.
Indonesia	70 tonnes of chicken intestines preserved in formalin seized. 310 000 illegal food products found behind piles of tiles in a warehouse and believed to be smuggled from Malaysia.
Italy	85 tonnes of olives seized, painted with copper sulfate solution to enhance colour. Counterfeit non-alcoholic sparkling wine, chocolates, sweets.*
Lithuania	Counterfeit non-alcoholic sparkling wine, chocolates, sweets.*
Malaysia	See Indonesia above.
Romania	Counterfeit non-alcoholic sparkling wine, chocolates, sweets.*
South Korea	Arrest made associated with the online sale of fake dietary supplements/weight loss products estimated to have generated US\$170 000 in a 10 month period.
Sudan	9 tonnes of counterfeit sugar contaminated with fertilizer.
Thailand	Four tonnes of meat smuggled by one individual from India. Further investigation led to discovery of illicit network operating across 10 provinces. Recovery and destruction of more than 30 tonnes of illegal beef and buffalo meat, unfit for human consumption and destined for sale in supermarkets.
Togo	24 tonnes of imported tilapia.
United Kingdom	10 000 litres of fake or adulterated alcohol, including wine, whisky and vodka.
Zambia	1 300 bottles of fake whisky in original packaging. Over 3 200 cartons of diet powder drinks with modified expiration dates.

* Aimed at children and destined for export to West Africa.

DNA-barcoding, named due to this technique using a specific region of the genome described as the DNA-barcode, is considered as one the most common identification systems for taxonomic discrimination [5]. However, the successful application of this approach for the separation of food and foodstuffs relies on the availability of comprehensive reference sequence libraries, such as the barcode of life database (www.barcodeoflife.org).

DNA-barcoding is of particular interest when it comes to authentication of seafood products [6]. This interest is mainly due to the presence of a wide-range of species, morphological similarities between species, as well as a loss of the structural and visible characteristics of the raw material during different food processing procedures (i.e. heat treatment, or cooking). Several studies have successfully applied this approach for seafood authenticity testing, such as those by Cutarelli and co-workers, who reported the application of mitochondrial cytochrome b (Cytb) and cytochrome oxidase subunit I (COI) genes as DNA markers for the identification of 58 Mediterranean marine fish samples sold on the Italian market. Whilst Pereira *et al.* demonstrated the efficacy of barcode methodology (COI gene), for the highly accurate (99.2%) identification of 254 species of freshwater fish samples, Kim and co-workers picked up the metaphorical baton, taking this approach a step further by employing a combination of DNA-barcoding and stable isotope analysis for

the identification and verification of the origin of Hairtail fish and shrimp. This strategy also allowed these authors to differentiate between natural and farmed shrimp [7], an important and significant area for fish authenticity and traceability, which was only possible as the stable isotope analyses allowed the phenotype of the organism to be measured.

As might be expected, DNA-based methodologies have been applied to the authentication and traceability of a wide-range of food products, including the detection of the mislabelling or cross-contamination of halal meat [8] and the detection of species such as horse in ground meats [9]. With one very recent article of significant interest involving the development of a real-time PCR approach for the relative quantitation of horse DNA in raw beef mixtures [10**]. Other studies have involved labelled milk and milk products, such as yogurt and cheese, which were traced through a production chain via DNA tags, in this case silica particles with encapsulated DNA (SPED) [11], with the applicability of synthetic and naturally occurring DNA sequences demonstrated. Identifying species specific differences in herbal medicines [12], chilli adulteration of traded black pepper powder down to 0.5% adulteration [13], and tracing/tagging of edible oils (e.g. olive oil) using encapsulated DNA in heat-resistant and inert magnetic particles [14] have also been reported. However, the application of DNA-barcoding

using the more traditional sequencing methods, such as Sanger sequencing, can suffer from some drawbacks including their low-throughput nature and the necessity for food samples with high DNA purity and concentration [15]. That said, such inherent limitations have been largely overcome using more recent technologies such as next generation sequencing (NGS), which present themselves as high-throughput and low cost approaches in comparison to earlier genomics methods, and allow for the sequencing of millions of DNA molecules in parallel [16]. These include microfluidic and nanofluidic devices such as the nanopore [17]. Readers with a particular interest in genomic methods for detecting food authenticity and integrity are directed to far more comprehensive reviews in this area [4,5,12,16].

The applications of analytical platforms common within other omics approaches have also been proven valuable in providing critical information regarding the biochemical composition of various food products. Proteomics, whilst a well-established and continually developing field in many areas of research, has been said to be emerging as a complementary methodology [18] to the DNA-based approaches (as well as antibody approaches [19,20]), for food authenticity detection, as the amino acid sequence, just like the DNA sequence, is species specific. Given the recent resurgence in public interest in food adulteration following the horsemeat scandal in the UK, it is perhaps to be expected that several of the recent proteomics approaches have involved adulteration of meat products. These have included the identification of novel heat stable peptides for horsemeat speciation in highly processed foods such as corned beef and baby food. After in-depth analyses of these two well-known products, muscle food mixtures of various meat species such as horse, cow, pig, and chicken were used to emulate these complex processed food matrices. This semi-targeted approach then utilised nanoflow liquid chromatography mass spectrometry (nLC–MS/MS) to produce a database of markers for a total of nine animal species [18], and was able to detect levels of processed and raw horse meats in meat mixtures as low as 0.5% (w/w).

Other analysts have used high-performance liquid chromatography–mass spectrometry (HPLC–MS) analysis of tryptic digests of protein extracts from a number of species, to detect horsemeat and pork in a range of these muscle foods, including halal beef. They developed multiple reaction monitoring (MRM) methods which allow for signal enhancement and the detection of 0.55% adulteration/contamination of horse or pork in a beef matrix, with still further enhancements in sensitivity (referred to as MRM³) allowing for the detection of 0.13% pork in halal beef [21].

More recently Obana and co-workers used LC–MS/MS shotgun spectral matching for the speciation as well as

quantification of a wide-range (16 mammalian and 10 avian species) of raw, cooked, and mixed meat types [22]. While others have recently trod a similar path of proteomic meat speciation and quantification but with less success in terms of the limit of detection (LOD) than the studies already discussed above, with targeted levels of meat speciation down to 1% (w/w) [23]. Of course proteomic detection methods involve many food products in addition to meats. Milk has been one of these products for example, where matrix-assisted laser desorption/ionisation mass spectrometry (MALDI-MS) methods have been used for the combined peptidomic and proteomic profiling of raw and thermally treated cow, goat, buffalo and sheep milk and combinations of these with skimmed milks and water at various concentrations. Results were said to permit rapid and accurate estimates of the extent of fraud at either the protein or peptide level, and were also said to extend previous MALDI-MS approaches in this area via the analysis of heat treatment and complementary peptide profiling, thereby potentially broadening the applicability of these methods [24].

There have also been a number of recent reviews in the field of proteomics for food fraud and related areas, including topics such as: biomarker discovery and applications for foods and beverages [25]; proteomics as a tool to understand the complexity of beers [26]; fruits and beverages [27]; and yet again, meat and muscle for animal origin authentication [28], as well as MS-based omics strategies concerning the analysis of the complex links between food science and nutrition, referred to as foodomics [29].

The more recent omics disciplines are of course metabolomics and lipidomics, the latter so relatively recent that very few standalone lipidomic studies of food fraud or authenticity have been reported to date (with the odd exception [30]). These do however include Fourier transform infrared (FT-IR) spectroscopy as a tool to authenticate the origin of wild and farmed European sea bass, as well as the additional benefit of providing useful information on the composition of nutritionally beneficial lipids [31]. That being said, there are additional lipidomics food authenticity methods that have been integrated within metabolomics studies, and we would expect this, as the lipidome is a subset of the more chemically diverse metabolome. Metabolomics is a rapidly expanding field within the omics finding useful and growing applications across a very broad range of disciplines. Metabolomics is downstream of genomics and proteomics (see Table 2 for a description of the omics approaches) and said to be the link between genotype and phenotype. Thus, it is not surprising that there are significantly more metabolomics studies related to food authenticity and integrity to be found during the last few years, than those from the other omics technologies

Table 2

A brief definition and description of the omics technologies used for the detection of food authenticity and integrity

Omics approaches	Description	Approaches and technologies
Genomics	The study and assessment of variability and function of DNA sequences	Whole-genome sequencing (WGS), Next generation sequencing (NGS), Amplified fragment length polymorphism (AFLP), Single nucleotide polymorphism (SNP), Simple sequence repeat (SSR)
Proteomics	The study of structure, function and abundance of different proteins and peptides (or complexes), post-translational modifications (PTM) in a system	Two-dimensional gel electrophoresis, MALDI-TOF-MS, LC-MS and high-resolution tandem mass spectrometry
Metabolomics	The comprehensive and systematic study of low molecular weight compounds (metabolites) involved in metabolism in a system	Metabolic fingerprinting (FT-IR and Raman spectroscopies), Metabolic profiling (GC/LC-MS, NMR), Targeted metabolomics analysis
Lipidomics	The study of pathways and networks of different lipid species in a system	Lipid profiling (GC/LC-MS, NMR), shot-gun MS, lipid fingerprinting (MALDI-TOF-MS), ultra-performance LC-MS (UPLC-MS)

mentioned here, especially as metabolites are often essential for human nutrition.

Indeed, these recent metabolomics studies appear to encompass not only a wider range of technological methods but also a more diverse range of applications within this area. These include nuclear magnetic resonance (NMR) spectroscopy for the determination of the country of origin of coffee by Arana and co-workers [32], using a fingerprinting approach [33] and showing high classification rates of large numbers of spectra of coffee extracts. Whilst others have used metabolomics to elucidate discriminant markers for the authentication of the world's most expensive coffee, Kopi Luwak, an exotic Indonesian blend made from coffee beans that have been eaten by the Asian palm civet (*Paradoxurus hermaphroditus*). Here gas chromatography–mass spectrometry (GC–MS) based multimarker profiling was used to elucidate significant metabolites as markers to determine original, fake Kopi Luwak, regular coffee, and blended samples containing only 50% Kopi Luwak [34]. In another study of a well-known high value food product the evaluation of the adulteration of saffron (*Crocus sativus* L.) by a variety of other cheaper plant bulking agents (i.e. *Crocus sativus* stamens, safflower, turmeric, and gardenia) was undertaken using NMR metabolite fingerprinting [35*]. Herbs and spices can be an expensive commodity and have, along with other sectors, long and complicated supply chains making them vulnerable to adulteration. One very recent study [36*] elegantly demonstrated a two-tier LC–MS-based metabolic profiling and FT-IR metabolic fingerprinting approach, which alarming showed that 24% of the 78 samples of commercially available oregano purchased from retail outlets in the UK and online had some form of adulterant present. These included myrtle, olive, sumac, cistus, and olive leaves, and ranged from 30 to 70% adulteration.

High resolution magic angle spinning nuclear magnetic resonance (HR-MAS-NMR) spectroscopy has also been

used to determine the metabolic profile of a highly prized and designated Sicilian lemon variety (Interdonato Lemon of Messina PGI) in order to determine commercial fraud [37]. With fruit juice authenticity also being the subject of a study by Jandric and co-workers [38]. Where 21 metabolites were selected and found to contribute to the separation of pineapple, orange, grapefruit, apple, clementine, and pomelo juices and their admixtures (pineapple juice adulterated with orange, apple, grapefruit and clementine at 1%, 5%, 10%, 15% adulteration, and orange juice adulterated with apple, grapefruit and clementine at the same levels of adulteration). Additional metabolomics studies published very recently of note also include: a salient reminder of the China melamine [39] crisis and ongoing issues (in some countries) with infant formula, via the classification and evaluation of the degradation and contamination (with melamine) of multiple infant formulas [40]; fully automated NMR analysis of wine authenticity [41] and honeys [42]; and direct analysis in real time mass spectrometry (DART-MS) of poultry meat with metabolomics for the retrospective control of feed fraud via contamination by banned bone meal [43] (yet another reminder of recent history, this time of the BSE crisis). A very recent integrated metabolic profiling and lipidomics approach led to a panel of metabolites for the identification of pork adulteration of minced beef, with additional insights from the analysis of lipids due to the integrated nature of this study [44**]. As with the other omics mentioned all too briefly above, those readers with an interest in metabolomics applied to the detection of food authenticity and integrity [45*] are directed to a number of more comprehensive reviews [33,46,47*,48*,49–51].

Discussion


In this very short overview, it has not been our intention to become entangled in discussions regarding the *nature* of food fraud or criminality, and we will leave the continually evolving and at times unhelpful terminology and quibbling semantics used to describe these various

practices to others. To those in the small, decreasing, and mumpsimus minority and new to this field of study (post Horsegate in 2013), whose comments regarding food fraud/crime include the refrain ‘where is the evidence?’, we kindly remind them that absence of evidence is not evidence of absence, and point them to the international reports, *record seizures* and other practices mentioned in the brief introduction above and accompanying Table 1. Evidence of these practices will not be found, or detected, if it is not being *actively* sought out.

Whilst we have only been able to show a ‘flavour’ of some of the recent applications of omics technologies for the detection of food authenticity and integrity in this article, we hope it is apparent that the omics have much to offer in this area. It must also be pointed out that there can be issues and caveats that are specific to each of the omics fields, such as the measurement issues associated with

molecular and mass spectrometry analysis of food matrices, with a very recent and timely reassessment of their quantitative potential [4]. As food and beverage products already present themselves as complex physical and (bio)chemical matrices, the composition of which can be strongly dependant on a variety of environmental factors such as climate, seasonality, and storage conditions, prior to any other analytical considerations specifically related to the detection of food fraud. Furthermore, one must think more laterally as food fraud involves more complex issues for analytical scientists (and regulators) than the simple adulteration of one high value food product with a less expensive one. Multiple factors can be involved and for example can include issues related to adulterated feed for livestock, and therefore the importance of the detection of chemical residues [52–54] in foods, resulting from either hormonal or antibiotic treatment of livestock, aquatic products [55], or herbicide/

Figure 1




Venn diagram which has been adapted from a graphical model of Routine Activity Theory [64,68] and more recently repurposed for its potential relevance to food fraud by Ellis *et al.* [61**]. This approach is originally based on the three necessary conditions for crime (such as food crime) to occur. The three conditions require: (1) a likely offender (i.e. potential food adulterer/fraudster); (2) a suitable target (i.e. food supply system/network); and finally (3) the absence of a capable guardian (i.e. omics and other disruptive detection technologies). The opportunities/vulnerable areas for crime to occur exist where the so-called capable guardian is absent. We have forwarded a technology-based capable guardian system. Whereby a wide-range of current and future sensor/detection platforms and technologies, along with future predictive computational methods could *together* take on the capable guardian role, and assist in significantly reducing the areas of vulnerability to fraud within food supply chains. Note the area vulnerable to food fraud is dynamic and the detection technologies not static.

pesticide [56] treatment of crops. One recent example from this area is a metabolomics screening method for the adulterated hormonal status of cattle [57]. Other food fraud issues can of course be related directly to consumer safety and health such as the inclusion, whether deliberately or inadvertently, of unwanted contaminants. These can have severe health impacts and include peptides, proteins or a variety of other compounds acting as food allergens [58]. With a recent review discussing the serious issues relating to the ability to measure food allergens reproducibly, their traceability, and the identification of substantial gaps within the international analytical community [59**].

The latter review by Michael Walker of the Government Chemist programme and colleagues from Manchester and Belfast [59**] takes an integrated approach, and whilst each of the main areas of omics briefly discussed here have their own *pros* and *cons*, the integration of several omics methods can be very effective indeed, and lead to more practicable knowledge and insights which can then have the potential for implementation within food supply chains. The study by Trivedi and co-workers [44**] mentioned above provides a very recent example of this integrated omics approach, as have others involving topical food pathogens such as *Campylobacter* [60]. New findings from the omics technologies such as the elucidation of omics markers of authenticity or adulteration can be used in knowledge transfer, and have the potential to be incorporated into a range of commercially available or future technologies for the rapid detection of food fraud. Whether these are to be used in laboratory based detection technologies, wi-fi connected and highly mobile point-and-shoot handheld devices out in supply chains [61**], or at/on/in-line sensors [62,63]. This of course would necessitate co-operation from other disciplines, with the hope that such disruptive detection technologies could act as so-called capable guardians [64] (Figure 1) within food supply chains, with the potential to reduce the areas vulnerable to food fraud. At some point in the future this may also include predictive analysis of points of vulnerability within food networks via one or more omics related techniques from the computational and informatics sciences [65], such as scale-free networks [66] for example. These forms of analyses may have the potential in future to be developed and assist in identifying/predicting nodes which are especially vulnerable to food fraud within complex food supply chains. Allowing for the rapid intervention of disruptive technologies, and/or be aided in this 'identify/predict' function via data automatically collected from omics and related technologies and relayed across networks (Figure 2). Data from these, as well as the other interdisciplinary approaches discussed here, will of course require large-scale and reliable open-access data repositories, and *significantly* more data sharing than is practised to date.

Figure 2



An example of a scale-free supply chain network from a breakfast cereal manufacturer (Chain 12 in Willems [69]) consisting of 88 nodes and 107 edges (links) representing various stages of food production within a company. Edge thickness is proportional to the average demand between nodes. The network is characterised by a number of dense clusters focused on the manufacturing processes linked via transport and procurement nodes.

Such interdisciplinary co-operation, across multiple and at times unrelated disciplines, including engineering, informatics, as well as the social sciences, would require all those involved to see well beyond the boundaries of their own respective fields of research and truly collaborate for the common good. As significant reductions in food fraud will have multiple benefits across international food supply chains. These benefits include *reductions* in: food waste, energy use, greenhouse gas (GHG) emissions, as well as negative health impacts. The integration of multiple disciplines we believe will as a direct consequence lead to *increases* in: food security via the maximisation of product yields, and improvements in food quality, as well as sustainability, with the result that food supply chains would have the potential to be far more resilient to withstand future food shocks [67].

In conclusion, the individual omics discussed here and their related approaches hold a great deal of promise for the detection of food authenticity and integrity, and especially so when using an integrated omics approach (in tandem with future technological and computational advances). With knowledge and expertise from a wide-range of sources leading to valuable new insights and applications; themselves inducing further technological leaps, and reaping beneficial outcomes for an equally wide-range of areas with complex intrinsic and extrinsic links to global food systems.

Acknowledgements

DIE, DPA, and RG wish to thank ESRC and FSA for funding (Food fraud: a supply network integrated systems analysis (Grant number ES/M003183/1)).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Interpol: Largest-ever seizures of fake food and drink in INTERPOL-Europol operation. 2016; vol. 2016.
 2. NFCU.: In *Food Crime Annual Strategic Assessment*. Edited by Unit NFC. Food Standards Agency and Food Standards Scotland; 2016:58.
 3. Balzan J: **Malta at the centre of organic food fraud**. *Malta Today* 2016.
 4. Burns M, Wiseman G, Knight A, Bramley P, Foster L, Rollinson S, Damant A, Primrose S: **Measurement issues associated with quantitative molecular biology analysis of complex food matrices for the detection of food fraud**. *Analyst* 2016, **141**:45-61.
 5. Galimberti A, De Mattia F, Losa A, Bruni I, Federici S, Casiraghi M, Martellos S, Labra M: **DNA barcoding as a new tool for food traceability**. *Food Res Int* 2013, **50**:55-63.
 6. Carvalho DC, Palhares RM, Drummond MG, Frigo TB: **DNA Barcoding identification of commercialized seafood in South Brazil: a governmental regulatory forensic program**. *Food Control* 2015, **50**:784-788.
 7. Kim H, Kumar KS, Hwang SY, Kang BC, Moon HB, Shin KNH: **Utility of stable isotope and cytochrome oxidase I gene sequencing analyses in inferring origin and authentication of hairtail fish and shrimp**. *J Agric Food Chem* 2015, **63**:5548-5556.
 8. Chuah LO, He XB, Effarizah ME, Syahariza ZA, Shamila-Syuhada AK, Rusul G: **Mislabelling of beef and poultry products sold in Malaysia**. *Food Control* 2016, **62**:157-164.
 9. Kane DE, Hellberg RS: **Identification of species in ground meat products sold on the US commercial market using DNA-based methods**. *Food Control* 2016, **59**:158-163.
 10. Nixon GJ, Wilkes TM, Burns MJ: **Development of a real-time PCR approach for the relative quantitation of horse DNA**. *Anal Methods* 2015, **7**:8590-8596.
- A designated hot research article. It being stated that the development of this non-proprietary novel real-time PCR method for the quantitation of horse DNA will inform and strengthen the decision making processes of food companies and regulators.
11. Bloch MS, Paunescu D, Stoessel PR, Mora CA, Stark WJ, Grass RN: **Labeling milk along its production chain with DNA encapsulated in silica**. *J Agric Food Chem* 2014, **62**:10615-10620.
 12. Chen SL, Pang XH, Song JY, Shi LC, Yao H, Han JP, Leon C: **A renaissance in herbal medicine identification: from morphology to DNA**. *Biotechnol Adv* 2014, **32**:1237-1244.
 13. Parvathy VA, Swetha VP, Sheeja TE, Leela NK, Chempakam B, Sasikumar B: **DNA barcoding to detect chilli adulteration in traded black pepper powder**. *Food Biotechnol* 2014, **28**:25-40.
 14. Puddu M, Paunescu D, Stark WJ, Grass RN: **Magnetically recoverable, thermostable, hydrophobic DNA/silica encapsulates and their application as invisible oil tags**. *ACS Nano* 2014, **8**:2677-2685.
 15. Shokralla S, Gibson JF, Nikbakht H, Janzen DH, Hallwachs W, Hajibabaei M: **Next-generation DNA barcoding: using next-generation sequencing to enhance and accelerate DNA barcode capture from single specimens**. *Mol Ecol Res* 2014, **14**:892-901.
 16. Moraes DFC, Still DW, Lum MR, Hirsch AM: **DNA-based authentication of botanicals and plant-derived dietary supplements: where have we been and where are we going?** *Planta Med* 2015, **81**:687-695.
 17. Schneider GF, Dekker C: **DNA sequencing with nanopores**. *Nat Biotechnol* 2012, **30**:326-328.
 18. Claydon AJ, Grundy HH, Charlton AJ, Romero MR: **Identification of novel peptides for horse meat speciation in highly processed foodstuffs**. *Food Addict Contam Part A-Chem Anal Control Exp Risk Assess* 2015, **32**:1718-1729.
 19. Chen YN, Kong DZ, Liu LQ, Song SS, Kuang H, Xu CL: **Development of an enzyme-linked immunosorbent assay (ELISA) for natamycin residues in foods based on a specific monoclonal antibody**. *Anal Methods* 2015, **7**:3559-3565.
 20. Wu XL, Wang WB, Liu LQ, Kuang H, Xu CL: **Monoclonal antibody-based cross-reactive sandwich ELISA for the detection of Salmonella spp. in milk samples**. *Anal Methods* 2015, **7**:9047-9053.
 21. von Bargaen C, Dojahn J, Waidelich D, Humpf HU, Brockmeyer J: **New sensitive high-performance liquid chromatography tandem mass spectrometry method for the detection of horse and pork in halal beef**. *J Agric Food Chem* 2013, **61**:11986-11994.
 22. Ohana D, Dalebout H, Marissen RJ, Wulff T, Bergquist J, Deelder AM, Palmblad M: **Identification of meat products by shotgun spectral matching**. *Food Chem* 2016, **203**:28-34.
 23. Orduna AR, Husby E, Yang CT, Ghosh D, Beaudry F: **Assessment of meat authenticity using bioinformatics, targeted peptide biomarkers and high-resolution mass spectrometry**. *Food Addict Contam Part A-Chem Anal Control Exp Risk Assess* 2015, **32**:1709-1717.
 24. Sassi M, Arena S, Scaloni A: **MALDI-TOF-MS platform for integrated proteomic and peptidomic profiling of milk samples allows rapid detection of food adulterations**. *J Agric Food Chem* 2015, **63**:6157-6171.
 25. Agrawal GK, Timperio AM, Zolla L, Bansal V, Shukla R, Rakwal R: **Biomarker discovery and applications for foods and beverages: proteomics to nanoproteomics**. *J Proteomics* 2013, **93**:74-92.
 26. Colgrave ML, Goswami H, Howitt CA, Tanner GJ: **Proteomics as a tool to understand the complexity of beer**. *Food Res Int* 2013, **54**:1001-1012.
 27. Fasoli E, Righetti PG: **Proteomics of fruits and beverages**. *Curr Opin Food Sci* 2015, **4**:76-85.
 28. Paredi G, Sentandreu MA, Mozzarelli A, Fadda S, Hollung K, de Almeida AM: **Muscle and meat: new horizons and applications for proteomics on a farm to fork perspective**. *J Proteomics* 2013, **88**:58-82.
 29. Herrero M, Simo C, Garcia-Canas V, Ibanez E, Cifuentes A: **Foodomics: MS-based strategies in modern food science and nutrition**. *Mass Spectrom Rev* 2012, **31**:49-69.
 30. Goodacre R, Vaidyanathan S, Bianchi G, Kell DB: **Metabolic profiling using direct infusion electrospray ionisation mass spectrometry for the characterisation of olive oils**. *Analyst* 2002, **127**:1457-1462.
 31. Vidal NP, Goicoechea E, Manzanos MJ, Guillen MD: **Fourier transform infrared spectroscopy as a tool to study farmed and wild sea bass lipid composition**. *J Sci Food Agric* 2014, **94**:1340-1348.
 32. Arana VA, Medina J, Alarcon R, Moreno E, Heintz L, Schafer H, Wist J: **Coffee's country of origin determined by NMR: the Colombian case**. *Food Chem* 2015, **175**:500-506.
 33. Ellis DI, Brewster VL, Dunn WB, Allwood JW, Golovanov AP, Goodacre R: **Fingerprinting food: current technologies for the detection of food adulteration and contamination**. *Chem Soc Rev* 2012, **41**:5706-5727.
 34. Jumhawan U, Putri SP, Yusianto, Marwani E, Bamba T, Fukusaki E: **Selection of discriminant markers for authentication of asian palm civet coffee (Kopi Luwak): a metabolomics approach**. *J Agric Food Chem* 2013, **61**:7994-8001.

35. Petrakis EA, Cagliani LR, Polissiou MG, Consonni R: **Evaluation of saffron (*Crocus sativus* L.) adulteration with plant adulterants by H-1 NMR metabolite fingerprinting.** *Food Chem* 2015, **173**:890-896.

A well cited NMR metabolite fingerprinting study on the adulteration of the highly prized and expensive saffron with other plant adulterants and bulking agents. Chemometric modelling approaches were used to authenticate and predict saffron adulteration which was said to compare well to the stated deficiencies in currently established methodologies.

36. Black C, Haughey SA, Chevallier OP, Galvin-King P, Elliott CT: **A comprehensive strategy to detect the fraudulent adulteration of herbs: the oregano approach.** *Food Chem* 2016, **210**:551-557.

An article which received a great deal of media attention even prior to its official publication. Using a metabolic profiling and metabolic fingerprinting approach, it alarmingly showed that 24% of the oregano samples analysed had some form of adulterant present, ranging from 30 to 70% adulteration.

37. Cicero N, Corsaro C, Salvo A, Vasi S, Giofre SV, Ferrantelli V, Di Stefano V, Mallamace D, Dugo G: **The metabolic profile of lemon juice by proton HR-MAS NMR: the case of the PGI Interdonato Lemon of Messina.** *Nat Prod Res* 2015, **29**:1894-1902.

38. Jandric Z, Roberts D, Rathor MN, Abraham A, Islam M, Cannavan A: **Assessment of fruit juice authenticity using UPLC-QToF MS: a metabolomics approach.** *Food Chem* 2014, **148**:7-17.

39. Lang TT, Pang S, He LL: **Integration of colorimetric and SERS detection for rapid screening and validation of melamine in milk.** *Anal Methods* 2015, **7**:6426-6431.

40. Inoue K, Tanada C, Sakamoto T, Tsutsui H, Akiba T, Min JZ, Todoroki K, Yamano Y, Toyooka T: **Metabolomics approach of infant formula for the evaluation of contamination and degradation using hydrophilic interaction liquid chromatography coupled with mass spectrometry.** *Food Chem* 2015, **181**:318-324.

41. Spraul M, Link M, Schaefer H, Fang F, Schuetz B: **Wine analysis to check quality and authenticity by fully-automated H-1-NMR.** In *38th World Congress of Vine and Wine*. Edited by JeanMarie A. E D P Sciences; 2015. [BIO Web of Conferences, vol 5].

42. Jamroz MK, Paradowska K, Zawada K, Makarova K, Kazmierski S, Wawer I: **H-1 and C-13 NMR-based sugar profiling with chemometric analysis and antioxidant activity of herb honeys and honeys.** *J Sci Food Agric* 2014, **94**:246-255.

43. Cajka T, Danhelova H, Zachariasova M, Riddellova K, Hajslova J: **Application of direct analysis in real time ionization-mass spectrometry (DART-MS) in chicken meat metabolomics aiming at the retrospective control of feed fraud.** *Metabolomics* 2013, **9**:545-557.

44. Trivedi DK, Hollywood KA, Rattray NJW, Ward H, Greenwood J, Ellis DI, Goodacre R: **Meat, the metabolites: an integrated metabolite profiling and lipidomics approach for the detection of the adulteration of beef with pork.** *Analyst* 2016, **141**:2155-2164.

A designated hot original research article which uses an integrated omics approach for the detection of beef adulterated with pork. Which not only demonstrated an excellent example of the integrated omics approach but also provided a panel of metabolites specific to this form of adulteration and interesting additional information on lipids.

45. Ellis DI, Goodacre R: **Detecting food authenticity and integrity.** *Anal Methods* 2016, **8**:3281-3283.

Could be considered as a useful fingerpost, pointing to most, though not all, of the articles and reviews within a very recent Royal Society of Chemistry themed collection on detecting food authenticity and integrity.

46. Cozzolino D, Porker K, Laws M: **An overview on the use of infrared sensors for in field, proximal and at harvest monitoring of cereal crops.** *Agric Basel* 2015, **5**:713-722.

47. Cubero-Leon E, Penalver R, Maquet A: **Review on metabolomics for food authentication.** *Food Res Int* 2014, **60**:95-107.

Another very well received review which describes metabolomics workflows for food authentication, recent applications of metabolomics methods, and again, their potential advantages and limitations.

48. Do TKT, Hadji-Minaglou F, Antoniotti S, Fernandez X: **Authenticity of essential oils.** *Trends Anal Chem* 2015, **66**:146-157.

A comprehensive overview of known adulterants and analytical methods for detecting adulterants in essential oils. Includes an overview of

fundamental principles and applications of analytical techniques as well as a comparison of their advantages and limitations.

49. Ibanez C, Garcia-Canas V, Valdes A, Simo C: **Novel MS-based approaches and applications in food metabolomics.** *Trends Anal Chem* 2013, **52**:100-111.

50. Senyuva HZ, Gokmen V, Sarikaya EA: **Future perspectives in Orbitrap (TM)-high-resolution mass spectrometry in food analysis: a review.** *Food Addit Contam Part A-Chem Anal Control Exp Risk Assess* 2015, **32**:1568-1606.

51. Trimigno A, Marincola FC, Dellarosa N, Picone G, Laghi L: **Definition of food quality by NMR-based foodomics.** *Curr Opin Food Sci* 2015, **4**:99-104.

52. Li ZZ, Li ZL, Li DM, Gao HL, Chen XJ, Cao L, Hou YZ, Li SB: **Molecularly imprinted polymer-based chemiluminescence imaging assay for the determination of ethopabate residues in chicken muscle.** *Anal Methods* 2015, **7**:9295-9303.

53. Ucles A, Garcia AV, Garcia MDG, del Real AMA, Fernandez-Alba AR: **Benzimidazole and imidazole fungicide analysis in grape and wine samples using a competitive enzyme-linked immunosorbent assay.** *Anal Methods* 2015, **7**:9158-9165.

54. Wang Q, Zheng YJ, Zhang XL, Han XX, Wang T, Zhang ZP: **A silica coated paper substrate: development and its application in paper spray mass spectrometry for rapid analysis of pesticides in milk.** *Analyst* 2015, **140**:8048-8056.

55. Uchida K, Konishi Y, Harada K, Okihashi M, Yamaguchi T, Do MHN, Bui LT, Nguyen TD, Khong NDT, Tran HT *et al.*: **Monitoring of antibiotic residues in aquatic products in urban and rural areas of Vietnam.** *J Agric Food Chem* 2016 <http://dx.doi.org/10.1021/acs.jafc.6b00091>. in press.

56. Hou RY, Pang S, He LL: **In situ SERS detection of multi-class insecticides on plant surfaces.** *Anal Methods* 2015, **7**:6325-6330.

57. Kouassi Nzougnet J, Dervilly-Pinel G, Chereau S, Biancotto G, Monteau F, Elliott C, Le Bizec B: **First insights into serum metabolomics of trenbolone/estradiol implanted bovines; screening model to predict hormone-treated and control animals' status.** *Metabolomics* 2015, **11**:1184-1196.

58. Sayers RL, Johnson PE, Marsh JT, Barran P, Brown H, Mills ENC: **The effect of thermal processing on the behaviour of peanut allergen peptide targets used in multiple reaction monitoring mass spectrometry experiments.** *Analyst* 2016, **141**:4130-4141.

59. Walker MJ, Burns DT, Elliott CT, Gowland MH, Mills ENC: **Is food allergen analysis flawed? Health and supply chain risks and a proposed framework to address urgent analytical needs.** *Analyst* 2016, **141**:24-35.

A timely and important article by a range of recognised food allergy experts discussing the urgent and serious issues relating to the ability to measure food allergens reproducibly, their traceability, and the identification of substantial gaps within the international analytical community.

60. Muhamadali H, Weaver D, Subaihi A, AlMasoud N, Trivedi DK, Ellis DI, Linton D, Goodacre R: **Chicken, beams, and Campylobacter: rapid differentiation of foodborne bacteria via vibrational spectroscopy and MALDI-mass spectrometry.** *Analyst* 2016, **141**:111-122.

61. Ellis DI, Muhamadali H, Haughey SA, Elliott CT, Goodacre R: **Point-and-shoot: rapid quantitative detection methods for on-site food fraud analysis — moving out of the laboratory and into the food supply chain.** *Anal Methods* 2015, **7**:9401-9414.

A well received and designated hot article discussing a range of spectroscopy/spectroscopy based approaches, as well as other methods currently under development. Forwards a future perspective of how this range of detection methods, along with developments in predictive computational and information sciences, could reduce the areas of vulnerability to food crime within food supply chains.

62. Sharma TK, Ramanathan R, Rakwal R, Agrawal GK, Bansal V: **Moving forward in plant food safety and security through NanoBioSensors: adopt or adapt biomedical technologies?** *Proteomics* 2015, **15**:1680-1692.

63. Zou XB, Huang XW, Povey M: **Non-invasive sensing for food reassurance.** *Analyst* 2016, **141**:1587-1610.

64. Hollis-Peel ME, Welsh BC: **What makes a guardian capable?. A test of guardianship in action.** *Security J* 2014, **27**:320-337.
65. Gubbi J, Buyya R, Marusic S, Palaniswami M: **Internet of Things (IoT): a vision, architectural elements, and future directions.** *Fut Gen Comput Syst – Int J Grid Comput Esci* 2013, **29**:1645-1660.
66. Barabasi AL: **Scale-free networks: a decade and beyond.** *Science* 2009, **325**:412-413.
67. Macfadyen S, Tylanakis JM, Letourneau DK, Benton TG, Tittone P, Perring MP, Gomez-Creutzberg C, Baldi A, Holland JM, Broadhurst L *et al.*: **The role of food retailers in improving resilience in global food supply.** *Global Food Security-Agric Policy Econ Environ* 2015, **7**:1-8.
68. Cohen LE, Felson M: **Social change and crime rate trends – a routine activity approach.** *Am Soc Rev* 1979, **44**:588-608.
69. Willems SP: **Real-world multiechelon supply chains used for inventory optimization.** *Manufact Serv Oper Manage* 2008, **10**:19-23.